

South Dakota

An Annual Report on Cancer Incidence and Mortality Among South Dakotans, 2014



Cancer In South Dakota 2014



Preface

“Cancer in South Dakota 2014” is the 22nd annual report from the South Dakota Cancer Registry (SDCR) in the Office of Chronic Disease Prevention and Health Promotion in the Division of Family and Community Health within the South Dakota Department of Health (DOH). The report contains 2014 cancer incidence and mortality data of South Dakota residents.

Acknowledgements

The South Dakota Cancer Registry acknowledges all the Certified Tumor Registrars in hospital cancer registries who work diligently to maintain quality data, hospitals, clinics, physicians, pathology laboratories and other providers and health entities that submit data to the SDCR, and co-workers in the Department of Health.

Funding Source

This publication was funded by the South Dakota Cancer Registry with a grant from the Centers for Disease Control and Prevention – grant number U58/DP003943.

Changes in Report

Age-adjusted rates were calculated using the 2000 US standard million and the US Census Bureau 2002 - 2014 population estimates for South Dakota. Reports published before 2012 used the 2000 South Dakota census population for age-adjusted calculations and are not comparable to this report.

Requests for Information

Data queries, cancer concerns, information, or requests for additional copies (if available) should be addressed to:

Kay Dosch, CTR, Coordinator
South Dakota Cancer Registry
615 East 4th Street
Pierre, SD 57501-1700
Phone: (605) 773-6345

Suggested Citation

Cancer in South Dakota, 2014. Department of Health, Pierre, SD. May 2017. All material in this report may be reproduced or copied without permission. However, citation as to source is appreciated.

Web Access: This report is available on the internet at the following URL:
<http://getscreened.sd.gov/documents/Cancer2014.pdf>

The following Department of Health personnel were the main contributors to the development of this report:

Mary Sarvis	SDCR Data Manager
Barb Buhler	Public Information Officer
Karen Cudmore	Cancer Programs Director
Kay Dosch	SDCR Coordinator
Mark Gildemaster	Management Analyst
Katie Hill	Communications and Community Coordinator
Ashley Miller	Epidemiologist

TABLE OF CONTENTS

PREFACE	i
TABLE OF CONTENTS	iii
LIST OF TABLES	iv
LIST OF FIGURES	v
I. EXECUTIVE SUMMARY	1
II. INTRODUCTION	3
III. TECHNICAL NOTES	5
IV. CANCER INCIDENCE	8
V. CANCER CASES AND DEATHS BY RANK	13
VI. STAGE AT DIAGNOSIS	16
VII. CANCER MORTALITY	18
VIII. YEARS OF POTENTIAL LIFE LOST	23
IX. AVERAGE YEARS OF LIFE LOST	25
X. SELECTED CANCER SITES INCIDENCE AND MORTALITY	27
Bladder	28
Breast (Female)	30
Cervix Uteri	32
Colorectal	34
Corpus and Uterus	36
Kidney and Renal Pelvis	38
Leukemia	40
Lung and Bronchus	42
Melanoma (Skin)	44
Myeloma	46
Non-Hodgkin's Lymphoma	48
Ovary	50
Pancreas	52
Prostate	54
Stomach	56
Thyroid	58
XI. APPENDICES	
Appendix A: 2000 US Standard Million Population	60
Appendix B: 2005-2014 South Dakota Estimated Population	60
Appendix C: Race in South Dakota by County, 2014 Estimated Population	61
Appendix D: SEER Incidence Site Analysis Categories	62-64
Appendix E: SEER Cancer Cause of Death Analysis Categories	65-66
REFERENCES	67

LIST OF TABLES

Table 1: Cancer Cases and Incidence Rates by County, South Dakota, 2014 and 2010-2014 Average.....	8
Table 2: Age-adjusted Incidence Rates by County for Selected Sites, 2014.....	9
Table 3: Age-adjusted Incidence Rates by Site, Gender, and Race, South Dakota, 2014.....	10
Table 4: Percentage of Selected Cancers by Age Groups in South Dakota, 2014.....	12
Table 5: South Dakota Stage at Diagnosis, All Cases, 2014.....	16
Table 6: Stage at Diagnosis for Selected Sites by Race, South Dakota, 2014.....	17
Table 7: Cancer Mortality and Rates by County, South Dakota, 2014 and 2010-2014	18
Table 8: Age-adjusted Mortality Rates by County for Selected Sites, South Dakota, 2014....	19
Table 9: Age-adjusted Mortality Rates by Site, Gender, and Race, South Dakota, 2014.....	20
Table 10: Percentage of Cancer Deaths by Age Groups and Selected Primary Sites, South Dakota, 2014	22
Table 11: Leading Causes of Years of Potential Life Lost (to age 75), South Dakota, 2014....	23
Table 12: Bladder Incidence and Mortality Summary, 2014.....	28
Table 13: Breast (Female) Incidence and Mortality Summary, 2014.....	30
Table 14: Cervix Uteri Incidence and Mortality Summary, 2014.....	32
Table 15: Colorectal Incidence and Mortality Summary, 2014	34
Table 16: Corpus and Uterus, NOS Incidence and Mortality Summary, 2014.....	36
Table 17: Kidney and Renal Pelvis Incidence and Mortality Summary, 2014.....	38
Table 18: Leukemia Incidence and Mortality Summary, 2014	40
Table 19: Lung and Bronchus Incidence and Mortality Summary, 2014	42
Table 20: Melanoma Incidence and Mortality Summary, 2014	44
Table 21: Myeloma Incidence and Mortality Summary, 2014	46
Table 22: Non-Hodgkin’s Lymphoma Incidence and Mortality Summary, 2014	48
Table 23: Ovarian Incidence and Mortality Summary, 2014	50
Table 24: Pancreas Incidence and Mortality Summary, 2014	52
Table 25: Prostate Incidence and Mortality Summary, 2014.....	54
Table 26: Stomach Incidence and Mortality Summary, 2014	56
Table 27: Thyroid Incidence and Mortality Summary, 2014	58

LIST OF FIGURES

Figure 1:	All Sites Cancer Incidence Rates by Race and Gender, South Dakota, 2014.....	12
Figure 2:	Percent Distribution of Cancer Cases and Deaths by Rank and Gender, South Dakota, 2014	13
Figure 3:	Cancer Incidence Rates by County, South Dakota, 2014 (Map)	14
Figure 4:	Cancer Mortality Rates by County, South Dakota, 2014 (Map)	15
Figure 5:	Stage at Diagnosis, All Sites, South Dakota, 2014.....	16
Figure 6:	Stage at Diagnosis in South Dakota, White, 2014.....	17
Figure 7:	Stage at Diagnosis in South Dakota, American Indian, 2014	17
Figure 8:	All Sites Cancer Mortality Rates by Race and Gender, South Dakota, 2014.....	22
Figure 9:	Years of Potential Life Lost (YPLL) Before Age 75 for Leading Causes of Death, by Race, South Dakota, 2014.....	23
Figure 10:	Years of Potential Life Lost (YPLL) Before Age 75 for Leading Causes of Death, by Race, South Dakota, 2010-2014	23
Figure 11:	YPLL for Selected Cancers Among Whites, South Dakota, 2014	24
Figure 12:	YPLL for Selected Cancers Among American Indians, South Dakota, 2014.....	24
Figure 13:	AYLL for Selected Cancers Among Whites, South Dakota, 2014	25
Figure 14:	AYLL for Selected Cancers Among American Indians, South Dakota, 2014.....	25
Figure 15:	Bladder Cancer Stage at Diagnosis, South Dakota, 2014.....	28
Figure 16:	Bladder Cancer Number of Cases and Deaths by Age, South Dakota, 2014.....	29
Figure 17:	Bladder Cancer Cases and Deaths by Year, South Dakota, 2001-2014	29
Figure 18:	Bladder Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014.....	29
Figure 19:	Female Breast Cancer Stage at Diagnosis, South Dakota, 2014	30
Figure 20:	Female Breast Cancer Number of Cases and Deaths by Age, South Dakota, 2014	31
Figure 21:	Female Breast Cancer Cases and Deaths by Year, South Dakota, 2001-2014	31
Figure 22:	Female Breast Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014.....	31
Figure 23:	Cervix Uteri Cancer Stage at Diagnosis, South Dakota, 2014.....	32
Figure 24:	Cervix Uteri Cancer Number of Cases and Deaths by Age, South Dakota, 2014 .	33
Figure 25:	Cervix Uteri Cancer Cases and Deaths by Year, South Dakota, 2001-2014.....	33
Figure 26:	Cervix Uteri Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014.....	33
Figure 27:	Colorectal Cancer Stage at Diagnosis, South Dakota, 2014	34
Figure 28:	Colorectal Cancer Number of Cases and Deaths by Age, South Dakota, 2014....	35
Figure 29:	Colorectal Cancer Cases and Deaths by Year, South Dakota, 2001-2014	35
Figure 30:	Colorectal Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014.....	35
Figure 31:	Corpus and Uterus, NOS Cancer Stage at Diagnosis, South Dakota, 2014.....	36
Figure 32:	Corpus and Uterus, NOS Cancer Number of Cases and Deaths by Age, South Dakota, 2014	37
Figure 33:	Corpus and Uterus, NOS Cancer Cases and Deaths by Year, South Dakota, 2001-2014	37
Figure 34:	Corpus and Uterus, NOS Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014	37
Figure 35:	Kidney and Renal Pelvis Cancer Stage at Diagnosis, South Dakota, 2014	38
Figure 36:	Kidney and Renal Pelvis Cancer Number of Cases and Deaths by Age, South Dakota, 2014.....	39

LIST OF FIGURES (cont'd)

Figure 37: Kidney and Renal Pelvis Cancer Cases and Deaths by Year, South Dakota, 2001-2014	39
Figure 38: Kidney and Renal Pelvis Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014	39
Figure 39: Leukemia Number of Cases and Deaths by Age, South Dakota, 2014	41
Figure 40: Leukemia Cases and Deaths by Year, South Dakota, 2001-2014	41
Figure 41: Leukemia Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014	41
Figure 42: Lung and Bronchus Cancer Stage at Diagnosis, South Dakota, 2014	42
Figure 43: Lung and Bronchus Cancer Number of Cases and Deaths by Age, South Dakota, 2014	43
Figure 44: Lung and Bronchus Cancer Cases and Deaths by Year, South Dakota, 2001-2014	43
Figure 45: Lung and Bronchus Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014.....	43
Figure 46: Melanoma of the Skin Stage at Diagnosis, South Dakota, 2014	44
Figure 47: Melanoma of the Skin Number of Cases and Deaths by Age, South Dakota, 2014	45
Figure 48: Melanoma of the Skin Cases and Deaths by Year, South Dakota, 2001-2014	45
Figure 49: Melanoma of the Skin Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014.....	45
Figure 50: Myeloma Number of Cases and Deaths by Age, South Dakota, 2014	47
Figure 51: Myeloma Cases and Deaths by Year, South Dakota, 2001-2014	47
Figure 52: Myeloma Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014	47
Figure 53: Non-Hodgkin's Lymphoma Stage at Diagnosis, South Dakota, 2014.....	48
Figure 54: Non-Hodgkin's Lymphoma Number of Cases and Deaths by Age, South Dakota, 2014	49
Figure 55: Non-Hodgkin's Lymphoma Cases and Deaths by Year, South Dakota, 2001-2014	49
Figure 56: Non-Hodgkin's Lymphoma Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014.....	49
Figure 57: Ovarian Cancer Stage at Diagnosis, South Dakota, 2014.....	50
Figure 58: Ovarian Cancer Number of Cases and Deaths by Age, South Dakota, 2014	51
Figure 59: Ovarian Cancer Cases and Deaths by Year, South Dakota, 2001-2014.....	51
Figure 60: Ovarian Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014.....	51
Figure 61: Pancreatic Cancer Stage at Diagnosis, South Dakota, 2014	52
Figure 62: Pancreatic Cancer Number of Cases and Deaths by Age, South Dakota, 2014 ...	53
Figure 63: Pancreatic Cancer Cases and Deaths by Year, South Dakota, 2001-2014.....	53
Figure 64: Pancreatic Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014.....	53
Figure 65: Prostate Cancer Stage at Diagnosis, South Dakota, 2014.....	54
Figure 66: Prostate Cancer Number of Cases and Deaths by Age, South Dakota, 2014.....	55
Figure 67: Prostate Cancer Cases and Deaths by Year, South Dakota, 2001-2014	55

LIST OF FIGURES (cont'd)

Figure 68: Prostate Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014.....	55
Figure 69: Stomach Cancer Stage at Diagnosis, South Dakota, 2014	56
Figure 70: Stomach Cancer Number of Cases and Deaths by Age, South Dakota, 2014.....	57
Figure 71: Stomach Cancer Cases and Deaths by Year, South Dakota, 2001-2014	57
Figure 72: Stomach Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014.....	57
Figure 73: Thyroid Cancer Stage at Diagnosis, South Dakota, 2014	58
Figure 74: Thyroid Cancer Number of Cases and Deaths by Age, South Dakota, 2014	59
Figure 75: Thyroid Cancer Cases and Deaths by Year, South Dakota, 2001-2014	59
Figure 76: Thyroid Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2014.....	59

I. EXECUTIVE SUMMARY

This report summarizes the burden of cancer in South Dakota and includes cancer incidence and mortality data. The data will enable the many organizations working with cancer prevention and control to identify public health problems, target goals for cancer control, and to inform citizens and health care professionals about risks, early detection, and treatment.

Incidence 2014

- 4,498 South Dakotans were diagnosed with invasive, reportable cases of cancer, which excludes the less life-threatening cancers such as *in situ* cancers (except *in situ* bladder cases) and the common skin cancers.
- Each day 12 cases of cancer were diagnosed in residents of South Dakota; this includes only cases of invasive cancer and *in situ* bladder.
- The five most diagnosed cancer sites (female breast, lung, prostate, colorectal, melanoma) accounted for 54% of all cancer cases.
- Female breast cancer was the most common reportable malignancy with 608 cases among women, accounting for 13.5% of all cases and 29% of cases for women.
- Lung cancer was the second most common reportable malignancy with 582 cases, 12.9% of all cases.
- Prostate cancer was the third most common reportable cancer with 549 cases, 12.2% of all cases and 22.9% of cases for men.
- Colon and rectal cancers were the fourth most common malignancy with 447 cases, 10% of all cases.
- Melanoma cancers were the fifth most common malignancy with 203 cases, 5.3% of all reported cases.
- 53% of all new cancers were diagnosed in males and 47% were in females.
- Males had an age-adjusted incidence rate of 505.7 per 100,000, which was higher than females who had an age-adjusted rate of 413.4 per 100,000.
- Whites accounted for 93.4% of cancer cases with 4,202 cases whereas American Indians were 5.4% with 245 cases.
- The American Indian age-adjusted incidence rate was 524.6, which is higher than the age-adjusted rate among whites of 451.5.
- The South Dakota age-adjusted incidence rate for 2014 was 451.5, significantly higher than the US SEER 2014 age-adjusted incidence rate of 428.6 per 100,000 persons.

Mortality 2014

- Overall, cancer was the second leading cause of death in South Dakota.
- In 2014, 1,679 South Dakotans died from cancer, accounting for one in every four deaths in South Dakota.
- Each day four South Dakotans died from cancer.
- The five cancer sites (lung, colorectal, pancreas, female breast, leukemia) caused 55.7% of all cancer deaths.
- Lung and bronchus cancers were the leading cause of cancer deaths at 439 deaths or 26.1% of all cancer deaths. Lung and bronchus cancers caused one in four deaths due to cancer.
- Colorectal cancer was the second leading cause of cancer deaths with 188 deaths, 11.2% of all cancer deaths.
- Pancreatic cancer was the third leading cause of death with 118 deaths, 7.0% of all cancer deaths.
- Female breast cancer was the fourth leading cause of cancer deaths with 100 deaths, 6.0% of all cancer deaths and 13.5% of all female cancer deaths.
- Leukemia cancer was the fifth leading cause of death with 90 deaths, 5.4% of all cancer deaths.
- Over half, 56% of all cancer deaths were males and 44% were females.
- Males had an age-adjusted death rate of 201.2 per 100,000 males, 52.4% higher than females with an age-adjusted rate of 132.0 deaths per 100,000 females.
- Whites accounted for 94.2% of deaths with 1,581 deaths, whereas American Indians were 5.2% with 88 deaths.
- The American Indian age-adjusted death rate was 217.9 which is 35.6% higher than the rate among whites at the age-adjusted death rate of 160.7 and is also significantly higher.
- South Dakota's age-adjusted death rate for 2014 was 161.4, no significant difference than the US SEER 2014 rate of 161.3.

Trends

- Melanoma incidence cases have increased significantly since 2009.
- For the last decade, female breast cancer mortality rates have remained steady.

II. INTRODUCTION

A limited cancer data collection system was established in 1992 under South Dakota Codified Laws, SDCL 1-43-1 to 1-43-18, and Administrative Rules ARSD 44:22:01 to 44:22:05, inclusive. The South Dakota Cancer Registry (SDCR) was established in 2001 to develop a statewide, population-based cancer surveillance system. However, the state legislature amended the law to expand reporting to reflect statewide surveillance. SDCL 1-43-14 has been in effect since July 1, 2005:

Any hospital licensed pursuant to chapter 34-12, physician licensed pursuant to chapter 36-4, physician assistant licensed pursuant to chapter 36-4A, nurse practitioner or nurse midwife licensed pursuant to chapter 36-9A, pathology laboratory, or free-standing radiology center that detects, diagnoses, or treats a cancer case in South Dakota shall submit a report to the Department of Health as required by § 1-43-11 to 1-43-17, inclusive.

Reportable cancers for 2014 include benign brain, benign central nervous system and all malignant neoplasms except basal and squamous cell carcinomas of the skin and *in situ* cervical cancers. In addition to receiving cases from hospital cancer registries the SDCR actively follows back pathology reports and abstracts cases from facilities without tumor registries.

The SDCR performs many quality assurance procedures to assure that the data are valid. The data are run through numerous edits and consolidated if received from more than one reporting source. In addition, the SDCR links the incidence data with mortality files to identify persons whose death records show cancer as a cause of death, but these cancers were not reported to the central registry. The SDCR also links the incidence file with the Indian Health Service database to identify any American Indian South Dakotan who was misclassified as another race.

The SDCR uses the cancer incidence data reported as well as the mortality data and health behavior surveys collected by the department's Health Statistics Office and Vital Records to provide useful information for cancer control and prevention programs, researchers, clinicians, and policy makers. The SDCR is able to answer several epidemiological questions such as:

- How many South Dakotans are diagnosed or die from cancer each year?
- What are the most common cancers?
- When are cancers being diagnosed, i.e. at what stage?
- Which cancers are the deadliest?
- Who is most affected by cancer?
- What are the trends in cancer incidence and mortality?
- Where are cancers occurring?
- Where and what are the disparities?
- Are screening efforts working?

Every life is touched by cancer in some way whether one is diagnosed with the disease or has a family member or friend with the disease. Although cancer is primarily a disease of people over 50 years old, the younger a person dies from cancer, the greater the impact on societal and economic costs. Cancer concerns voiced by South Dakotans are a priority for the SDCR.

As the SDCR continues to collect population based data and as more health care entities and providers report cases, more questions can be answered with the data. For example, SDCR can look at modifiable risk factors such as obesity and exercise. The Harvard Report on Cancer Prevention in 1996 researched the risk factors for cancer. The estimated percentages (which are still relevant today) of total cancer deaths attributed to established causes of cancer were:

Risk Factor	Percentage
Tobacco	30%
Adult diet/obesity	30%
Sedentary lifestyle	5%
Occupational factors	5%
Family history of cancer	5%
Viruses/biological agents	5%
Perinatal factors/growth	5%
Reproductive factors	3%
Alcohol	3%
Socioeconomic status	3%
Environmental pollution	2%
Ionizing /UV radiation	2%
Prescription drugs/medical	1%
Salt/food additives/contaminants	1%

The most successful way to prevent cancer is to limit the number of modifiable risk factors by following these guidelines:

- Do not smoke
- Maintain a healthy weight
- Get at least 30 minutes of physical activity every day
- Eat a healthy diet
- Limit alcohol intake to less than one drink a day
- Protect yourself from the sun

Some risk factors for cancer can be avoided, but many cannot. For example, both smoking and inheriting certain genes are risk factors for some types of cancer, but only smoking can be avoided. Risk factors that a person can control are called modifiable risk factors.

Source: National Cancer Institute

The SDCR continuously expands data collection. This will help prevention and control programs to target at risk populations as well as support epidemiologic studies. The end goal is to reduce the cancer burden by producing valid and accurate data reflecting the complete assessment of cancer in South Dakota, and to disseminate the information in a timely manner.

III. TECHNICAL NOTES

Age-adjusted death rates: Death rates are calculated for total cases and separately for males and females. The death rates are age-adjusted to the 2000 US standard population using five-year groups and are per 100,000 persons. Rates are presented for 2014 and for the five-year period, 2010-2014.

Age-adjusted incidence rate: Age-adjusted incidence rates are calculated using the direct method and standardized to the age distribution of the 2000 US standard population (Appendix A). Age adjustment allows rates for one geographic area to be compared with rates from other geographic areas that may have differences in age distributions. Any observed differences in age-adjusted incidence rates between populations are not due to different age structures. Reports prior to 1999 used the 1970 US standard population. In conformity with the National Cancer Institute's (NCI) Surveillance, Epidemiology, and End Results (SEER) Program guidelines, the incidence rates for cancer sites exclude the following:

- *In situ* cases, except bladder
- Basal and squamous cell skin cancers
- Cases with unknown age
- Cases with unknown gender

Age-specific incidence rates: Age specific rates are calculated by dividing the number of cases for a given age group by the total population of that age group and are expressed as an average annual rate per 100,000 persons by age group. Age specific rates exclude the same types of cases that are excluded from age-adjusted incidence rates. These rates, however, are crude rates, i.e. not age-adjusted.

Annual percent change (APC): The annual percent change is the average rate of change in a cancer rate per year in a given time frame indicating how fast or how slowly a cancer rate has increased or decreased each year over a period of years. A negative APC describes a decreasing trend, and a positive APC describes an increasing trend.

Average years of life lost (AYLL): This is the extent to which life is cut short due to premature death. This is obtained by dividing the years of potential life lost (YPLL) by the number of deaths. On average each person who dies from cancer loses 15 years of their life.

Cancer case definitions: A "cancer case" is defined as the primary cancer site, i.e., the site where the cancer started. Since an individual can have more than one primary cancer site, the number of incident cancer cases could be greater than the number of persons who are diagnosed with cancer. A *metastasis* is not a primary site.

Changes in diagnostic criteria: Early detection resulting from either screening or early response to symptoms may result in increasing diagnosis of small tumors that are not yet life-threatening. This may raise incidence and survival rates but without changes in mortality rates. Cancers likely to be affected are breast, colon, cervix uteri, prostate, and melanoma. Prostate cancer is particularly prone to changing diagnostic criteria.

Confidence intervals (CI): A confidence interval tells how confident we are of the accuracy of the calculated rates. The SDCR uses a computed interval with a given probability of 95%, i.e., the true value of the calculated rate is contained within the interval. Thus, given a calculated rate of 191.4 and a confidence interval of 182.1 to 200.8, it is better to say that the true rate will fall between 182.1 and 200.8. The larger the sample size, the shorter the interval size, giving us more certainty that the rate is correct. When CI for percentages contains zero, the rate is considered to be stable. Above zero, the statistical significance is higher and below zero it is lower.

Data source: All data, tables, and figures come from the South Dakota Department of Health, *American Cancer Society Facts and Figures 2014* or *SEER Cancer Statistics Review 1975-2013* and should be cited as such if taken out of this report in part. SEER data represents approximately 10% of the U.S population.

Disparity: Health disparities are differences in the incidence, prevalence, mortality, and burden of diseases and other adverse health conditions that exist among specific population groups in the United States.¹ Health disparities can be defined as a specific group bearing a disproportionate share of negative health outcomes compared to the general population, i.e., disease, disability, and death.² Disparity can occur as a result of factors such as poverty, living in geographically underserved areas and belonging to specific minority groups.

Early detection/screening: Improved early detection/screening may produce increases in both incidence and survival rates. Increases may occur as a result of the introduction of new procedures. The interval between the time a cancer is diagnosed by a screening procedure and the time when it would have been diagnosed in the absence of screening procedures is called the lead-time. Changes in lead-time, for example, in breast cancer diagnosis, have led to increased survival rates and reduction of mortality.

Limitations to data interpretation and comparison: A number of factors need to be considered when reviewing cancer statistics and interpreting them. A cancer registry database is a fluid and dynamic database, therefore, the reported number of new cases in a particular race, gender, and age cancer category may change for the calendar year for which the data have already been reported in a previous publication. Additional cancer cases which have been previously overlooked for a given diagnosis year may be found and reported to the central registry. There may also be elimination of duplicate records for the same patient, often due to name changes or spelling corrections.

Metastasis: When cancer spreads from the primary site to other organs or tissues of the body, it is said to metastasize. Cancer usually spreads through the blood or the lymphatic system.

Mortality/incidence ratio (M/I): This ratio is calculated by dividing the number of deaths in a given year by the number of new cancers diagnosed in the same year. The death to case ratio provides a crude indication of the prognosis for patients. A ratio approaching 1.0, when the number of deaths equals the number of cases for a particular type of cancer, indicates a poor prognosis. A lower ratio indicates fewer deaths relative to the number of cases and suggests a better prognosis.

Percent change: This is the difference between two rates expressed as a percentage.

Racial misclassifications: When race is not specified in a source document and the default is to record these cases as white or unknown, the results are considered biased. Numerator error can occur because of misclassification.

Rate comparisons: When comparing age-adjusted rates and age-specific rates based on fewer than 10 cases, rate comparisons are difficult to interpret. In comparing rates among geographic areas such as counties, states and health districts, the absolute numbers and differences in demographics should be considered, as well as clinical significance of the disease. Data quality indicators for each registry should also be reviewed. Interpretations made without considering these factors may be misleading. There will also be differences between mortality statistics published by various agencies and the mortality rates in this report.

Risks and associated risk factors: These were developed using the "American Cancer Society Textbook of Oncology," and the Harvard Cancer Center, Causes of Human Cancer.

Stage at time of diagnosis: Staging is the process of describing the extent or spread of disease from the origin, which is the primary site. Summary staging is the standard used for comparison nationally. SEER Summary Stages 2000 are defined as follows:

- **In Situ:** Malignant cells are within the cell group from which they arose, without penetration of the basement membrane of the tissue and no stromal invasion. *In situ* is “in place”.
- **Localized:** The malignant cells are limited to the organ of origin and have spread no farther than the organ in which they started.
- **Regional:** The tumor is beyond the limits of the organ of origin by direct extension to adjacent areas with or without lymph node involvement.
- **Distant:** The primary tumor has broken away and has traveled, growing secondary tumors in other parts of the body. It has metastasized.

In situ and localized stages are the **early stages** of diagnosis. Regional and distant stages are **late stage** diagnoses.

Staging: Stage is based on an assessment of the size of the primary tumor, whether it has spread, and, if so, how far. Because an accurate diagnosis is so important to effective treatment, physicians might use physical exams, imaging, lab tests, a biopsy, an analysis of the patient’s body fluids, and surgery in various combinations in the staging process. Advancement in diagnostic procedures may change in due time. These advancements might increase the chance that a given cancer will be diagnosed at a more advanced stage, for example with new scanning methods metastases can be detected. Therefore, if someone was previously diagnosed with a localized tumor, they may now be staged as distant. This is called stage migration and can affect the analysis of all solid tumors.

Statistical significance: This determines whether an event happens by chance alone. The null hypothesis states that in a given place and a period of time, all events occur randomly by chance. If not, then there is statistical significance. Confidence intervals are used to test statistical significance in this report. If the confidence intervals of two different rates intersect each other, then there is no statistical difference between the two rates.³ However, if the confidence intervals do not intersect one another, there is statistical significance. This report looks at the South Dakota rates as compared to the US national rates using SEER data.

In South Dakota, case counts can be very low; therefore, magnitude bias is inherent with confidence intervals and z- tests. For example, in the year 2001, cervical cancer rates were 10 per 100,000 American Indian women, a cervical cancer age-adjusted rate six times higher than white women in South Dakota. However, the case counts were two for American Indians and 10 whites. Small numbers result in wider confidence intervals, thus less confidence in the data.

Years of potential life lost (YPLL): The years of potential life lost is calculated for each individual who dies of a cancer of interest by determining the number of years of additional expected life if that person had lived to 75 years. The YPLL in the general population associated with a particular cancer is the sum of this expectation over all those individuals who died of that cancer in a particular year. YPLL reflects the burden of cancer on younger persons while death rates reflect the burden on older persons.

¹<http://epi.grants.cancer.gov/ResPort/HDOverview.html>

²<http://www.omni.org/docs/CMHFProceedings.pdf>

³BIostatistics *The Bare Essentials*, 2nd edition Norman and Shreiner Page 512

**Table 1 : Cancer Cases and Incidence Rates by County
South Dakota, 2014 and 2010-2014 Average**

IV. CANCER INCIDENCE

South Dakota collected 4,498 new reportable cancer cases in 2014. Data at the county level ranged from a low incidence rate of 92.7 in Ziebach County to a high of 775.0 in Kingsbury County. There were eight counties with rates significantly lower than the state incidence rate of 451.5. Four counties had a significantly higher rate, compared to two counties in 2012.

The United States incidence rate for 2014 was 428.6 and the 2014 South Dakota incidence rate was 451.5 per 100,000 persons.

On page 14 of this report a map displaying the 2014 incidence rates by county can be located. On page 15 is a map displaying the mortality rates by county.

County	2014		2010-2014 [^]	
	Cases	Rate	Cases	Rate
South Dakota	4,498	451.5	4,415	457.5
Aurora	21	555.8	20	529.6
Beadle	110	470.0	106	458.2
Bennett	16	499.7	15	467.6
Bon Homme	52	527.5	38	382.7 ▼
Brookings	141	508.6	127	470.5
Brown	214	462.9	201	436.2 ▼
Brule	33	454.1	31	448.1
Buffalo	6	484.4	7	553.3
Butte	61	448.6	58	435.6
Campbell	7	337.1	8	313.6 ▼
Charles Mix	70	586.7	60	513.6
Clark	24	411.5	24	441.3
Clay	57	466.6	52	443.3
Codington	137	407.6	149	459.4
Corson	15	390.1	14	374.6 ▼
Custer	58	405.7	51	370.2 ▼
Davison	112	427.1	125	499.5
Day	35	381.6	39	429.3
Deuel	33	539.9	28	444.9
Dewey	19	377.5	27	570.4 ▲
Douglas	30	655.4	24	506.9
Edmunds	17	255.3 ▼	27	428.7
Fall River	72	585.0	61	492.2
Faulk	17	418.5	19	482.4
Grant	39	356.8	48	460.0
Gregory	41	712.4 ▲	41	629.2 ▲
Haakon	15	437.5	18	581.6
Hamlin	25	341.0	29	406.7
Hand	27	497.2	23	397.6
Hanson	23	646.6	18	526.1
Harding	5	310.4	5	276.5 ▼
Hughes	104	480.8	104	502.2
Hutchinson	42	364.4	49	429.0
Hyde	10	480.7	10	508.9
Jackson	22	643.8	12	356.8 ▼
Jerauld	19	565.0	16	477.9
Jones	10	746.4	6	385.9
Kingsbury	58	775.0 ▲	49	644.2 ▲
Lake	79	454.1	69	429.4
Lawrence	114	336.2 ▼	123	383.8 ▼
Lincoln	220	465.8	174	416.0 ▼
Lyman	23	525.0	20	461.2
McCook	46	756.0 ▲	38	524.7
McPherson	19	219.2 ▼	16	276.1 ▼
Marshall	21	473.8	25	404.0
Meade	110	357.7 ▼	114	398.6 ▼
Mellette	6	302.3	10	448.4
Miner	18	540.0	18	499.0
Minnehaha	944	512.4 ▲	927	527.3 ▲
Moody	29	343.3	30	371.9 ▼
Oglala Lakota	50	502.3	39	424.9
Pennington	513	406.9	514	433.3 ▼
Perkins	7	144.8 ▼	18	377.3 ▼
Potter	20	407.4	22	529.3
Roberts	41	312.6 ▼	52	395.2 ▼
Sanborn	14	376.9	17	534.5
Spink	44	424.9	43	445.7
Stanley	14	356.2	20	495.4
Sully	3	109.5 ▼	6	277.1 ▼
Todd	30	435.7	26	409.5
Tripp	38	462.7	38	442.9
Turner	62	563.2	60	520.3
Union	90	504.1	96	554.5 ▲
Walworth	32	331.5	32	347.0 ▼
Yankton	111	391.2	120	418.7 ▼
Ziebach	3	92.7 ▼	4	163.2 ▼

* Counts less than three are suppressed. Mortality rates with counts less than 20 are generally considered unstable. ▲ Rate significantly higher. ▼ Rate significantly lower.
[^] Number of the cases and rates are averaged over the five-year period.
 Rates per 100,000 age-adjusted to 2000 US standard population and SD 2014 estimated population. Source: South Dakota Department of Health

Table 2: Age-adjusted Incidence Rates by County for Selected Sites, 2014

	Colorectal		Lung and Bronchus		Female Breast		Prostate		Bladder		NHL	
	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
South Dakota	447	44.2	582	56.4	608	122.2	549	107.7	203	19.7	188	18.9
Aurora	5	118.3	*	54.6	*	62.3	3	139.4	*	43.2	0	0.0
Beadle	12	50.9	16	68.7	9	81.3	14	115.8	3	12.2	3	11.3
Bennett	0	0.0	0	0.0	4	271.2	*	48.3	0	0.0	*	25.5
Bon Homme	6	58.1	4	32.4	5	112.8	5	92.8	*	21.1	*	9.2
Brookings	12	43.7	8	31.0	28	205.2	11	78.4	6	21.8	*	9.2
Brown	17	37.6	36	72.6	35	149.7	27	118.2	5	11.1	6	12.3
Brule	4	45.7	6	76.9	*	55.5	4	111.6	*	27.6	*	16.4
Buffalo	*	141.0	4	293.0	0	0.0	*	94.6	0	0.0	0	0.0
Butte	11	76.6	12	73.7	7	102.7	6	81.9	*	8.0	*	15.0
Campbell	0	0.0	0	0.0	0	0.0	4	240.5	0	0.0	0	0.0
Charles Mix	6	43.1	7	53.5	7	138.0	15	244.6	5	36.1	*	14.6
Clark	*	16.7	*	20.0	*	27.1	*	69.2	3	59.4	0	0.0
Clay	6	47.2	6	47.4	4	72.5	9	152.2	5	43.0	6	47.9
Codington	22	70.1	19	54.5	11	65.5	15	88.9	9	25.2	9	28.9
Corson	*	18.7	0	0.0	3	161.5	4	225.6	*	24.3	0	0.0
Custer	7	38.5	8	45.7	7	134.4	7	88.3	0	0.0	*	12.9
Davison	9	31.5	20	75.2	9	68.2	20	166.8	4	9.6	*	3.9
Day	4	41.3	3	30.2	4	80.2	5	100.8	*	19.4	*	6.8
Deuel	6	82.5	*	10.8	5	169.4	*	58.1	0	0.0	3	38.0
Dewey	0	0.0	*	17.3	7	281.9	5	211.4	0	0.0	0	0.0
Douglas	5	116.0	5	75.4	5	258.0	4	190.3	0	0.0	*	18.7
Edmunds	*	13.4	*	30.6	*	57.9	*	55.3	0	0.0	*	15.2
Fall River	6	44.0	15	112.4	10	190.8	7	102.4	3	18.4	4	27.4
Faulk	3	65.6	*	31.8	*	111.5	*	99.3	0	0.0	*	25.0
Grant	*	8.1	5	42.8	9	151.9	*	32.7	*	9.1	4	36.9
Gregory	4	42.3	5	69.1	8	287.2	8	268.4	0	0.0	4	71.7
Haakon	*	57.6	4	56.9	4	234.0	*	137.7	*	36.1	0	0.0
Hamlin	*	29.6	3	45.7	*	77.7	7	165.2	3	31.5	0	0.0
Hand	*	16.9	*	34.4	*	13.0	6	204.6	4	57.2	*	40.4
Hanson	3	108.5	4	104.5	4	212.2	*	112.5	*	62.9	*	69.7
Harding	0	0.0	*	41.8	*	100.9	0	0.0	0	0.0	0	0.0
Hughes	11	51.3	11	51.1	16	142.5	11	103.7	9	43.3	4	14.9
Hutchinson	5	42.2	*	11.5	4	70.5	6	96.9	*	29.7	3	20.8
Hyde	*	43.9	*	37.0	*	264.3	*	73.0	0	0.0	*	30.2
Jackson	5	130.7	*	67.2	*	141.3	4	210.0	0	0.0	*	26.5
Jerault	*	57.4	4	127.4	4	88.6	*	147.8	*	23.5	*	118.8
Jones	5	379.2	*	55.1	0	0.0	0	0.0	*	112.6	*	61.2
Kingsbury	13	181.1	8	95.8	9	226.9	4	109.8	*	18.8	3	35.7
Lake	3	16.2	8	36.3	14	185.9	15	156.9	*	7.3	5	28.1
Lawrence	18	49.9	14	39.8	21	116.2	11	66.1	6	15.9	*	6.0
Lincoln	14	29.3	21	45.3	39	165.6	23	103.9	11	24.9	11	22.7
Lyman	*	28.1	5	105.6	4	202.8	4	158.0	3	61.1	*	22.1
McCook	*	18.0	9	132.8	3	146.1	8	220.3	*	32.3	*	22.2
McPherson	*	19.7	*	26.1	*	41.8	*	25.0	*	11.9	*	24.4
Marshall	*	36.9	*	29.2	*	41.8	4	187.8	*	14.6	*	36.9
Meade	8	22.6	25	79.2	12	86.3	9	57.7	7	22.5	3	9.7
Mellette	*	36.9	*	32.1	*	118.4	0	0.0	*	36.9	*	40.6
Miner	3	53.2	*	57.9	*	142.8	3	185.1	0	0.0	*	23.2
Minnehaha	77	41.8	111	62.5	132	135.7	121	125.9	45	24.8	28	15.0
Moody	*	21.7	3	33.3	*	47.1	6	152.8	0	0.0	*	28.6
Oglala Lakota	6	51.4	9	118.8	9	183.8	3	59.8	*	10.8	0	0.0
Pennington	46	38.6	81	62.6	59	86.4	41	59.8	29	23.0	29	22.2
Perkins	*	16.2	*	28.1	*	54.0	*	34.5	0	0.0	0	0.0
Potter	*	38.0	3	71.1	*	74.6	7	301.3	*	24.1	*	45.3
Roberts	4	35.7	7	44.2	5	76.4	9	124.9	0	0.0	0	0.0
Sanborn	*	51.0	*	38.3	0	0.0	3	175.2	0	0.0	*	25.5
Spink	6	67.4	4	38.7	9	145.7	5	98.7	0	0.0	3	21.6
Stanley	0	0.0	4	97.5	*	128.2	3	166.1	*	20.6	0	0.0
Sully	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Todd	3	49.9	*	10.6	7	173.9	0	0.0	*	33.0	*	22.4
Tripp	9	106.8	0	0.0	6	162.9	3	67.5	0	0.0	4	46.9
Turner	8	70.2	7	55.4	7	111.6	8	129.3	3	19.3	3	41.3
Union	9	50.8	12	64.2	12	138.7	10	108.3	4	21.9	3	18.4
Walworth	*	10.2	7	70.2	4	59.0	7	156.1	0	0.0	*	15.9
Yankton	17	64.0	14	43.9	18	123.6	13	94.9	3	9.8	6	20.3
Ziebach	0	0.0	0	0.0	0	0.0	*	57.0	0	0.0	0	0.0

Note: * Counts less than 3 are suppressed. Incidence rates with counts less than 20 are generally considered unstable.
 Rates per 100,000 age-adjusted to the 2000 US standard population and 2014 SD estimated population.
 Source: South Dakota Department of Health.

Table 3: Age-adjusted Incidence Rates by Site, Gender and Race, South Dakota, 2014

2014	TOTAL		MALE		FEMALE		WHITE		AMERICAN INDIAN	
	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
Total	4,498	451.5	2,401	505.7	2,097	413.4	4,202	454.8	245	532.9
Oral Cavity	145	14.2	113	23.0	32	6.2	139	14.8	5	8.0
Lip	25	2.4	17	3.4	8	1.6	25	2.6	0	0.0
Tongue	32	3.3	25	5.2	7	1.4	30	3.5	2	3.2
Salviary Gland	15	1.4	12	2.4	3	0.5	15	1.4	0	0.0
Floor of Mouth	9	0.9	6	1.2	3	0.6	8	0.9	1	1.6
Gum and Other Mouth	7	0.6	3	0.6	4	0.6	6	0.5	1	1.6
Nasopharynx	4	0.4	4	0.8	0	0.0	2	0.2	1	1.6
Tonsil	38	3.8	35	7.1	3	0.7	38	4.1	0	0.0
Oropharynx	8	0.8	5	1.2	3	0.5	8	0.9	0	0.0
Hypopharynx	5	0.5	4	0.7	1	0.2	5	0.5	0	0.0
Other Oral Cavity and Pharynx	2	0.2	2	0.4	0	0.0	2	0.2	0	0.0
Digestive System	815	80.1	469	98.6	346	64.0	753	79.5	48	99.0
Esophagus	46	4.2	41	7.9	5	0.8	45	4.4	1	1.6
Stomach	46	4.4	28	5.7	18	3.3	40	4.2	6	14.2
Small Intestine	23	2.5	12	2.6	11	2.3	23	2.7	0	0.0
Colorectal	447	44.2	247	53.0	200	37.0	415	44.3	25	52.8
Colon Excluding Rectum	321	31.6	169	37.2	152	27.2	305	32.3	14	30.5
Rectum and Rectosigmoid	126	12.6	78	15.8	48	9.8	110	12.1	11	22.3
Anus, Anal Canal and Anorectum	13	1.2	4	0.8	9	1.7	12	1.3	1	1.6
Liver and Intrahepatic Bile Duct	60	5.9	41	8.1	19	3.6	49	5.1	8	14.7
Gallbladder	9	0.9	4	0.9	5	0.8	8	0.9	1	2.9
Other Biliary	24	2.2	14	2.9	10	1.7	24	2.3	0	0.0
Pancreas	131	12.9	73	15.4	58	10.6	121	12.7	6	11.2
Retroperitoneum	4	0.4	3	0.7	1	0.2	4	0.5	0	0.0
Peritoneum, Omentum and Mesentery	12	1.2	2	0.4	10	1.9	12	1.2	0	0.0
Respiratory	623	60.8	359	76.3	264	48.9	582	60.3	37	108.7
Nose, Nasal Cavity and Middle Ear	4	0.3	2	0.3	2	0.3	4	0.3	0	0.0
Larynx	34	3.7	28	6.5	6	1.1	30	3.5	4	11.1
Lung and Bronchus	582	56.4	327	69.0	255	47.4	545	56.1	33	97.5
Pleura	1	0.1	1	0.2	0	0.0	1	0.1	0	0.0
Mediastinum and Other Resp Organs	2	0.2	1	0.2	1	0.2	2	0.2	0	0.0
Bones and Joints	7	0.8	3	0.7	4	0.9	6	0.8	1	0.8
Soft Tissue (Including Heart)	25	3.0	16	4.2	9	1.8	22	2.8	3	4.8
Skin	257	27.8	136	29.2	121	27.6	254	30.2	3	5.5
Melanomas of the Skin	239	25.8	127	27.0	112	25.6	237	28.2	2	3.4
Other Skin	18	2.0	9	2.2	9	1.9	17	2.0	1	2.1
Breast	610	63.1	2	0.4	608	122.3	559	62.7	46	100.7
Breast, Female	608	122.3			608	122.3	557	121.7	46	183.5
Breast, Male	2	0.4	2	0.4			2	0.4	0	0.0
Female	260	52.1			260	52.1	238	50.7	19	69.4
Vulva	16	2.8			16	2.8	15	2.8	1	6.1
Vagina	1	0.2			1	0.2	1	0.2	0	0.0
Cervix Uteri	36	8.8			36	8.8	27	6.9	9	34.4
Corpus and Uterus, NOS	147	28.9			147	28.9	140	29.5	6	20.1
Corpus Uteri	145	28.5			145	28.5	138	29.1	6	20.1
Uterus, NOS	2	0.4			2	0.4	2	0.4	0	0.0
Ovary	55	10.5			55	10.5	51	10.5	3	8.8
Other Female Genital Organs	5	0.9			5	0.9	4	0.8	0	0.0

Table 3: Age-adjusted Incidence Rates by Site, Gender and Race, South Dakota, 2014 (continued)

	TOTAL		MALE		FEMALE		WHITE		AMERICAN INDIAN	
	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
Male	593	118.6	593	118.6			557	120.2	33	150.7
Penis	10	2.1	10	2.1			10	2.2	0	0.0
Prostate	549	107.7	549	107.7			515	108.0	31	143.5
Testis	32	8.3	32	8.3			31	9.5	1	3.9
Other Male Genital Organs	2	0.6	2	0.6			1	0.4	1	3.3
Urinary	385	38.4	290	63.3	95	17.7	356	37.9	23	51.5
Bladder	203	19.7	161	35.2	42	7.4	197	20.2	5	12.7
Kidney and Renal Pelvis	175	17.9	125	27.0	50	9.7	153	17.0	18	38.8
Ureter	4	0.5	1	0.3	3	0.6	4	0.5	0	0.0
Other Urinary Organs	3	0.3	3	0.8	0	0.0	2	0.2	0	0.0
Eye and Orbit	7	0.6	4	0.7	3	0.5	6	0.6	1	2.9
Brain and CNS	68	7.1	41	8.8	27	5.5	61	6.8	3	2.9
Brain	65	6.7	39	8.3	26	5.3	60	6.6	2	2.1
Meninges and CNS	3	0.4	2	0.5	1	0.2	1	0.2	1	0.8
Endocrine	112	13.0	33	7.5	79	18.7	109	14.3	2	3.5
Thyroid	103	12.0	26	6.1	77	18.2	100	13.2	2	3.5
Other Endocrine	9	1.0	7	1.4	2	0.5	9	1.1	0	0.0
Lymphomas	205	20.9	121	26.9	84	15.7	194	21.4	6	15.2
Hodgkin's Lymphoma	17	1.9	12	2.7	5	1.2	14	1.8	0	0.0
Non-Hodgkin's Lymphoma	188	18.9	109	24.1	79	14.4	180	19.6	6	15.2
Multiple Myeloma	47	4.6	35	7.6	12	2.4	42	4.4	4	6.9
Leukemia	153	15.2	90	19.0	63	12.5	148	16.0	3	5.6
Acute Lymphocytic	11	1.3	6	1.4	5	1.2	9	1.4	0	0.0
Chronic Lymphocytic	60	5.6	39	8.0	21	3.7	60	6.0	0	0.0
Other Lymphocytic	6	0.6	3	0.7	3	0.6	5	0.5	1	1.6
Acute Myeloid	38	3.9	20	4.3	18	3.7	37	4.1	1	2.0
Acute Monocytic	2	0.2	0	0.0	2	0.4	2	0.2	0	0.0
Chronic Myeloid	32	3.3	21	4.5	11	2.5	31	3.4	1	2.1
Other Myeloid/Monocytic	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other Acute Leukemia	1	0.1	0	0.0	1	0.1	1	0.1	0	0.0
Other Leukemia	3	0.2	1	0.1	2	0.3	3	0.3	0	0.0
Myeloproliferative Myelodysplastic	85	8.7	42	9.5	43	8.0	81	8.8	3	4.5
Mesothelioma	5	0.4	5	1.0	0	0.0	5	0.4	0	0.0
Other Sites	96	9.5	49	10.5	47	8.7	90	9.5	5	9.6

Rates per 100,000 age-adjusted to the 2000 US standard population and 2014 SD estimated population.
Source: South Dakota Department of Health

Table 3 above shows incidence and age-adjusted incidence rates for South Dakota in 2014 by primary sites, gender, and race according to SEER site category recodes (Appendix C).

Table 4: Percentage of Selected Cancers by Age Groups in South Dakota, 2014


	0-19	20-34	35-49	50-64	65-74	75-84	85+
All Sites	1%	3%	9%	32%	26%	21%	10%
Bladder	0%	0%	3%	19%	32%	28%	18%
Breast, Female	0%	2%	14%	37%	23%	17%	7%
Colorectal	0%	2%	7%	29%	25%	23%	15%
Corpus and Uterus, NOS	0%	1%	14%	48%	22%	13%	1%
Hodgkin's Lymphoma	6%	47%	18%	24%	0%	0%	6%
Kidney and Renal Pelvis	2%	0%	12%	35%	24%	20%	7%
Leukemia	8%	1%	7%	23%	24%	20%	16%
Lung and Bronchus	0%	0%	3%	28%	28%	31%	11%
Melanomas of the Skin	0%	8%	19%	33%	18%	15%	8%
Non-Hodgkin's Lymphoma	1%	2%	9%	29%	24%	26%	10%
Pancreas	0%	1%	5%	24%	32%	22%	16%
Prostate	0%	0%	1%	39%	38%	17%	5%
Thyroid	2%	16%	31%	28%	17%	3%	4%

Source: South Dakota Department of Health

In 2014, 57% of all cancers were diagnosed between ages 50 to 74 (Table 4). Notable were the 14% of female breast and corpus and uterus, NOS cancers and 31% of thyroid cancers diagnosed between the ages of 35 to 49. In 2014, 47% of the Hodgkin's lymphoma cases were diagnosed in persons under 35 years old, compared to 39% in 2012.

Figure 1 below shows that incidence rates for American Indians in South Dakota were higher than those for whites in 2014. Of the 4,498 newly diagnosed cases in 2014, 245 or 5.4% were American Indians, 119 males and 126 females.

Figure 1: All Sites Cancer Incidence Rates by Race and Gender, South Dakota, 2014




Note: Rates per 100,000 age-adjusted to the 2000 US standard population and 2014 SD estimated population.
Source: South Dakota Department of Health

V. CANCER CASES AND DEATHS BY RANK

Female breast cancer was the most common cancer diagnosed during 2014. The five most diagnosed cancers were female breast, lung and bronchus, prostate, colorectal, and melanoma which accounted for 53.9% of the new cases diagnosed and 49.6% of cancer deaths. Figure 2 shows the percent of new cancer cases and deaths by rank and gender.

Figure 2: Percent Distribution of Cancer Cases and Deaths by Rank and Gender, South Dakota, 2014




Source: South Dakota Department of Health

VI. STAGE AT DIAGNOSIS

SEER Summary Staging:

- ***In Situ*** - Malignant cells are within the cell groups from which they arose, without penetration of the basement membrane of the tissue and stromal invasion.
- **Localized** - The malignant cells are limited to the organ of origin and have spread no farther than the organ where they began.
- **Regional** - The tumor is beyond the limits of the organ of origin by direct extension to adjacent areas such as the regional lymph nodes, adjacent organs, or tissue.
- **Distant** - The tumor cells have broken away from the primary tumor and traveled to other parts of the body.
- **Unknown** - If extension or metastatic, there is not sufficient evidence available to assign a stage.

Figure 5: Stage at Diagnosis, All Sites, South Dakota, 2014



Source: South Dakota Department of Health


The figure above demonstrates the number of cases diagnosed at each stage of disease. For 2014, there were a total of 4,905 cases reported to the South Dakota State Cancer Registry, which includes *in situ* cases.

Table 5: South Dakota Stage at Diagnosis, All Cases, 2014

Stage	Number of Cases	Percent of Total
<i>In Situ</i>	520	11%
Localized	2031	41%
Regional	895	18%
Distant	1175	24%
Unknown	284	6%


When analyzed by race, 41% of all cancer cases diagnosed for whites were diagnosed at localized stage and 44% for American Indians. American Indians had another 46% diagnosed at the regional and distant stages combined while the same stages for whites were a combined 42%. Stage at diagnosis by race was as follows:

Figure 6: Stage at Diagnosis in South Dakota, White, 2014



Source: South Dakota Department of Health

Figure 7: Stage at Diagnosis in South Dakota, American Indians, 2014



Source: South Dakota Department of Health

Cases of non-Hodgkin's lymphoma, myeloma and leukemias are usually at distant stages, and therefore can skew the proportion of all sites diagnosed at distant stages. Some differences in case counts by stage for selected sites are shown in Table 6.

Table 6: Stage at Diagnosis for Selected Sites by Race, South Dakota, 2013

	White						American Indian					
	Localized		Regional		Distant		Localized		Regional		Distant	
	Cases	%	Cases	%	Cases	%	Cases	%	Cases	%	Cases	%
Female Breast	396	20.9%	130	15.6%	26	2.4%	29	25.9%	15	28.3%	1	1.6%
Lung and Bronchus	117	6.2%	114	13.7%	286	26.1%	5	4.5%	6	11.3%	22	34.4%
Prostate	423	22.3%	51	6.1%	31	2.8%	28	25.0%	1	1.9%	2	3.1%
Melanoma of the Skin	202	10.6%	11	1.3%	18	1.6%	2	1.8%	0	0.0%	0	0.0%
Colorectal	158	8.3%	152	18.2%	83	7.6%	5	4.5%	14	26.4%	6	9.4%
Bladder	53	2.8%	17	2.0%	13	1.2%	2	1.8%	1	1.9%	0	0.0%
Non-Hodgkin's Lymphoma	39	2.1%	29	3.5%	94	8.6%	1	0.9%	1	1.9%	4	6.3%
Kidney and Renal Pelvis	73	3.8%	31	3.7%	35	3.2%	13	11.6%	2	3.8%	2	3.1%
Corpus and Uterus, NOS	106	5.6%	22	2.6%	11	1.0%	4	3.6%	2	3.8%	0	0.0%
Leukemia	0	0.0%	1	0.1%	146	13.3%	0	0.0%	0	0.0%	3	4.7%
Thyroid	57	3.0%	37	4.4%	3	0.3%	1	0.9%	1	1.9%	0	0.0%
Pancreas	15	0.8%	32	3.8%	60	5.5%	0	0.0%	0	0.0%	5	7.8%

Source: South Dakota Department of Health

Table 7 : Cancer Deaths and Mortality Rates by County
South Dakota, 2014 and 2010-2014 Average

VII. CANCER MORTALITY

Cancer age-adjusted death rates for 2014 ranged from a low of 28.5 in Ziebach County to a high of 362.8 in Buffalo County. South Dakota's age adjusted death rate was 161.4 in 2014 compared to a 5-year death rate of 162.4.

In 2014, nine counties had a significantly lower rate than that of the entire state and one had a significantly higher rate. The five-year rates show nine counties having significantly lower rates and four counties with significantly higher rates. South Dakota's mortality rate for 2010-2014 was 162.4 per 100,000 persons.

The United States mortality rate for 2014 was 161.3 and the South Dakota rate was 161.4 per 100,000 persons. When the United States and South Dakota rates are compared for 2014 there is no significant difference.

The South Dakota 2014 mortality rates are displayed in a state map on page 15 of this report.

County	2014		2010-2014 [^]	
	Deaths	Rate	Deaths	Rate
South Dakota	1,679	161.4	1,636	162.4
Aurora	6	137.6	7	159.7
Beadle	47	181.2	41	164.6
Bennett	8	230.9	6	179.2
Bon Homme	16	146.7	15	136.3
Brookings	49	167.8	47	166.8
Brown	96	191.1	80	165.0
Brule	11	135.6	12	154.2
Buffalo	5	362.8	3	285.9
Butte	14	92.3	25	175.8
Campbell	*	69.2	3	95.8
Charles Mix	24	170.9	21	160.6
Clark	17	301.5	11	170.9
Clay	24	193.0	21	174.9
Codington	70	195.9	60	174.1
Corson	9	225.5	8	211.3
Custer	15	99.6	21	148.5
Davison	59	208.7	48	172.5
Day	10	116.2	13	124.5
Deuel	5	67.7	7	103.0
Dewey	10	211.8	10	231.0
Douglas	9	190.3	9	182.4
Edmunds	5	66.6	10	138.7
Fall River	21	153.7	24	188.1
Faulk	9	169.8	8	164.1
Grant	12	102.6	15	130.7
Gregory	16	189.2	13	170.3
Haakon	4	152.8	5	171.7
Hamlin	10	111.9	11	137.5
Hand	8	109.4	9	133.2
Hanson	8	325.4	6	180.3
Harding	*	41.8	*	67.0
Hughes	31	151.5	32	155.5
Hutchinson	19	130.7	18	127.0
Hyde	3	107.9	3	121.6
Jackson	7	209.4	6	179.2
Jerauld	9	195.6	7	167.9
Jones	3	183.2	*	122.8
Kingsbury	13	162.0	15	167.1
Lake	25	127.5	26	152.6
Lawrence	47	140.8	55	160.6
Lincoln	62	137.5	49	122.3
Lyman	8	178.4	6	137.7
McCook	20	330.2	17	211.7
McPherson	8	85.4	6	99.0
Marshall	9	204.9	12	183.7
Meade	45	155.1	42	150.7
Mellette	5	187.1	6	237.2
Miner	4	78.2	7	153.3
Minnehaha	298	160.1	313	179.5
Moody	16	195.8	14	169.7
Oglala Lakota	17	197.5	17	219.9
Pennington	207	162.8	195	159.9
Perkins	11	205.4	12	207.8
Potter	4	72.3	7	117.6
Roberts	28	190.0	27	196.3
Sanborn	6	181.2	6	180.2
Spink	10	100.2	16	144.9
Stanley	7	205.6	6	156.0
Sully	4	161.5	4	163.3
Todd	14	230.4	12	212.4
Tripp	8	76.0	14	146.0
Turner	26	198.9	21	166.3
Union	30	165.6	28	157.4
Walworth	19	216.4	18	171.5
Yankton	53	161.1	44	144.1
Ziebach	*	28.5	*	54.6

* Counts less than three are suppressed. Mortality rates with counts less than 20 are generally considered unstable. ▲ Rate significantly higher; ▼ Rate significantly lower

[^] Number of the cases and rates are averaged over the five-year period.

Rates per 100,000 age-adjusted to 2000 US standard population and SD 2014 estimated population. Source: South Dakota Department of Health

Table 8: Age-adjusted Mortality Rates by County for Selected Sites, 2014

	Colorectal		Lung and Bronchus		Female Breast		Prostate		Bladder		NHL	
	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate
South Dakota	188	17.6	439	41.9	100	18.2	439	93.2	39	3.7	52	4.9
Aurora	*	23.9	*	40.6	0	0.0	*	96.4	0	0.0	0	0.0
Beadle	4	12.7	14	56.5	*	9.5	14	128.0	3	10.6	0	0.0
Bennett	3	84.3	*	25.8	*	47.4	*	48.3	0	0.0	*	23.1
Bon Homme	*	14.3	3	29.1	3	66.3	3	62.9	0	0.0	0	0.0
Brookings	8	26.8	6	20.4	3	20.2	6	47.9	*	7.3	*	4.4
Brown	7	15.5	24	46.3	10	36.1	24	111.6	0	0.0	*	3.8
Brule	*	29.3	*	22.0	0	0.0	*	57.7	0	0.0	0	0.0
Buffalo	*	80.2	3	219.1	0	0.0	3	394.6	0	0.0	0	0.0
Butte	*	13.9	3	16.2	*	10.7	3	32.3	0	0.0	0	0.0
Campbell	0	0.0	0	0.0	*	61.4	0	0.0	0	0.0	0	0.0
Charles Mix	*	6.6	11	79.6	*	15.4	11	177.4	*	5.9	0	0.0
Clark	*	38.8	*	40.0	*	50.8	*	83.6	*	20.0	0	0.0
Clay	*	7.2	5	37.0	*	26.4	5	91.5	0	0.0	*	8.8
Codington	5	12.6	22	62.6	5	28.1	22	139.5	0	0.0	4	12.5
Corson	*	58.9	*	20.2	0	0.0	*	39.6	0	0.0	0	0.0
Custer	*	8.7	3	16.4	*	10.0	3	39.6	0	0.0	0	0.0
Davison	5	18.9	13	46.8	4	16.7	13	112.0	0	0.0	4	16.8
Day	*	11.9	3	27.7	3	90.6	3	57.4	0	0.0	*	8.4
Deuel	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	3	42.6
Dewey	*	37.4	4	82.1	*	50.2	4	172.2	0	0.0	0	0.0
Douglas	*	22.7	0	0.0	*	45.8	0	0.0	0	0.0	*	16.7
Edmunds	0	0.0	0	0.0	*	24.1	0	0.0	0	0.0	0	0.0
Fall River	6	46.2	4	28.8	0	0.0	4	73.9	0	0.0	*	8.0
Faulk	0	0.0	3	56.2	0	0.0	3	111.7	0	0.0	*	14.6
Grant	*	6.7	4	34.6	0	0.0	4	76.5	0	0.0	0	0.0
Gregory	3	23.7	*	30.1	*	37.7	*	68.3	0	0.0	*	7.9
Haakon	0	0.0	*	56.9	0	0.0	*	111.5	0	0.0	0	0.0
Hamlin	*	16.4	*	29.6	*	11.2	*	59.4	*	8.2	0	0.0
Hand	*	31.8	*	11.7	0	0.0	*	27.0	0	0.0	0	0.0
Hanson	3	151.0	*	69.6	0	0.0	*	127.3	0	0.0	0	0.0
Harding	0	0.0	*	41.8	0	0.0	*	71.3	0	0.0	0	0.0
Hughes	4	17.9	7	35.5	3	34.4	7	79.8	0	0.0	*	4.7
Hutchinson	*	3.7	4	28.7	*	5.7	4	61.8	0	0.0	*	3.7
Hyde	0	0.0	*	37.0	0	0.0	*	73.0	0	0.0	0	0.0
Jackson	0	0.0	3	88.7	0	0.0	3	186.0	0	0.0	0	0.0
Jerauld	*	25.6	3	70.2	0	0.0	3	183.2	*	23.5	0	0.0
Jones	0	0.0	3	183.2	0	0.0	3	404.8	0	0.0	0	0.0
Kingsbury	*	7.4	3	34.0	*	11.8	3	64.2	0	0.0	0	0.0
Lake	*	10.4	10	51.0	0	0.0	10	113.6	*	3.6	0	0.0
Lawrence	5	13.6	10	31.1	3	20.5	10	67.6	*	5.9	*	5.8
Lincoln	3	5.8	22	49.1	4	15.1	22	106.7	*	2.8	*	1.4
Lyman	*	46.7	*	37.6	0	0.0	*	70.4	0	0.0	0	0.0
McCook	4	49.9	6	90.2	*	97.3	6	195.3	0	0.0	0	0.0
McPherson	*	7.9	*	11.9	0	0.0	*	26.9	*	7.9	0	0.0
Marshall	0	0.0	*	29.2	0	0.0	*	67.3	*	43.1	0	0.0
Meade	5	18.2	12	39.6	3	17.5	12	86.1	*	6.2	0	0.0
Mellette	*	36.9	*	32.1	0	0.0	*	52.4	0	0.0	*	40.6
Miner	*	19.0	*	44.0	0	0.0	*	104.4	0	0.0	0	0.0
Minnehaha	37	19.6	78	42.5	20	19.3	78	96.5	8	4.3	8	4.1
Moody	3	33.5	*	24.3	0	0.0	*	50.8	*	12.7	0	0.0
Oglala Lakota	4	41.4	5	65.9	*	13.4	5	176.1	0	0.0	0	0.0
Pennington	18	13.8	63	47.9	8	11.9	63	106.3	7	6.7	10	7.6
Perkins	*	43.7	*	11.0	*	37.2	*	31.6	0	0.0	0	0.0
Potter	0	0.0	*	38.0	0	0.0	*	91.3	0	0.0	0	0.0
Roberts	4	25.2	11	71.6	*	21.9	11	152.4	0	0.0	0	0.0
Sanborn	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Spink	*	24.9	3	27.7	0	0.0	3	65.4	0	0.0	0	0.0
Stanley	0	0.0	3	73.9	0	0.0	3	133.4	0	0.0	0	0.0
Sully	0	0.0	0	0.0	*	80.8	0	0.0	*	43.5	*	50.0
Todd	*	26.5	0	0.0	0	0.0	0	0.0	0	0.0	*	21.1
Tripp	*	11.3	*	6.3	0	0.0	*	18.5	0	0.0	0	0.0
Turner	7	53.7	7	54.6	4	59.3	7	127.1	*	7.6	0	0.0
Union	*	4.5	7	37.4	0	0.0	7	78.4	*	8.1	0	0.0
Walworth	*	30.4	6	57.7	0	0.0	6	127.3	0	0.0	*	10.2
Yankton	6	18.1	15	45.9	3	15.9	15	98.0	*	4.1	*	5.6
Ziebach	*	28.5	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

Note: * Counts less than 3 are suppressed. Mortality rates with counts less than 20 are generally considered unstable. Rates per 100,000 age-adjusted to the 2000 US standard population and 2014 SD estimated population.

Source: South Dakota Department of Health

Table 9: Age-adjusted Mortality Rates by Site, Gender, and Race, South Dakota, 2014

	TOTAL		MALE		FEMALE		WHITE		AMERICAN INDIAN	
	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate
Total	1,679	161.4	938	201.2	741	132.0	1,581	160.7	88	217.9
Oral Cavity	23	2.4	16	3.5	7	1.5	22	2.5	1	1.6
Lip	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Tongue	6	0.8	1	0.3	5	1.3	6	0.9	0	0.0
Salivary Gland	1	0.1	1	0.2	0	0.0	1	0.1	0	0.0
Floor of Mouth	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Gum and Other Mouth	4	0.4	2	0.4	2	0.2	4	0.4	0	0.0
Nasopharynx	3	0.4	3	0.7	0	0.0	2	0.3	1	1.6
Hypopharynx	2	0.2	2	0.4	0	0.0	2	0.2	0	0.0
Tonsil	2	0.2	2	0.4	0	0.0	2	0.2	0	0.0
Oropharynx	1	0.1	1	0.2	0	0.0	1	0.1	0	0.0
Other Oral Cavity and Pharynx	4	0.3	4	0.7	0	0.0	4	0.4	0	0.0
Digestive System	448	42.3	276	57.4	172	29.4	420	41.7	23	51.6
Esophagus	41	3.8	32	6.3	9	1.5	39	3.8	2	5.3
Stomach	24	2.2	11	2.3	13	2.2	22	2.1	2	3.8
Small Intestine	6	0.7	2	0.4	4	0.8	6	0.7	0	0.0
Colorectal	188	17.6	110	23.5	78	13.2	176	17.4	10	21.8
Colon Excluding Rectum	155	14.5	85	18.3	70	11.8	146	14.4	8	18.2
Rectum and Rectosigmoid	33	3.1	25	5.2	8	1.4	30	3.0	2	3.5
Anus, Anal Canal and Anorectum	1	0.1	0	0.0	1	0.3	1	0.2	0	0.0
Liver and Intrahepatic Bile Duct	58	5.5	39	7.7	19	3.5	52	5.2	4	9.1
Gallbladder	8	0.9	3	0.7	5	0.9	6	0.6	2	5.5
Other Biliary	4	0.3	3	0.6	1	0.1	4	0.3	0	0.0
Pancreas	118	11.1	76	16.1	42	6.7	114	11.4	3	6.2
Retroperitoneum	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Peritoneum, Omentum and Mesentery	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Respiratory	446	42.6	261	55.2	185	33.5	413	41.6	33	95.6
Nose, Nasal Cavity and Middle Ear	3	0.2	2	0.4	1	0.2	3	0.3	0	0.0
Larynx	4	0.4	4	0.8	0	0.0	4	0.4	0	0.0
Lung and Bronchus	439	41.9	255	54.1	184	33.3	406	40.9	33	95.6
Pleura	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Mediastinum and Other Resp Organs	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Bones and Joints	7	0.8	3	0.7	4	0.9	5	0.6	2	3.9
Soft Tissue	13	1.3	9	1.9	4	0.7	10	1.0	3	6.5
Skin	45	4.6	32	7.6	13	2.2	44	4.9	1	4.4
Melanoma of the Skin	31	3.3	22	5.3	9	1.7	30	3.5	1	4.4
Other Nonepithelial Skin	14	1.3	10	2.3	4	0.6	14	1.4	0	0.0
Breast	101	9.9	1	0.2	100	18.2	98	10.3	3	5.4
Breast, Female	100	18.2			100	18.2	97	18.8	3	10.0
Breast, Male	1	0.2	1	0.2			1	0.2	0	0.0
Female	81	15.5			81	15.5	75	15.2	5	20.5
Vulva	2	0.3			2	0.3	2	0.4	0	0.0
Vagina	0	0.0			0	0.0	0	0.0	0	0.0
Cervix Uteri	11	2.8			11	2.8	8	2.2	3	10.3
Corpus and Uterus, NOS	24	4.2			24	4.2	24	4.4	0	0.0
Corpus Uteri	15	2.7			15	2.7	15	2.9	0	0.0
Uterus, NOS	9	1.5			9	1.5	9	1.6	0	0.0
Ovary	41	7.7			41	7.7	38	7.6	2	10.2
Other Female Genital Organs	3	0.6			3	0.6	3	0.6	0	0.0

**Table 9: Age-adjusted Mortality Rates by Site, Gender, and Race, South Dakota, 2014
(continued)**

	TOTAL		MALE		FEMALE		WHITE		AMERICAN INDIAN	
	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate
Male	77	17.3	77	17.3			72	16.9	5	29.2
Penis	1	0.1	1	0.1			1	0.2	0	0.0
Prostate	75	17.0	75	17.0			71	16.7	4	25.9
Testis	1	0.2	1	0.2			0	0.0	1	3.3
Other Male Genital Organs	0	0.0	0	0.0			0	0.0	0	0.0
Urinary	83	7.8	57	12.2	26	4.3	80	8.0	3	4.8
Bladder	39	3.7	29	6.5	10	1.7	39	3.9	0	0.0
Kidney and Renal Pelvis	40	3.7	26	5.3	14	2.2	37	3.6	3	4.8
Ureter	1	0.1	0	0.0	1	0.2	1	0.1	0	0.0
Other Urinary Organs	3	0.3	2	0.5	1	0.2	3	0.3	0	0.0
Eye and Orbit	2	0.2	1	0.2	1	0.2	2	0.2	0	0.0
Brain and CNS	53	5.2	36	7.5	17	3.2	52	5.6	1	1.6
Brain	53	5.2	36	7.5	17	3.2	52	5.6	1	1.6
Meninges and CNS	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Endocrine	9	0.9	2	0.4	7	1.3	7	0.7	1	1.6
Thyroid	9	0.9	2	0.4	7	1.3	7	0.7	1	1.6
Other Endocrine	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Lymphomas	52	4.9	34	7.4	18	3.1	49	4.9	2	5.4
Hodgkin's Disease	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Non-Hodgkin's Lymphomas	52	4.9	34	7.4	18	3.1	49	4.9	2	5.4
Multiple Myeloma	27	2.6	18	3.9	9	1.5	26	2.6	1	2.9
Leukemia	90	8.8	51	11.5	39	6.6	89	9.2	1	2.0
Acute Lymphocytic	3	0.4	2	0.6	1	0.1	3	0.4	0	0.0
Chronic Lymphocytic	22	1.9	11	2.5	11	1.5	22	2.0	0	0.0
Other Lymphocytic	4	0.4	3	0.6	1	0.2	4	0.4	0	0.0
Acute Myeloid	37	3.7	24	5.3	13	2.5	36	3.9	1	2.0
Acute Monocytic	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Chronic Myeloid	6	0.5	4	1.0	2	0.2	6	0.6	0	0.0
Other Myeloid/Monocytic	5	0.6	1	0.2	4	0.9	5	0.6	0	0.0
Other Acute Leukemia	5	0.5	1	0.3	4	0.7	5	0.5	0	0.0
Other Leukemia	8	0.8	5	1.1	3	0.5	8	0.8	0	0.0
Mesothelioma	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Immunoproliferative Diseases	5	0.4	5	1.0	0	0.0	5	0.5	0	0.0
III-Defined and Unspecified Sites	117	11.2	59	13.1	58	9.8	112	11.3	3	7.0

Rates per 100,000 age-adjusted to the 2000 US standard population and 2013 SD estimated population.
Source: South Dakota Department of Health

Table 9 shows death and age-adjusted death rates by SEER recode primary sites (Appendix D), gender and race. Approximately 1,600 persons die from cancer in South Dakota each year with little or no change in counts. Overall more males than females die from cancer.


Table 10: Percentage of Cancer Deaths by Age Groups and Selected Primary Sites, South Dakota, 2014

Age Group	0-19	20-34	35-49	50-64	65-74	75-84	85+
All Sites	0%	0%	4%	22%	24%	28%	21%
Bladder	0%	3%	5%	18%	13%	36%	26%
Female Breast	0%	0%	9%	24%	17%	25%	25%
Colorectal	0%	0%	4%	18%	19%	28%	31%
Corpus and Uterus, NOS	0%	0%	8%	13%	33%	13%	33%
Meninges, Brain and CNS	4%	0%	8%	45%	25%	17%	2%
Kidney and Renal Pelvis	0%	0%	3%	35%	15%	20%	28%
Leukemia	1%	0%	6%	11%	26%	30%	27%
Lung and Bronchus	0%	0%	1%	24%	30%	30%	15%
Melanoma of the Skin	0%	0%	23%	23%	10%	23%	23%
Non-Hodgkin's Lymphoma	0%	0%	4%	17%	21%	29%	29%
Pancreas	0%	1%	2%	23%	27%	25%	23%
Prostate	0%	0%	0%	5%	21%	33%	40%

Source: South Dakota Department of Health

Overall, in 2014 more persons 75 to 84 years of age died from cancer in South Dakota than any other age group (Table 10). However, there would be cause for concern if too many people died from cancer at a young age.

Figure 8: All Sites Cancer Mortality Rates by Race, and Gender, South Dakota, 2014



Note: Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population.
Source: South Dakota Department of Health

Figure 8 illustrates that American Indian males had a higher death rate than any other group. In 2012 American Indian females had a higher death rate than their male counterparts. American Indian females had a lower death rate than white males or males in total.

VIII. YEARS OF POTENTIAL LIFE LOST


In both 2014 and the five-year period 2010 to 2014, cancer ranked number one in years of potential years lost (YPLL). The number of years lost are shown in Figures 9 and 10 by race. There were 11,017 years of potential years lost due to cancer in 2014, compared to 9,904 years in 2008.

Table 11: Leading Causes of Years of Potential Life Lost (to 75 years of age), South Dakota, 2014

Cancer	11,017
Accidents	9,932
Heart Disease	7,552
Suicide and Self-Inflicted Injury	4,806
Chronic Liver and Cirrhosis	3,187
Diabetes Mellitus	1,560
Chronic Lung Diseases	1,185
Assault (Homicide)	1,066
Cerebrovascular Diseases	1,063
Influenza and Pneumonia	996
Septicemia	730
All Other Causes	12,945


Source: South Dakota Department of Health

Figure 9: Years of Potential Life Lost (YPLL) Before Age 75 for the Leading Causes of Death by Race, South Dakota, 2014



Source: South Dakota Department of Health


Figure 10: Years of Potential Life Lost (YPLL) Before Age 75 for the Leading Causes of Death by Race, South Dakota 2010-2014 Average*



* Number of years are averaged over the five-year period.
Source: South Dakota Department of Health


The differences in YPLL in Figures 11 and 12 reflect the number of cancer cases by primary sites by race. For example, the YPLL for lung and bronchus for whites was 2,262 for 221 deaths whereas the YPLL for American Indians was 196 years for 20 deaths which occurred during 2014. Not all cancers were present among the American Indian population during 2014, hence the differences in the cancer sites presented.

Figure 11: Years of Potential Life Lost for Selected Cancers Among Whites, South Dakota, 2014



Source: South Dakota Department of Health

Figure 12: Years of Potential Life Lost for Selected Cancers Among American Indians, South Dakota, 2014




Source: South Dakota Department of Health

IX. AVERAGE YEARS OF LIFE LOST


South Dakota's average years of life lost (AYLL) due to cancer in 2014 was 12.9 years, a slight decrease from 13.2 years in 2013. Leukemia cancer ranked 15th among cancer sites for American Indians at 9.0 years compared to whites where it ranked 9th with an average of 13.8 years.

Figure 13: Average Years of Life Lost for Selected Cancers Among Whites, South Dakota, 2014



Source: South Dakota Department of Health

Figure 14: Average Years of Life Lost for Selected Cancers Among American Indians, South Dakota, 2014



Source: South Dakota Department of Health

This page intentionally left blank.

X. SELECTED CANCER SITES INCIDENCE AND MORTALITY

This section covers the following cancers: bladder, female breast, cervix uteri, colorectal cancer, corpus uterus, kidney and renal pelvis, leukemia, lung and bronchus, melanoma (skin), myeloma, non-Hodgkin's lymphoma, ovary, pancreas, prostate, stomach, and thyroid.

These cancers were selected because of the ranking in the cancer sites reported as well as the importance and impact to society.

Topics for each cancer include incidence and mortality data along with age-adjusted rates, trends, comparison with national data (if available), risks, associated risk factors, and prevention.


BLADDER

Table 12: Bladder Incidence and Mortality Summary, 2014

Bladder Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	203	161	42	39	29	10
		Age Adjusted Rate	19.7	35.2	7.4	3.7	6.5	1.7
	White	# Cases / Deaths	197	156	41	39	29	10
		Age Adjusted Rate	20.2	35.9	7.7	3.9	6.8	1.8
	American Indian	# Cases / Deaths	5	4	1	0	0	0
		Age Adjusted Rate	12.7	18.0	6.4	0.0	0.0	0.0
United States	Total	Age Adjusted Rate	19.1	33.4	8.2	4.3	7.5	2.1
	White	Age Adjusted Rate	20.8	36.4	8.7	4.6	7.9	2.1
	American Indian	Age Adjusted Rate	9.2	16.1	0.0	2.5	2.8	2.3

¹Includes *in situ* bladder; Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population. US rates www.seer.cancer.gov Source: South Dakota Department of Health

Figure 15: Bladder Cancer Stage at Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: Cancer is categorized as noninvasive and invasive. There were 113 noninvasive bladder cancers reported in 2014. There were 90 invasive. More than half, 56%, of bladder cancer cases in South Dakota were diagnosed at noninvasive, *in situ* stage. Nationally 50% of the cases of urinary bladder cancer are diagnosed at the *in situ* stage. In South Dakota, 6% of the cases were not diagnosed until the disease had spread to distant sites. In the United States, distant stage accounted for 4% of the bladder cancers reported.

Incidence: In 2014, it was estimated that over 74,690 cases of bladder cancer would be diagnosed in the United States. There were 203


cases of bladder cancer reported in South Dakota. There were 161 men and 42 women diagnosed with bladder cancer in 2014. Statistically, men were diagnosed about three times as often as women. There were only five American Indian cases diagnosed in 2014. In the United States it was the eighth most frequent cancer. In South Dakota it was the sixth most frequent cancer diagnosed.

Mortality: Advances in intravesical therapy and in the treatment of advanced disease with chemotherapy have reduced the percentage of mortality from bladder cancer. In South Dakota, ages from 70 and above have the highest mortality. In 2014, the South Dakota mortality rate was 3.7 compared to the US rate which was 4.3.

Risk and Associated Factors: Bladder cancer was one of the first malignancies associated with industrialization. Cigarette smoking increases the risk for bladder cancer by two times that of a nonsmoker. Work exposure to certain chemicals also increases risk. Some of those with the highest risk are makers of rubber, leather, textiles, paint products, and printing compounds.

Prevention and Early Detection: Avoiding exposure to chemicals and cigarette smoking are two of the most common suggestions for prevention.

Figure 16: Bladder Cancer Number of Cases and Deaths by Age, South Dakota, 2014



Source: South Dakota Department of Health




Figure 17: Bladder Cancer Cases and Deaths by Year, South Dakota, 2001 - 2014

Bladder cancer cases declined in 2012 and 2013 after tying an all-time high of 221 cases in 2011, the same number as in 2002.

Source: South Dakota Department of Health




Figure 18: Bladder Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014

Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health


BREAST (FEMALE)

Table 13: Female Breast Incidence and Mortality Summary, 2014

Female Breast Cancer			Incidence	Mortality
South Dakota	Total	# Cases / Deaths	608	100
		Age Adjusted Rate	122.2	18.2
	White	# Cases / Deaths	556	97
		Age Adjusted Rate	121.5	18.8
	American Indian	# Cases / Deaths	46	3
		Age Adjusted Rate	183.5	10.0
United States	Total	Age Adjusted Rate	124.9	20.5
	White	Age Adjusted Rate	127.3	20.0
	American Indian	Age Adjusted Rate	85.5	13.2

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population. US rates www.seer.cancer.gov. Source: South Dakota Department of Health

Figure 19: Female Breast Cancer Stage at Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: Including *in situ* female breast cancer cases there were 747 cases diagnosed in 2014, of which 429 cases were diagnosed at localized stage. This represents 57% of all reported breast cancer cases. There were 146 cases that had progressed beyond the breast. There were 27 that were diagnosed as a distant stage and six that were staged as unknown. The 139 *in situ* female breast cancer cases were reported but were not used in calculating incidence rates.

Incidence: Female breast cancer is the most common malignant tumor among women. The incidence rate increased from 1947-1990. The rates fell 3.5% per year from 2001-2005. This decrease may be in part due to the lower number of women using hormone replacement therapy. There were 608 cases of invasive female breast cancer reported


in 2014. In South Dakota, 13.5% of all invasive cancer cases reported in 2014 were female breast. Nationally, 15% of all cancer cases are female breast cancer. Breast cancer represented 29% of the cancer cases diagnosed for South Dakota women in 2014.

Mortality: Breast cancer is the fourth leading cause of death attributed to cancer in South Dakota. Nationwide, breast cancer mortality has been relatively stable overall since 1950. In cancers only of women, it is the second leading cause of cancer deaths. Although mortality has increased among women older than 55 years, it has decreased among women younger than 55 years of age. In 2014, there were 100 deaths. Of those deaths, 97 were white and three were American Indian.

Risk and Associated Factors: Among the known risk factors for breast cancer are early onset of menarche, late onset of menopause, never having been pregnant, first full term pregnancy after age 30 and fewer number of children. These factors increase the risk because of cumulative exposure of breast tissue to estrogen. Other risk factors include high fat diets, obesity, alcohol consumption, history of fibrocystic disease, having a mother or sister with breast cancer, a personal history of ovarian or endometrial cancer and specific tumor suppressor genes such as BRCA1 and BRCA2.

Prevention and Early Detection: Prevention and early detection is the key to survival of breast cancer. Breast cancers found during screening exams are more likely to be smaller and still confined to the breast. Women should talk to their doctor for individualized screening recommendations.

Figure 20: Female Breast Cancer Number of Cases and Deaths by Age, South Dakota, 2014



Source: South Dakota Department of Health




Figure 21: Female Breast Cancer Cases and Deaths by Year, South Dakota, 2001 - 2014

The incidence for female breast cancer was at an all-time high in 2013.

Source: South Dakota Department of Health




Figure 22: Female Breast Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014

Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health


CERVIX UTERI

Table 14: Cervix Uteri Incidence and Mortality Summary, 2014

Cervix Uteri Cancer			Incidence	Mortality
South Dakota	Total	# Cases / Deaths	36	11
		Age Adjusted Rate	8.7	2.8
	White	# Cases / Deaths	26	8
		Age Adjusted Rate	6.5	2.2
American Indian	# Cases / Deaths	8	3	
	Age Adjusted Rate	30.8	10.3	
United States	Total	Age Adjusted Rate	7.4	2.3
	White	Age Adjusted Rate	7.4	2.1
	American Indian	Age Adjusted Rate	7.4	2.3

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population. US rates www.seer.cancer.gov. Source: South Dakota Department of Health

Figure 23: Cervix Uteri Cancer Stage at Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: Early stage of diagnosis clearly provides the best opportunity for cure. In South Dakota, 32% of the cases reported were diagnosed at localized stage. SEER reports that 46% of the cases diagnosed nationally were at the localized stage.

Incidence: The incidence rate in South Dakota was 8.7 and in the United States it was 7.4. Both nationally and in South Dakota cervical cancer was the third most common female genital tract malignancy. Invasive cervical cancer accounted for 0.8% of all cases reported and 1.7% of all females diagnosed with cancer in South Dakota in 2014. SEER incidence reports that 0.2% of cases were younger than 20 years of age.


Mortality: The death rate in South Dakota was 2.8 for cancer of the cervix uteri. In the United States, the rate was 2.3. The stage of disease at diagnosis affects the mortality rate. Cases diagnosed at a localized stage have a 92% survival rate according to the American Cancer Society. Nationally, when diagnosed at distant stage, the percentage of survival drops to 17% at five years. In South Dakota, there were six cases in 2014 diagnosed at distant stage.

Risk and Associated Factors: Risk factors associated with cervical cancer suggest that a sexually transmitted agent is involved in the pathogenesis of the disease. In the last decade the Human Papilloma virus (HPV) has been identified as the most likely cause. Among the other risk factors are nutritional deficiencies (Vitamin C and Vitamin B), low socioeconomic status, being sexually active at a young age, high-risk male partner, tobacco use as well as the use of oral contraceptives.

Prevention and Early Detection: The US Preventive Services Task Force (USPSTF) recommends screening for cervical cancer in women age 21 to 65 years with cytology (Pap smear) every 3 years or, for women age 30 to 65 years who want to lengthen the screening interval, screening with a combination of cytology and human papillomavirus (HPV) testing every 5 years.

For more information on cervical cancer visit <http://www.cancer.gov/cancertopics/types/cervical/>.

Figure 24: Cervix Uteri Cancer Number of Cases and Deaths by Age, South Dakota, 2014



Source: South Dakota Department of Health

Half of the incidence of cervical cancer occurred in women under the age of 50 .




Figure 25: Cervix Uteri Cancer Cases and Deaths by Year, South Dakota, 2001 - 2014

The incidence peak for female cervix uteri cancer was in 2014.

Source: South Dakota Department of Health




Figure 26: Cervix Uteri Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014

Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health


COLORECTAL

Table 15: Colorectal Incidence and Mortality Summary, 2014

Colorectal Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	447	247	200	188	110	78
		Age Adjusted Rate	44.2	53.0	37.0	17.6	23.5	13.2
	White	# Cases / Deaths	413	231	182	176	100	76
		Age Adjusted Rate	44.1	53.4	36.4	17.4	22.6	13.6
	American Indian	# Cases / Deaths	24	10	14	10	9	1
		Age Adjusted Rate	51.1	43.8	55.8	21.8	39.7	6.6
United States	Total	Age Adjusted Rate	38.9	44.7	34.0	14.1	16.9	11.9
	White	Age Adjusted Rate	38.4	44.2	33.5	13.8	16.4	11.7
	American Indian	Age Adjusted Rate	41.2	44.3	39.0	15.2	20.0	11.6

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population.
US rates www.seer.cancer.gov Source: South Dakota Department of Health

Figure 27: Colorectal Cancer Stage at Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: The prognosis of the patient is greatly influenced by the stage of disease at diagnosis. In 2014, 36% (167) of the cases of colorectal cancer were diagnosed at localized stage. Localized is defined as when the disease is still confined to the colon. The remaining 258 invasive cases (55%) were diagnosed after the disease had spread beyond the colon. Of those 258 cases, 91 were diagnosed at distant stage when the disease had spread further involving other organs. The SEER National Cancer Institute website states that the 5-year survival rate for those who have distant stage at diagnosis is 13.9% for the 2007-2013 time period.


Incidence: Colorectal cancer accounted for 9.9% of all cases reported in South Dakota in 2014. The median age at diagnosis was 70. There were 247 men and 200 women diagnosed with colorectal cancer in 2014 in South Dakota. Overall, colorectal cancer was the fourth most diagnosed cancer. When reviewed by gender, it was the third most diagnosed cancer with 10.3% of the cancers reported in males and 9.5% of the cancers reported in females.

Mortality: Overall incidence and mortality rates for colorectal cancer are decreasing. The overall 5-year survival rate for 2007-2013 from SEER was 64.9% for men and women. In 2014, there were a total of 166 deaths that were attributed to colorectal cancer in South Dakota; more than half were men. Of that number, 176 were white and 10 were American Indian. The median age at death was 75. The SEER National Cancer Institute website states that the United States mortality rate was 14.1.


Risk and Associated Factors: Risk for colorectal cancer increases with age. Over 90% of cases occur in people who are 50 years old or older. Lifestyle factors that may contribute to an increased risk include: lack of regular physical activity, a diet low in fruit and vegetables, a low-fiber and high-fat diet, overweight and obesity, alcohol consumption, and tobacco use.

Prevention and Early Detection: The USPSTF recommends screening for colorectal cancer using high-sensitivity fecal occult blood testing (FOBT), sigmoidoscopy with interval FOBT, or colonoscopy for adults beginning at age 50 years and continuing until age 75 years. The risks and benefits of these screening methods vary.

Figure 28: Colorectal Cancer Number of Cases and Deaths by Age, South Dakota, 2014



Source: South Dakota Department of Health



Source: South Dakota Department of Health

Figure 29: Colorectal Cancer Cases and Deaths by Year, South Dakota, 2001 - 2014

The incidence peak for colorectal cancer occurred in 2002. Incidences of colorectal cancer appear to escalate after age 50.




Figure 30: Colorectal Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014

Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health


CORPUS and UTERUS, NOS

Table 16: Corpus and Uterus, NOS Incidence and Mortality Summary, 2014

Corpus and Uterus, NOS Cancer			Incidence	Mortality
South Dakota	Total	# Cases / Deaths	147	24
		Age Adjusted Rate	28.9	4.2
	White	# Cases / Deaths	134	24
		Age Adjusted Rate	28.4	4.4
	American Indian	# Cases / Deaths	5	0
		Age Adjusted Rate	16.4	0.0
United States	Total	Age Adjusted Rate	26.4	4.7
	White	Age Adjusted Rate	27.1	4.4
	American Indian	Age Adjusted Rate	16.2	3.3

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population. US rates www.seer.cancer.gov Source: South Dakota Department of Health

Figure 31: Corpus and Uterus, NOS Cancer Stage at Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: Cancer in the uterus is treated surgically. Staging for these diseases is done following surgery, unless it is obvious that the disease has progressed and advanced. Cases with obvious advanced disease do not benefit from surgical procedures and are staged by physical examination. These cases are treated without operative staging. In South Dakota, during 2014, 73% of corpus uteri cases were diagnosed at localized stage. Eleven cases were diagnosed at distant stage, less than in 2013.

Incidence: The uterine cervix is the small cylindrical neck that leads from the uterus, or womb, into the vagina. A knob of the cervix protrudes into the vagina and can be visualized on physical examination. It is lined with epithelial and stromal cells creating a site for epithelial, stromal and mixed cell malignancies. Endometrial carcinoma is one of the female genital cancers.


It is ranked fourth among females reported with cancer in South Dakota in 2014. Cancer of the corpus uteri represented 7.0% of all of the cancers diagnosed in South Dakota females in 2014. Endometrial cancer affects primarily postmenopausal women. The median age at diagnosis in the United States is 65. In South Dakota, the median age is 61 years of age.

Mortality: The death rate in South Dakota for the reporting period was 4.2 for deaths attributed to uterine cancer. In the United States the 2014 rate was 4.7. South Dakota had 24 female deaths attributed to cancer of the uterus in 2014. The stage of disease at diagnosis affects the mortality rate. Overall (all stages included), the five-year relative survival rate was 81.3% in the United States.

Risk and Associated Factors: Risk factors associated with corpus uteri cancer suggest that exposure to estrogen for long periods of time play a critical role. The use of exogenous estrogen replacement therapy accounted for a dramatic rise in the incidence of endometrial cancer in the United States in the 1970s. The use of combination estrogen-progesterone oral contraceptive pills confers protection against endometrial hyperplasia and subsequent development of cancer.

Prevention and Early Detection: Other factors associated with an increased risk of developing uterine cancer include obesity, a high-fat diet and a prolonged exposure to the female hormone, estrogen. One pregnancy appears to lower the risk of uterine cancer by 50%.

Figure 32: Corpus and Uterus, NOS Cancer Number of Cases and Deaths by Age, South Dakota, 2014



Source: South Dakota Department of Health

In South Dakota, in 2014 the incidence peaked in the 60-64 age group.




Figure 33: Corpus and Uterus, NOS Cancer Cases and Deaths by Year, South Dakota, 2001 - 2014

The incidence peak for female corpus and uterus, NOS cancer was in 2013.

Source: South Dakota Department of Health




Figure 34: Corpus and Uterus, NOS Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014

Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health


KIDNEY AND RENAL PELVIS

Table 17: Kidney and Renal Pelvis Incidence and Mortality Summary, 2014

Kidney and Renal Pelvis Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	175	125	50	40	26	14
		Age Adjusted Rate	17.9	27.0	9.7	3.7	5.3	2.2
	White	# Cases / Deaths	150	108	42	37	23	14
		Age Adjusted Rate	16.7	25.4	8.9	3.6	5.0	2.4
	American Indian	# Cases / Deaths	18	11	7	3	3	0
		Age Adjusted Rate	38.8	53.0	27.4	4.8	9.9	0.0
United States	Total	Age Adjusted Rate	15.6	21.4	10.6	3.8	5.6	2.3
	White	Age Adjusted Rate	16.1	22.0	10.9	3.9	5.7	2.3
	American Indian	Age Adjusted Rate	13.5	17.3	10.7	5.3	8.0	3.1

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population.
US rates www.seer.cancer.gov Source: South Dakota Department of Health

Figure 35: Kidney and Renal Pelvis Cancer Stage at Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: As with all malignancies, early diagnosis is the key to better prognosis and possible cure. In 2014 52% of the cases were diagnosed at localized stage, with another 20% diagnosed at distant stage. Unfortunately, symptoms do not always reflect the stage of disease. Blood in the urine is one of the symptoms that frequently presents at diagnosis. As with other cancers, renal cancer can spread through the blood stream and/or lymphatic system. Survival rates associated with kidney cancer depend on how far the disease has progressed, the size of tumor, and whether or not it has metastasized. The latest five-year survival rate for localized stage kidney cancer is 92.6%. The survival rate for distant stage is 11.7%.


Incidence: In 2014 the American Cancer Society estimated there would be 63,920 new cases of kidney cancer in the United States. This accounts for 3.9% of all reported malignancies in the United States. In South Dakota there were 175 reported cases of kidney cancer in 2014 representing 3.9% of all cancer cases. Kidney cancer develops most often in people over 40. There is no known cause of this disease. Doctors can seldom explain why one person develops kidney cancer and another does not. The median age at diagnosis is 65 in South Dakota and 64 in the United States.

Mortality: This cancer was the 14th leading cause of cancer death for South Dakota in 2014. In the United States for 2009-2013 it was the 12th leading cause of death with a median age of death of 71 years. Death rates decreased by 0.7% per year from 2004 to 2013.

Risk and Associated Factors: Cigarette smoking increases the risk of developing kidney cancer. The risk seems to increase by the amount one smokes. Obesity is associated with risk, as are exposures to occupational substances such as aniline dyes, benzene, and naphthalene.


Prevention and Early Detection: The main preventive measure is to stop smoking and maintain a healthy weight. It is difficult to diagnose kidney cancer until it becomes symptomatic. There are no known screenings recommended at this time.

Figure 36: Kidney and Renal Pelvis Cancer Number of Cases and Deaths by Age, South Dakota, 2014



Source: South Dakota Department of Health


Figure 37: Kidney and Renal Pelvis Cancer Cases and Deaths by Year, South Dakota, 2001 - 2014



Source: South Dakota Department of Health

The incidence peak for kidney and renal pelvis cancer occurred in 2014.

Figure 38: Kidney and Renal Pelvis Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health

LEUKEMIA

Table 18: Leukemia Incidence and Mortality Summary, 2014

Leukemia			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	153	90	63	90	51	39
		Age Adjusted Rate	15.2	19.0	12.5	8.8	11.5	6.6
	White	# Cases / Deaths	147	89	58	89	51	38
		Age Adjusted Rate	15.8	20.2	12.5	9.2	12.2	6.9
	American Indian	# Cases / Deaths	3	0	3	1	0	1
		Age Adjusted Rate	5.6	0.0	10.6	2.0	0.0	3.6
United States	Total	Age Adjusted Rate	13.5	17.2	10.5	6.6	8.8	5.0
	White	Age Adjusted Rate	14.1	18.0	11.0	6.9	9.2	5.2
	American Indian	Age Adjusted Rate	7.6	10.8	4.8	4.0	4.9	3.2

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population.
US rates www.seer.cancer.gov Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: Leukemias are not staged because they may involve bone marrow throughout the body. Doctors classify them by type and subtype in an attempt to determine the prognosis and a recommended level of treatment. Chronic myelogenous leukemia is grouped by phases and chronic lymphocytic leukemia (CLL) uses a Rai classification. Leukemia is a type of cancer of the blood. It is defined by how quickly the disease progresses. Leukemia is either chronic (disease progresses slowly) or acute (disease progresses quickly).

Incidence: Leukemias are a diverse group of cancers and are subtyped by histology. Subtypes have different etiology, treatment, and prognosis. Leukemias accounted for 3.4% of the cancers reported in 2014 for South Dakota. The American Cancer Society estimated that there would be 160 new cases of leukemia in South Dakota during 2014 and 52,380 cases nationwide.

Mortality: Leukemia accounted for 5.4% of the cancer deaths in South Dakota in 2014. The subtype of acute myeloid leukemia was the most frequent cause of leukemia death. Over 74% of the deaths associated with leukemia occurred at the age of 65 or older.

Leukemia is clinically and pathologically subdivided into a variety of large groups. The first division is between the acute and chronic forms.

Chronic Leukemia: Early in the disease process, the abnormal blood cells still have normal processes. Slowly, chronic leukemia does get worse. It causes symptoms as the number of abnormal cells in the blood rises. In South Dakota in 2014, there were 92 new cases of chronic leukemia.


Acute Leukemia: The blood cells are very abnormal. The blood cells cannot carry out their normal processes. The number of abnormal cells increases rapidly. Acute leukemia worsens quickly as do the symptoms. There were 50 new cases of acute leukemia in South Dakota in 2014.

These types of leukemia are further divided by the type of white blood cell that is affected.

Risk and Associated Factors: People who are exposed to very high levels of radiation are more likely to develop leukemia. Working with certain chemicals and exposure to high levels of benzene in the workplace can cause leukemia. Benzene is used widely in the chemical industry. Workers exposed to formaldehyde may also be at greater risk of leukemia. Chromosomal abnormalities, such as Down's syndrome and certain other genetic diseases may increase the risk of leukemia.

Prevention and Early Detection: There are no early detection or prevention strategies. Often symptoms are the same as for many other health problems, thus early detection is difficult. Diagnosis is made using blood tests and bone marrow biopsies.


Figure 39: Leukemia Number of Cases and Deaths by Age, South Dakota, 2014



Source: South Dakota Department of Health

Leukemia is frequently diagnosed in children, but as with all malignancies it is a disease of the elderly. Deaths from leukemia increase after the age of 64.


Figure 40: Leukemia Cases and Deaths by Year, South Dakota, 2001 - 2014



Source: South Dakota Department of Health

The incidence peak for leukemia occurred in 2011.

Figure 41: Leukemia Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health


LUNG AND BRONCHUS

Table 19: Lung and Bronchus Incidence and Mortality Summary, 2014

Lung and Bronchus Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths Age Adjusted Rate	582 56.4	327 69.0	255 47.4	439 41.9	255 54.1	184 33.3
	White	# Cases / Deaths Age Adjusted Rate	544 56.0	306 68.0	238 47.3	406 40.9	236 52.4	170 32.8
	American Indian	# Cases / Deaths Age Adjusted Rate	33 97.5	17 129.6	16 76.1	33 95.6	19 145.3	14 62.4
United States	Total	Age Adjusted Rate	53.1	61.5	46.7	42.2	52.0	34.7
	White	Age Adjusted Rate	54.5	61.7	49.0	43.0	52.1	36.0
	American Indian	Age Adjusted Rate	30.2	39.5	22.8	36.0	44.3	29.5

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population.
US rates www.seer.cancer.gov Source: South Dakota Department of Health

Figure 42: Lung and Bronchus Cancer Stage at Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: The presentation of lung cancer is extremely variable and depends on local manifestations of the tumor, distant metastases or associated paraneoplastic syndromes. In 2014, 21% of lung cancer patients were diagnosed at localized stage. The more advanced the stage, the poorer the prognosis is for the patient. In 2014, 309 (53%) cases were diagnosed when disease had progressed beyond the lung and metastasized to a distant location. Approximately 74% of cases in 2014 were diagnosed after the disease had progressed beyond the lung to lymph nodes, regional areas, or distant sites, such as brain or bone.

Incidence: Lung cancer is a major public health concern, with an estimated 224,210 new cases in


the United States in 2014. Despite the well documented link between tobacco product use and respiratory diseases, including cancer, the outcomes of such efforts to curb the use of tobacco products have been mixed. In South Dakota, there were 592 new invasive lung cancer cases diagnosed in 2014.

Mortality: There were 439 lung cancer deaths in South Dakota in 2014. Incidence and mortality rates have significantly increased during the last century. Lung cancer accounts for approximately 26.5% of all United States deaths attributed to cancer. In South Dakota, lung cancer accounts for 26.1% of deaths from cancer. Lung cancer is the leading cause of cancer deaths in both men and women.

Risk and Associated Factors: Cigarette smoking is the number one risk factor for lung cancer and is linked to approximately 90% of lung cancers. Other risk factors include secondhand smoke exposure, occupational or environmental exposures to substances such as radon, arsenic, benzene, and asbestos, a personal or family history of lung cancer, and previous radiation therapy to the chest.

Prevention and Early Detection: The USPSTF recommends annual screening for lung cancer with low-dose computed tomography (LDCT) in adults aged 55 to 80 years who have a 30 pack-year smoking history and currently smoke or have quit within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.

Figure 43: Lung and Bronchus Cancer Number of Cases and Deaths by Age, South Dakota, 2014



Source: South Dakota Department of Health




Figure 44: Lung and Bronchus Cancer Cases and Deaths by Year, South Dakota, 2001 - 2014

The number of cases and deaths associated with lung and bronchus cancer remain constant.

Source: South Dakota Department of Health




Figure 45: Lung and Bronchus Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014

Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health


MELANOMA OF THE SKIN

Table 20: Melanoma of the Skin Incidence and Mortality Summary, 2014

Melanoma of the Skin			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	239	127	112	31	22	9
		Age Adjusted Rate	25.8	27.0	25.6	3.3	5.3	1.7
	White	# Cases / Deaths	232	122	110	30	21	9
		Age Adjusted Rate	27.7	28.2	28.2	3.5	5.5	1.8
	American Indian	# Cases / Deaths	2	1	1	1	1	0
		Age Adjusted Rate	3.4	3.3	3.6	4.4	14.1	0.0
United States	Total	Age Adjusted Rate	23.6	30.7	18.4	2.6	3.8	1.6
	White	Age Adjusted Rate	28.2	36.1	22.4	3.0	4.4	1.8
	American Indian	Age Adjusted Rate	4.9	0.0	4.8	0.8	1.6	0.0

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population. US rates www.seer.cancer.gov Source: South Dakota Department of Health

Figure 46: Melanoma of the Skin Stage at Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: Melanoma is staged by the depth of invasion and the extension of the lesion. In 2014, almost half of the melanoma of the skin cases reported for South Dakota were localized. Another 43% were staged as *in situ* disease. The survival rate for localized melanoma is 98.4%. For distant disease, the survival rate is 17.9% at 5 years.

Incidence: In the United States in 2014, the American Cancer Society estimated that there would be 76,100 new cases of melanoma of the skin. There are three forms of skin cancer: basal cell, squamous cell, and melanoma. Melanoma is by far the most dangerous form of skin cancer.

Melanoma is primarily a cancer of the white populations. In 2014, South Dakota's incidence rate was 25.8 and the United States had an incidence rate of 23.6.

Mortality: There were 31 deaths attributed to melanoma of the skin in South Dakota in 2014 with a mortality rate of 3.3 and the United States mortality rate was 2.6. The median age for death in South Dakota for this cancer was 72 in 2014. Nationwide, the median age at death was 70 for melanoma of the skin.


Risk and Associated Factors: Certain factors are more likely to contribute to a higher risk. These are:

- Lighter natural skin color
- Family history of skin cancer
- Personal history of skin cancer
- Exposure to the sun
- History of sunburns early in life
- Skin that burns, freckles, reddens easily
- Blue or green eyes, blond or red hair
- Large number of moles

Early Detection and Prevention: The best way to prevent skin cancer is to protect the skin from the sun. The CDC recommends five easy options for protection from sunburn:


- use sun glasses
- use sunscreen regularly
- stay in the shade
- cover skin
- wear a hat

Figure 47: Melanoma of the Skin Number of Cases and Deaths by Age, South Dakota, 2014




Source: South Dakota Department of Health

Figure 48: Melanoma of the Skin Cases and Deaths by Year, South Dakota, 2001 - 2014



Source: South Dakota Department of Health

Figure 49: Melanoma of the Skin Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health

Table 21: Myeloma Incidence and Mortality Summary, 2014

Myeloma §			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	47	35	12	27	18	9
		Age Adjusted Rate	4.6	7.6	2.4	2.6	3.9	1.5
	White	# Cases / Deaths	42	32	10	26	18	8
		Age Adjusted Rate	4.4	7.4	2.0	2.6	4.1	1.3
	American Indian	# Cases / Deaths	4	2	2	1	0	1
		Age Adjusted Rate	6.9	6.9	7.0	2.9	0.0	5.0
United States	Total	Age Adjusted Rate	6.4	8.0	5.1	3.3	4.2	2.7
	White	Age Adjusted Rate	5.9	7.5	4.5	3.1	4.0	2.4
	American Indian	Age Adjusted Rate	4.9	0.0	5.5	3.6	3.3	3.8

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population. § can include NOS, multiple, plasma cell and solitary. US rates www.seer.cancer.gov Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: Stage of disease for myeloma is always distant per the SEER Summary Staging Manual.

Incidence: Myeloma is a systemic malignancy of plasma cells that is highly treatable, but rarely curable. It is potentially curable when it presents as a solitary plasmacytoma of the bone or as an extramedullary plasmacytoma. In South Dakota during 2014, myeloma accounted for 1.6% of total cancer cases reported. Median age at diagnosis in South Dakota was 73 and the United States was 69. The national incidence rate is higher in men (8.0) than women (5.1). In South Dakota the incidence rate is also higher in men (7.6) than women (2.4). Myeloma is more common among the elderly. African Americans have approximately twice the incidence and mortality rates of whites.


Mortality: The median survival prior to the common use of chemotherapy was about seven months. After the introduction of chemotherapy, prognosis improved significantly with a median survival of 24 to 30 months and a 10-year survival of 3%. During 2014, there were 27 deaths attributed to myeloma in South Dakota. Eighteen were male and nine were female. The mortality rate for South Dakota was 2.6. The rate for men was 3.9 and 1.5 for women. These

rates compare to United States mortality rates in 2014 of 3.3 overall, 4.2 for men and 2.7 for women.


Risk and Associated Factors: The etiology of myeloma is relatively unknown. There are many research studies evaluating the exposure of individuals with myeloma to various substances.

Early Detection and Prevention: There is no known test for screening for early detection. Some cases of myeloma progress very slowly, and they are referred to as smoldering or indolent myeloma. The presence of plasma cells and proteinuria do not automatically lead to myeloma, but it can be an early symptom. This disease is often asymptomatic in early stages of the disease. Myeloma is most often diagnosed clinically by radiological procedures and through cytology.

Figure 50: Myeloma Number of Cases and Deaths by Age, South Dakota, 2014




Source: South Dakota Department of Health



Source: South Dakota Department of Health

Figure 51: Myeloma Cancer Cases and Deaths by Year, South Dakota, 2001 - 2014

The incidence count for myeloma cancers took a sharp drop from 2004 to 2005 and 2013 to 2014, with an all-time high in 2011.



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health

Figure 52: Myeloma Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014

NON-HODGKIN'S LYMPHOMA


Table 22: Non-Hodgkin's Lymphoma Incidence and Mortality Summary, 2014

Non-Hodgkin's Lymphoma			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	188	109	79	52	34	18
		Age Adjusted Rate	18.9	24.1	14.4	4.9	7.4	3.1
	White	# Cases / Deaths	180	102	78	49	33	16
Age Adjusted Rate		19.6	24.4	15.4	4.9	7.5	3.0	
American Indian	# Cases / Deaths	6	5	1	2	1	1	
	Age Adjusted Rate	15.2	33.6	2.9	5.4	10.1	2.9	
United States	Total	Age Adjusted Rate	19.3	23.6	15.9	5.7	7.3	4.4
		White	20.2	24.7	16.6	5.9	7.6	4.6
	American Indian	Age Adjusted Rate	10.4	10.6	9.9	3.9	5.0	3.1

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population.

US rates www.seer.cancer.gov Source: South Dakota Department of Health

Figure 53: Non-Hodgkin's Lymphoma Stage at Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: Stage is based on where lymphoma cells are found (in the lymph or in other organs or tissues). The stage also depends on how many areas are involved. Localized stage only involves a single lymph node region or single extralymphatic organ. When two or more lymph node regions are involved and the regions are on both sides of the diaphragm the cancer is staged as distant. In 2014, 53% of the cases were diagnosed at distant stage, an increase from 2011 when 47% were diagnosed at distant stage.

Incidence: Non-Hodgkin's lymphoma is a cancer that originates in the lymphatic system, the disease-fighting network that spreads throughout the body. It develops in lymphocytes, a type of white blood cell. Non-Hodgkin's lymphoma is more than five times as common as the other general type of lymphoma,


Hodgkin's disease. The incidence rate has been increasing in the United States since the 1970s. The incidence of non-Hodgkin's disease in South Dakota was 188 cases in 2014. The median age at diagnosis in South Dakota in 2014 was 71.

Mortality: There were 52 deaths reported in South Dakota that were attributed to non-Hodgkin's lymphoma. The median age at death for those whose death was attributed to non-Hodgkin's lymphoma in South Dakota was 82 years of age. Nationally, the five-year survival rate is 71.0% for non-Hodgkin's lymphoma.


Risk and Associated Factors: Age is a strong risk factor for this disease, with most cases occurring from age 60 and older. Some studies suggest that exposure to chemicals such as benzene and certain herbicides and insecticides may be linked to an increased risk. Some chemotherapy used to treat other cancers can increase the risk as well as patients having been treated with radiation. The risk is higher for those having been treated with both. Certain infections increase the risk, such as HIV, Epstein - Barr virus, H. pylori bacteria, and Hepatitis C virus.

Early Detection and Prevention: Non-Hodgkin's lymphoma may present with various symptoms. Symptoms may include signs resulting from local effects of cancer growth. Non-Hodgkin's lymphoma can also produce generalized symptoms, such as unexplained weight loss, fever, drenching night sweats, and severe itching.

Figure 54: Non-Hodgkin's Lymphoma Number of Cases and Death by Age, South Dakota, 2014




Source: South Dakota Department of Health



Source: South Dakota Department of Health

Figure 55: Non-Hodgkin's Lymphoma Cases and Deaths by Year, South Dakota, 2001 – 2014



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health


Figure 56: Non-Hodgkin's Lymphoma Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014

Table 23: Ovarian Incidence and Mortality Summary, 2014

Ovarian Cancer			Incidence	Mortality
South Dakota	Total	# Cases / Deaths	55	41
		Age Adjusted Rate	5.4	4.1
	White	# Cases / Deaths	51	38
		Age Adjusted Rate	5.4	4.0
	American Indian	# Cases / Deaths	3	2
		Age Adjusted Rate	4.5	5.5
United States	Total	Age Adjusted Rate	11.3	7.0
	White	Age Adjusted Rate	11.7	7.3
	American Indian	Age Adjusted Rate	8.6	5.4

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population. US rates www.seer.cancer.gov Source: South Dakota Department of Health

Figure 57: Ovarian Cancer Stage at Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: Staging of ovarian cancer is done by a surgical procedure to remove as much of the cancer as possible. Surgical staging is of critical importance in management of this disease. The morbidity associated with ovarian carcinoma is partially attributable to the fact that in the United States two-thirds of the patients present with advanced-stage disease at the time of diagnosis. In 2014, in South Dakota 30 (54%) of the 55 cases were diagnosed at distant stage.

Incidence: The incidence of ovarian cancer varies greatly. There were 55 cases of ovarian cancer reported in 2014 in South Dakota. This accounted for 2.6% of the cancer cases diagnosed in 2014 for South Dakota women. The lifetime risk of a woman developing ovarian cancer is 1.4%. Seven cases were diagnosed at

younger than 49 years of age. There were 14 cases diagnosed in the 60-69 age group. The median age at diagnosis in South Dakota was 64; nationally it was 63.


Mortality: Doctors are using dramatic new therapies to fight ovarian cancer, extending the lives of women who five or 10 years ago would have died from the disease. Survival rates for the last several decades are only about 25% for those with advanced disease. Most ovarian cancer presents at advanced disease. Only 28.9% of those diagnosed at late stage survive five years. For those who are diagnosed early, before the disease spreads beyond the ovaries, the disease is 92.5% curable. In South Dakota, 41 patients died in 2014. The mortality rate was 4.1 for women in South Dakota.

Risk and Associated Factors: Women who have a higher risk for developing ovarian cancer are those with a family history of the disease, those who have used fertility drugs, those who had their first baby after age 30 and those over the age of 65.

Prevention and Early Detection: Factors that may reduce the risk of ovarian cancer are pregnancy lasting full term, use of oral contraceptives, breastfeeding, tubal ligation, hysterectomy, or removal of ovaries in women with inherited risks.


No early stage screening tests have been proven for ovarian cancer and it can be difficult to detect until it has advanced. A combination of imaging and lab tests are the most useful diagnostic tools.

Figure 58: Ovarian Cancer Number of Cases and Deaths by Age, South Dakota, 2014




Source: South Dakota Department of Health

Figure 59: Ovarian Cancer Cases and Deaths by Year, South Dakota, 2001 - 2014



Source: South Dakota Department of Health

Figure 60: Ovarian Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health


PANCREAS

Table 24: Pancreas Incidence and Mortality Summary, 2014

Pancreas Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	131	73	58	118	76	42
		Age Adjusted Rate	12.9	15.4	10.6	11.1	16.1	6.7
	White	# Cases / Deaths	120	65	55	114	73	41
		Age Adjusted Rate	12.6	14.8	10.4	11.4	16.4	6.9
	American Indian	# Cases / Deaths	6	4	2	3	2	1
		Age Adjusted Rate	11.2	17.7	6.9	6.2	10.8	3.3
United States	Total	Age Adjusted Rate	12.6	14.3	11.1	10.9	12.6	9.5
	White	Age Adjusted Rate	12.5	14.3	11.0	10.9	12.6	9.4
	American Indian	Age Adjusted Rate	9.1	11.4	7.7	8.5	9.9	7.5

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population.
US rates www.seer.cancer.gov Source: South Dakota Department of Health

Figure 61: Pancreatic Cancer Stage at Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: Often pancreatic cancer is diagnosed late in the disease process. Patients who have local stage disease may be acceptable for resection. Only 10 to 20% of patients with pancreatic cancer are candidates for surgical resection. In South Dakota, 76% of new cases were diagnosed at late stage (regional and distant) in 2014.

Incidence: The incidence of pancreatic cancer increases steadily with age. An estimated 46,420 new cases of pancreatic cancer were expected to be diagnosed in 2014 in the United States. The majority of the cases occurred in South Dakotans 65 years old or older. Ninety-two cases (70.2%) were diagnosed in 2014 in that age group. In the United States and South Dakota this cancer


occurred more in males than in females. Nationally, there was a higher incidence rate in blacks of both genders. The median age at diagnosis was 71 years in South Dakota and 70 in the United States.

Mortality: The overall survival for cancer of the pancreas is poor. Studies reveal that the 5-year survival rate is approximately 8.2%. More recently, prospective studies show survival improvement with postoperative chemotherapy. In 2014, there were 118 deaths and the median age at death was 74 in South Dakota.

Risk and Associated Factors: The exact causes of pancreatic cancer are unknown. Studies have found that certain factors increase a person's risk for developing pancreatic cancer. As one ages, the incidence of pancreatic cancer increases, especially after the age of 60. Cigarette smokers are two to three times more likely than nonsmokers to develop this cancer. Pancreatic cancer occurs frequently in those with diabetes. Also, African Americans are more likely than Asians, Hispanics, American Indians, or whites to have pancreatic cancer. The risk triples if the person's mother, father, sister, or brother had the disease. Also, a history of colon or ovarian cancer increases the risk. Some evidence shows that chronic pancreatitis may increase the risk.


Prevention and Early Detection: Currently, there are no known screenings for pancreatic cancer. Also, there is no specific prevention except to avoid smoking.

Figure 62: Pancreatic Cancer Number of Cases and Deaths by Age, South Dakota, 2014




Source: South Dakota Department of Health

Figure 63: Pancreatic Cancer Cases and Deaths by Year, South Dakota, 2001 - 2014



Source: South Dakota Department of Health

Figure 64: Pancreatic Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health


PROSTATE

Table 25: Prostate Incidence and Mortality Summary, 2014

Prostate Cancer			Incidence	Mortality
South Dakota	Total	# Cases / Deaths	549	75
		Age Adjusted Rate	107.7	17.0
	White	# Cases / Deaths	512	71
		Age Adjusted Rate	107.4	16.7
	American Indian	# Cases / Deaths	31	4
		Age Adjusted Rate	143.5	25.9
United States	Total	Age Adjusted Rate	97.6	19.1
	White	Age Adjusted Rate	90.8	17.9
	American Indian	Age Adjusted Rate	50.4	20.1

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population. US rates www.seer.cancer.gov Source: South Dakota Department of Health

Figure 65: Prostate Cancer Stage at Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: The greatest number of cases was diagnosed at an early stage. In 2014, 82% of the cases were diagnosed as localized (not extending outside the prostate). Frequently older cases may simply be monitored (watchful waiting) by their physician to assess the rate of growth; others may be given hormonal therapy. New treatments for prostate cancer include the de Vinci Robotic assisted prostatectomy, proton therapy, and brachytherapy radiation.

Incidence: Carcinoma of the prostate is predominately a tumor of older men. The median age at diagnosis in South Dakota is 66. Also, in South Dakota the incidence of prostate cancer begins to increase in the 60's age group. Nationwide, eight out of 10 men diagnosed with prostate cancer are over the age of 65. Prostate


cancer is the third most diagnosed site of all cancers reported in the state.

Mortality: Prostate cancer was the fourth leading cancer death in men in South Dakota in 2014. Prostate cancer can be a slow progressing disease and can be cured or at least controlled in the early stages. The median age of death in South Dakota in 2014 was 80 years old. Many patients have co-morbid conditions and will die of other causes rather than prostate cancer.

Risk and Associated Factors: A number of risk factors for prostate cancer have been identified. Studies suggest that prostate cancer risk is increased two to five-fold in relatives of men with prostate cancer. Environmental factors, including exposure to heavy metals may increase risk. Smoking has also been indicated as a risk. Diets high in saturated fat intake may also contribute.


Early Detection and Prevention: The availability of the Prostate Specific Antigen (PSA) test as a diagnostic test coupled with increased awareness by the public of the disease has produced an increase in the number of new cases diagnosed each year in the United States. Disease detected by elevated PSA in the presence of a palpable normal gland is the most common presentation of prostate cancer. The American Cancer Society recommends the PSA and digital rectal exam should be offered annually beginning at the age of 50.

Figure 66: Prostate Cancer Number of Cases and Deaths by Age, South Dakota, 2014




Source: South Dakota Department of Health

Figure 67: Prostate Cancer Cases and Deaths by Year, South Dakota, 2001 - 2014



Source: South Dakota Department of Health

Figure 68: Prostate Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota 2001 - 2014



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health


STOMACH

Table 26: Stomach Incidence and Mortality Summary, 2014

Stomach Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	46	28	18	24	11	13
		Age Adjusted Rate	4.4	5.7	3.3	2.2	2.3	2.2
	White	# Cases / Deaths	39	25	14	22	11	11
Age Adjusted Rate		4.1	5.5	3.0	2.1	2.4	1.8	
American Indian	# Cases / Deaths	6	3	3	2	0	2	
	Age Adjusted Rate	14.2	14.1	12.8	3.8	0.0	7.2	
United States	Total	Age Adjusted Rate	7.2	9.8	5.1	3.1	4.2	2.3
		White	Age Adjusted Rate	6.4	8.8	4.4	2.7	3.7
	American Indian	Age Adjusted Rate	10.2	13.0	7.8	6.0	8.7	3.9

Rates per 100,000 age-adjusted to 2000 US standard population and 2013 SD estimated population. US rates www.seer.cancer.gov Source: South Dakota Department of Health

Figure 69: Stomach Cancer Stage of Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: In 2014 data demonstrates that 10 (22%) cases were diagnosed at localized stage. When a patient is diagnosed at an early stage prognosis is much better. Eight cases (17%) were diagnosed at regional stage. There were 22 (48%) of the cases in South Dakota diagnosed at distant stage. Prognosis for distant stage is very poor. The stage is based on whether the tumor has invaded nearby tissues, where the cancer has spread, and if so, to what extent.

Incidence: Stomach cancer continues to account for approximately 1.0% of all cancers in South Dakota. Of the 46 cases diagnosed in 2014, 28 were male and 18 were female. It is

predominately a disease of men. Gastric (stomach) cancer is found more commonly in people between the ages of 50 and 70 years of age. The median age at diagnosis was 68 in the United States and 71 in South Dakota.

Mortality: Stomach cancer accounted for 1.4% of cancer deaths in South Dakota in 2014. The median age at death was 76.5 in South Dakota and 72 in the United States. The age-adjusted death rate was 2.3 for men and 2.2 in women in South Dakota. These rates are based on patients who died in 2014 in South Dakota. There were two American Indian stomach cancer deaths.


Risk and Associated Factors: Men have twice the risk of women for developing stomach cancer. In recent years, *Helicobacter pylori* bacteria have received considerable attention as a potential factor. Some researchers suspect this bacterium, which causes stomach inflammation and ulcers, may be an important stomach cancer risk factor. Individuals with pernicious anemia (a vitamin B-12-related disorder) and achlorhydria or gastric atrophy, both of which result in lower than normal amounts of gastric juices, may be at higher risk.

Prevention and Early Detection: Excessive salt intake has been identified as a possible risk factor for stomach cancer. Having a high intake of fresh fruits and vegetables may be associated with a decreased risk of stomach cancer. Studies have suggested that eating foods that contain **beta-carotene**¹ and **vitamin C**² may decrease the risk of stomach cancer.

¹<http://www.cancer.gov/Common/PopUps/popDefinition.aspx?id=45328&version=Patient&language=English>


²<http://www.cancer.gov/Common/PopUps/popDefinition.aspx?id=439435&version=Patient&language=English>

Figure 70: Stomach Cancer Number of Cases and Deaths by Age, South Dakota, 2014




Source: South Dakota Department of Health

Figure 71: Stomach Cancer Cases and Deaths by Year, South Dakota, 2001 - 2014



Source: South Dakota Department of Health

Figure 72: Stomach Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health

THYROID


Table 27: Thyroid Incidence and Mortality Summary, 2014

Thyroid Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	103	26	77	9	2	7
		Age Adjusted Rate	12.0	6.1	18.2	0.9	0.4	1.3
	White	# Cases / Deaths	99	26	73	7	2	5
		Age Adjusted Rate	13.1	6.8	19.6	0.7	0.4	1.0
	American Indian	# Cases / Deaths	2	0	2	1	0	1
		Age Adjusted Rate	3.5	0.0	6.5	1.6	0.0	3.3
United States	Total	Age Adjusted Rate	14.6	7.3	21.7	0.5	0.5	0.5
	White	Age Adjusted Rate	15.4	7.7	23.1	0.5	0.5	0.5
	American Indian	Age Adjusted Rate	9.7	6.1	13.4	0.0	0.0	0.0

Rates per 100,000 age-adjusted to 2000 US standard population and 2014 SD estimated population.

US rates www.seer.cancer.gov Source: South Dakota Department of Health

Figure 73: Thyroid Cancer Stage at Diagnosis, South Dakota, 2014



Source: South Dakota Department of Health

Descriptive Epidemiology

Stage at Diagnosis: In 2014 data demonstrates that 59 (57%) of cases were diagnosed at localized stage. When a patient is diagnosed at an early stage, prognosis is much better for a cure. There were 38 (37%) cases diagnosed at regional stage. Only three cases (3%) were diagnosed at a distant stage.

Incidence: The American Cancer Society estimated 62,980 thyroid cancer cases would be diagnosed in the United States in 2014. Thyroid cancer continues to account for approximately 2.3% of all cancers in South Dakota. Of the 103 cases diagnosed in 2014, 26 were male and 77 were female. The median age at diagnosis was 52. In the United States the median age was 51. Thyroid cancer is

found more commonly in people between the ages of 45 and 70 years of age, with 77% diagnosed before age 65. It is predominately a disease of females as the statistics for South Dakota confirm.


Mortality: South Dakota had nine deaths attributed to thyroid cancer in 2014. Nationally, the 5-year relative survival rates were 99.9% for localized, 98.0% for regional, and 88.6% for unknown stage.

Risk and Associated Factors: Thyroid cancer accounted for only 2.3% of the cancer cases in South Dakota in 2014. Risk factors include being exposed to radiation to the head and neck in childhood. Other risk factors for the development of thyroid cancer include a history of goiter, family history of thyroid disease, and Asian race.

Early Detection and Prevention: Early detection of cancer of the thyroid is extremely important. There are currently no tests or screenings for early detection of thyroid cancer. Physical examinations may reveal a lump on the side of neck, hoarseness of the voice, and difficulty swallowing. Most cancerous thyroid tumors are slow growing and curable. Prompt attention to signs and symptoms is the best approach to early diagnosis of most thyroid cancers. Signs or symptoms include:


- A lump in the neck, sometimes growing rapidly
- A pain in the neck, sometimes going up to the ears
- Hoarseness
- Trouble swallowing
- Breathing problems (feeling as if one were breathing through a straw)
- A cough that persists and is not due to a cold

Figure 74: Thyroid Cancer Number of Cases and Deaths by Age, South Dakota, 2014



Source: South Dakota Department of Health


Figure 75: Thyroid Cancer Cases and Deaths by Year, South Dakota, 2001 - 2014



The incidence count for thyroid cancers peaked in 2013.

Source: South Dakota Department of Health

Figure 76: Thyroid Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2014



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.
Source: South Dakota Department of Health

XI: APPENDICES

Appendix A. 2000 United States Standard Million Population

Age Group	Number in Group
Total	1,000,000
<5	69,135
5-9	72,533
10-14	73,032
15-19	72,169
20-24	66,478
25-29	64,529
30-34	71,044
35-39	80,762
40-44	81,851
45-49	72,118
50-54	62,716
55-59	48,454
60-64	38,793
65-69	34,264
70-74	31,773
75-79	26,999
80-84	17,842
85+	15,508

Appendix B. 2005-2014 South Dakota Estimated Population

Year	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Total	780,084	788,519	797,035	804,532	812,383	814,180	824,082	833,354	844,877	853,175
<5	54,334	55,640	57,448	58,582	59,640	59,621	59,591	59,202	59,957	60,610
5-9	51,478	51,545	51,944	52,692	53,496	55,531	56,389	58,010	59,832	60,148
10-14	54,880	54,450	53,630	52,954	52,576	53,960	54,096	53,956	54,899	55,807
15-19	60,843	60,084	59,858	59,184	58,571	57,628	57,916	57,439	57,419	57,823
20-24	62,742	63,756	62,601	62,764	62,919	57,596	58,178	59,174	60,849	61,697
25-29	47,482	48,972	51,977	53,851	56,270	55,570	56,020	56,397	56,103	55,859
30-34	44,406	43,985	44,439	45,387	46,540	49,859	52,216	53,875	55,411	56,075
35-39	46,103	46,184	46,287	46,003	45,707	45,766	45,524	46,326	47,452	49,007
40-44	56,131	54,090	51,639	49,163	47,626	47,346	47,375	47,570	47,255	46,709
45-49	58,361	58,701	58,653	58,526	57,845	57,519	54,849	52,681	50,182	48,446
50-54	53,782	55,507	56,682	57,673	57,850	59,399	59,960	60,037	59,699	58,976
55-59	44,626	47,667	49,558	51,199	52,996	54,231	56,261	57,577	58,434	58,768
60-64	34,232	35,504	38,305	40,441	42,615	43,573	47,054	48,927	51,222	53,027
65-69	27,699	28,381	29,312	30,956	32,303	31,944	33,144	35,738	37,753	40,473
70-74	24,567	24,588	24,670	24,879	25,236	25,683	25,882	26,396	27,834	28,875
75-79	22,441	22,634	22,574	22,287	21,866	21,724	21,660	21,766	21,939	22,209
80-84	17,826	18,043	18,162	18,253	18,122	18,004	18,102	18,112	18,017	17,795
85+	18,151	18,788	19,296	19,738	20,205	19,226	19,865	20,171	20,620	20,871

Appendix C: Race in South Dakota by County, 2014 Estimated Population

	Total		White		Black		American Indian		Asian		Other	
South Dakota	853,175	731,359	86%	15,971	2%	75,879	9%	10,739	1%	19,227	2%	
Aurora	2,745	2,635	96%	13	0%	54	2%	17	1%	26	1%	
Beadle	18,169	16,107	89%	315	2%	231	1%	1,153	6%	363	2%	
Bennett	3,430	1,192	35%	15	0%	2,016	59%	22	1%	185	5%	
Bon Homme	7,023	6,303	90%	91	1%	522	7%	13	0%	94	1%	
Brookings	33,314	30,970	93%	505	2%	441	1%	941	3%	457	1%	
Brown	38,408	35,019	91%	617	2%	1,267	3%	735	2%	770	2%	
Brule	5,309	4,617	87%	20	0%	489	9%	16	0%	167	3%	
Buffalo	2,077	383	18%	12	1%	1,634	79%	2	0%	46	2%	
Butte	10,298	9,723	94%	73	1%	209	2%	33	0%	260	3%	
Campbell	1,386	1,346	97%	4	0%	11	1%	7	1%	18	1%	
Charles Mix	9,287	5,940	64%	36	0%	3,013	32%	22	0%	276	3%	
Clark	3,645	3,550	97%	52	1%	8	0%	3	0%	32	1%	
Clay	13,932	12,665	91%	229	2%	423	3%	299	2%	316	2%	
Codington	27,938	26,499	95%	214	1%	631	2%	221	1%	373	1%	
Corson	4,182	1,348	32%	10	0%	2,675	64%	16	0%	133	3%	
Custer	8,445	7,879	93%	58	1%	283	3%	42	0%	183	2%	
Davison	19,885	18,715	94%	202	1%	551	3%	109	1%	308	2%	
Day	5,588	4,926	88%	35	1%	490	9%	34	1%	103	2%	
Deuel	4,312	4,205	98%	43	1%	16	0%	6	0%	42	1%	
Dewey	5,662	1,292	23%	21	0%	4,120	73%	10	0%	219	4%	
Douglas	2,973	2,856	96%	17	1%	72	2%	7	0%	21	1%	
Edmunds	3,983	3,902	98%	9	0%	23	1%	7	0%	42	1%	
Fall River	6,845	6,003	88%	77	1%	479	7%	60	1%	226	3%	
Faulk	2,357	2,318	98%	9	0%	4	0%	5	0%	21	1%	
Grant	7,241	6,998	97%	69	1%	63	1%	28	0%	83	1%	
Gregory	4,217	3,746	89%	16	0%	322	8%	18	0%	115	3%	
Haakon	1,847	1,719	93%	16	1%	38	2%	13	1%	61	3%	
Hamlin	5,989	5,864	98%	32	1%	29	0%	19	0%	45	1%	
Hand	3,345	3,288	98%	4	0%	13	0%	12	0%	28	1%	
Hanson	3,419	3,366	98%	7	0%	12	0%	12	0%	22	1%	
Harding	1,250	1,189	95%	7	1%	27	2%	3	0%	24	2%	
Hughes	17,642	14,983	85%	181	1%	1,945	11%	115	1%	418	2%	
Hutchinson	7,200	6,968	97%	53	1%	98	1%	15	0%	66	1%	
Hyde	1,396	1,231	88%	6	0%	129	9%	4	0%	26	2%	
Jackson	3,274	1,393	43%	1	0%	1,700	52%	2	0%	159	5%	
Jerauld	2,007	1,967	98%	4	0%	10	0%	5	0%	21	1%	
Jones	975	899	92%	3	0%	35	4%	8	1%	30	3%	
Kingsbury	5,075	4,930	97%	27	1%	35	1%	21	0%	62	1%	
Lake	12,368	11,858	96%	125	1%	103	1%	120	1%	162	1%	
Lawrence	24,657	23,148	94%	245	1%	570	2%	178	1%	516	2%	
Lincoln	51,548	49,165	95%	653	1%	280	1%	684	1%	766	1%	
Lyman	3,877	2,254	58%	19	0%	1,484	38%	11	0%	109	3%	
Marshall	4,683	4,113	88%	57	1%	434	9%	12	0%	0	0%	
McCook	5,649	5,532	98%	25	0%	37	1%	10	0%	2	0%	
McPherson	2,429	2,386	98%	10	0%	9	0%	4	0%	3	0%	
Meade	26,951	24,646	91%	519	2%	744	3%	252	1%	19	0%	
Mellette	2,100	873	42%	7	0%	1,100	52%	6	0%	0	0%	
Miner	2,316	2,255	97%	15	1%	12	1%	9	0%	0	0%	
Minnehaha	182,882	161,484	88%	8,482	5%	4,988	3%	3,495	2%	199	0%	
Moody	6,367	5,189	81%	57	1%	856	13%	81	1%	0	0%	
Oglala Lakota	14,218	835	6%	32	0%	13,109	92%	16	0%	3	0%	
Pennington	108,242	90,415	84%	1,704	2%	10,812	10%	1,314	1%	115	0%	
Perkins	3,033	2,935	97%	10	0%	51	2%	8	0%	0	0%	
Potter	2,340	2,252	96%	10	0%	42	2%	11	0%	0	0%	
Roberts	10,374	6,158	59%	63	1%	3,798	37%	33	0%	0	0%	
Sanborn	2,336	2,281	98%	3	0%	9	0%	5	0%	1	0%	
Spink	6,598	6,349	96%	47	1%	127	2%	7	0%	0	0%	
Stanley	2,983	2,642	89%	27	1%	230	8%	8	0%	0	0%	
Sully	1,438	1,367	95%	12	1%	28	2%	1	0%	0	0%	
Todd	9,882	1,072	11%	28	0%	8,523	86%	25	0%	0	0%	
Tripp	5,512	4,553	83%	22	0%	774	14%	28	1%	1	0%	
Turner	8,272	8,029	97%	42	1%	86	1%	20	0%	4	0%	
Union	15,029	14,362	96%	160	1%	108	1%	159	1%	13	0%	
Walworth	5,511	4,538	82%	29	1%	747	14%	20	0%	0	0%	
Yankton	22,684	21,044	93%	437	2%	689	3%	157	1%	9	0%	
Ziebach	2,826	690	24%	9	0%	2,011	71%	10	0%	1	0%	

US Census Bureau Estimated Race Population

Appendix D: SEER Incidence Site Analysis Categories

Site Group	ICD-O-3 Site	ICD-O-3 Histology (Type)	Recode	
Oral Cavity and Pharynx				
Lip	C000-C009	excluding 9050-9055, 9140, 9590-9992	20010	
Tongue	C019-C029		20020	
Salivary Gland	C079-C089		20030	
Floor of Mouth	C040-C049		20040	
Gum and Other Mouth	C030-C039, C050-C059, C060-C069		20050	
Nasopharynx	C110-C119		20060	
Tonsil	C090-C099		20070	
Oropharynx	C100-C109		20080	
Hypopharynx	C129, C130-C139		20090	
Other Oral Cavity and Pharynx	C140, C142, C148		20100	
Digestive System				
Esophagus	C150-C159	excluding 9050-9055, 9140, 9590-9992	21010	
Stomach	C160-C169		21020	
Small Intestine	C170-C179		21030	
Colon and Rectum				
Colon excluding Rectum				
Cecum	C180	excluding 9050-9055, 9140, 9590-9992	21041	
Appendix	C181		21042	
Ascending Colon	C182		21043	
Hepatic Flexure	C183		21044	
Transverse Colon	C184		21045	
Splenic Flexure	C185		21046	
Descending Colon	C186		21047	
Sigmoid Colon	C187		21048	
Large Intestine, NOS	C188-C189, C260		21049	
Rectum and Rectosigmoid Junction				
Rectosigmoid Junction	C199	excluding 9050-9055, 9140, 9590-9992	21051	
Rectum	C209		21052	
Anus, Anal Canal and Anorectum	C210-C212, C218		21060	
Liver and Intrahepatic Bile Duct				
Liver	C220	excluding 9050-9055, 9140, 9590-9992	21071	
Intrahepatic Bile Duct	C221		21072	
Gallbladder	C239		21080	
Other Biliary	C240-C249		21090	
Pancreas	C250-C259		21100	
Retroperitoneum	C480		21110	
Peritoneum, Omentum and Mesentery	C481-C482		21120	
Other Digestive Organs	C268-C269, C488		21130	
Respiratory System				
Nose, Nasal Cavity and Middle	C300-C301, C310-C319		excluding 9050-9055, 9140, 9590-9992	22010
Larynx	C320-C329	22020		
Lung and Bronchus	C340-C349	22030		
Pleura	C384	22050		
Trachea, Mediastinum and Other Respiratory Organs	C339, C381-C383, C388, C390, C398, C399	22060		

Appendix D: SEER Incidence Site Analysis Categories (continued)

Site Group	ICD-O-3 Site	ICD-O-3 Histology (Type)	Recode
Bones and Joints	C400-C419	excluding 9050-9055, 9140, 9590-9992	23000
Soft Tissue including Heart	C380, C470-C479, C490-C499	excluding 9050-9055, 9140, 9590-9992	24000
Skin excluding Basal and Squamous			
Melanoma of the Skin	C440-C449	8720-8790	25010
Other Non-Epithelial Skin	C440-C449	excluding 8000-8005, 8010-8046, 8050-8084, 8090-8110, 8720-8790, 9050-9055, 9140, 9590-9992	25020
Breast	C500-C509	excluding 9050-9055, 9140, 9590-9992	26000
Female Genital System			
Cervix Uteri	C530-C539	excluding 9050-9055, 9140, 9590-9992	27010
Corpus and Uterus, NOS			
Corpus Uteri	C540-C549	excluding 9050-9055, 9140, 9590-9992	27020
Uterus, NOS	C559		27030
Ovary	C569		27040
Vagina	C529		27050
Vulva	C510-C519		27060
Other Female Genital Organs	C570-C579, C589		27070
Male Genital System			
Prostate	C619	excluding 9050-9055, 9140, 9590-9992	28010
Testis	C620-C629		28020
Penis	C600-C609		28030
Other Male Genital Organs	C630-C639		28040
Urinary System			
Urinary Bladder	C670-C679	excluding 9050-9055, 9140, 9590-9992	29010
Kidney and Renal Pelvis	C649, C659		29020
Ureter	C669		29030
Other Urinary Organs	C680-C689		29040
Eye and Orbit	C690-C699	excluding 9050-9055, 9140, 9590-9992	30000
Brain and Other Nervous System			
Brain	C710-C719	excluding 9050-9055, 9140, 9530-9539, 9590-9992	31010
Cranial Nerves Other Nervous System	C710-C719	9530-9539	31040
	C700-C709, C720-C729	excluding 9050-9055, 9140, 9590-9992	
Endocrine System			
Thyroid	C739	excluding 9050-9055, 9140, 9590-9992	32010
Other Endocrine including Thymus	C379, C740-C749, C750-C759		32020

Appendix D: SEER Incidence Site Analysis Categories (continued)

Site Group	ICD-O-3 Site	ICD-O-3 Histology (Type)	Recode
Lymphoma			
Hodgkin's Lymphoma			
Hodgkin's - Nodal	C024,C098-C099,C111, C142,C379,C422,C770-C779	9650-9667	33011
Hodgkin's - Extranodal	All other sites		33012
Non-Hodgkin's Lymphoma			
NHL - Nodal	C024, C098, C099, C111, C142,C379, C422, C770-C779	9590-9597, 9670-9671, 9673, 9675, 9678-9680, 9684, 9687-9691, 9695, 9698-9702, 9705, 9708-9709, 9712, 9714-9719, 9724-9729, 9735, 9737-9738, 9811-9818, 9823, 9827, 9837	33041
NHL - Extranodal	All sites except C024, C098-C099, C111, C142, C379, C422, C770-C779	9590-9597, 9670-9671, 9673, 9675, 9678-9680, 9684, 9687, 9688, 9689-9691, 9695, 9698-9702,9705, 9708-9709, 9712, 9714-9719, 9724-9729, 9735, 9737,9738	33042
	All sites except C024, C098-C099, C111, C142, C379, C420-C422, C424, C770-C779	9811-9818, 9823, 9827, 9837	
Myeloma		9731-9732, 9734	34000
Leukemia			
Lymphocytic Leukemia			
Acute Lymphocytic Leukemia		9826, 9835-9836	35011
	C420, C421, C424	9811-9818, 9837	
Chronic Lymphocytic Leukemia	C420, C421, C424	9823	35012
Other Lymphocytic Leukemia		9820, 9832-9834, 9940	35013
Myeloid and Monocytic Leukemia			
Acute Myeloid Leukemia		9840, 9861, 9865-9867, 9869, 9871-9874, 9895-9897, 9898, 9910-9911, 9920	35021
Acute Monocytic Leukemia		9891	35031
Chronic Myeloid Leukemia		9863, 9875-9876, 9945-9946	35022
Other Myeloid/Monocytic Leukemia		9860, 9930	35023
Other Leukemia			
Other Acute Leukemia		9801, 9805-9809, 9931	35041
Aleukemic, subleukemic and NOS		9733, 9742, 9800, 9831, 9870, 9948, 9963-9964	35043
	C420, C421, C424	9827	
Mesothelioma +		9050-9055	36010
Kaposi Sarcoma +		9140	36020
Miscellaneous		9740-9741, 9750-9769, 9950, 9960-9962, 9965-9967, 9970-9971, 9975, 9980, 9982-9987, 9989, 9991-9992	37000
	C760-C768, C809	Excluding 9050-9055, 9140, 9590-9992	
	C420-C424 C770-C779		
Invalid	Site or histology code not within valid range or site code not found in this table.		99999

Source: <http://seer.cancer.gov/siterecode>

+ The Site Recode variable can be created with or without Mesothelioma (9050-9055) and Kaposi Sarcoma (9140) as separate groupings. The table above documents both possibilities.

Appendix E: SEER Cancer Cause of Death Analysis Categories

Cancer Causes of Death	ICD-10
All Malignant Cancers	C00-C97
Oral Cavity and Pharynx	
Lip	C00
Tongue	C01-C02
Salivary Gland	C07-C08
Floor of Mouth	C04
Gum and Other Mouth	C03, C05-C06
Nasopharynx	C11
Tonsil	C09
Oropharynx	C10
Hypopharynx	C12-C13
Other Oral Cavity and Pharynx	C14
Digestive System	
Esophagus	C15
Stomach	C16
Small Intestine	C17
Colon and Rectum	
Colon Excluding Rectum	C18, C26.0
Rectum and Rectosigmoid Junction	C19-C20
Anus, Anal Canal and Anorectum	C21
Liver and Intrahepatic Bile Duct	
Liver	C22.0, C22.2-C22.4, C22.7, C22.9
Intrahepatic Bile Duct	C22.1
Gallbladder	C23
Other Biliary	C24
Pancreas	C25
Retroperitoneum	C48.0
Peritoneum, Omentum and Mesentery	C45.1+, C48.1-C48.2
Other Digestive Organs	C26.8-C26.9, C48.8
Respiratory System	
Nose, Nasal Cavity and Middle Ear	C30-C31
Larynx	C32
Lung and Bronchus	C34
Pleura	C38.4, C45.0+
Trachea, Mediastinum and Other Respiratory Organs	C33, C38.1-C38.3, C38.8, C39
Bones and Joints	C40-C41
Soft Tissue including Heart	C47, C49, C38.0, C45.2+
Skin excluding Basal and Squamous	
Melanoma of the Skin	C43
Other Non-Epithelial Skin	C44, C46+
Breast	C50
Female Genital System	
Cervix Uteri	C53
Corpus and Uterus, NOS	
Corpus Uteri	C54
Uterus, NOS	C55
Ovary	C56
Vagina	C52
Vulva	C51
Other Female Genital Organs	C57-C58

Appendix E: SEER Cancer Cause of Death Analysis Categories (Continued)

Cancer Causes of Death	ICD-10
Male Genital System	
Prostate	C61
Testis	C62
Penis	C60
Other Male Genital Organs	C63
Urinary System	
Bladder	C67
Kidney and Renal Pelvis	C64-C65
Ureter	C66
Other Urinary Organs	C68
Eye and Orbit	C69
Brain and Other Nervous System	C70, C71, C72
Endocrine System	
Thyroid	C73
Other Endocrine Including Thymus	C37, C74-C75
Lymphoma	
Hodgkin's Lymphoma	C81
Non-Hodgkin's Lymphoma	C82-C85, C96.3
Myeloma	C90.0, C90.2
Leukemia	
Lymphocytic Leukemia	
Acute Lymphocytic Leukemia	C91.0
Chronic Lymphocytic Leukemia	C91.1
Other Lymphocytic Leukemia	C91.2-C91.4, C91.7, C91.9
Myeloid and Monocytic Leukemia	
Acute Myeloid	C92.0, C92.4-C92.5, C94.0, C94.2
Acute Monocytic Leukemia	C93.0
Chronic Myeloid Leukemia	C92.1
Other Myeloid/Monocytic Leukemia	C92.2-C92.3, C92.7, C92.9, C93.1-C93.2, C93.7, C93.9
Other Acute Leukemia	C94.4, C94.5, C95.0
Aleukemic, Subleukemic and NOS	C90.1, C91.5, C94.1, C94.3, C94.7, C95.1, C95.2, C95.7, C95.9
Mesothelioma (ICD-10 only)+	C45+
Kaposi Sarcoma (ICD-10 only)+	C46+
Miscellaneous Malignant Cancer	C26.1, C45.7+, C45.9+, C76-C80, C88, C96.0-C96.2, C96.7, C96.9, C97

Source: <http://seer.cancer.gov/codrecode>

REFERENCES

- Bast et Al. Cancer Edition 5. American Cancer Society, 2000.
- Beahrs, O.H. et al, AJCC Staging Manual, 5th Edition. American Joint Committee on Cancer, Lippincott, Williams and Wilkins. 1995.
- Cancer Facts and Figures, 2014. Atlanta. American Cancer Society. 2014.
- Cancer Progress Report, US Department of Health and Human Services, National Cancer Institute. 2005.
- Devita, Vincent et al. CANCER Principles and Practice of Oncology. Lippincott. Williams & Wilkins. 2001.
- Harvard Center for Cancer Prevention. Harvard Report on Cancer Prevention Volume 1: Causes of Human Cancer, 1996:7 (S1): 7-15.
- Howlader N, Noone AM, Krapcho M, Garshell J, Miller D, Altekruse SF, Kosary CL, Yu M, Ruhl J, Tatalovich Z, Mariotto A, Lewis DR, Chen HS, Feuer EJ, Cronin KA (eds). SEER Cancer Statistics Review, 1975-2011, National Cancer Institute. Bethesda, MD, http://seer.cancer.gov/csr/1975_2011/, based on November 2013 SEER data submission, posted to the SEER web site, April 2014.
- Jemal A, Simard EP, Dorell C, Noone AM, Markowitz LE, Kohler B, Ehemann C, Saraiya M, Bandi P, Saslow D, Cronin KA, Watson M, Schiffman M, Henley SJ, Schymura MJ, Anderson RN, Yankey D, and Edwards BK. Annual Report to the Nation on the Status of Cancer, 1975-2009, Featuring the Burden and Trends in HPV-Associated Cancers and HPV Vaccination Coverage Levels. *J Natl Ca Inst* 2013 Feb;105(3). Epub 2013 Jan 7.
- Lenhard, Raymond et al. Clinical Oncology. The American Cancer Society. Bethesda, MD. 2001.
- Nascar, Philip and Harris Pastides. Fundamentals of Cancer Epidemiology. Aspen Publications, Maryland. 2001.
- Schottenfeld, David and Joseph Fraumenis, Cancer Epidemiology and Prevention. 2nd Edition. Oxford University Press. 1996.
- Surveillance, Epidemiology, and End Results (SEER) Program of the National Cancer Institute, Cancer Query Systems, <http://seer.cancer.gov/canques/>

This document is available online at
<http://getscreened.sd.gov/documents/Cancer2014.pdf>

A limited number were printed and publication, funded by the South Dakota Cancer Registry with a grant from the Centers for Disease Control and Prevention – grant number U58/DP003943. For more information or additional copies (if available), contact the South Dakota Department of Health at 605.773.3361

Per SDCL 5-18D-15: 75 copies of this publication have been printed on recycled paper by the South Dakota Department of Health at a cost of **\$7.38** each.