

# *South Dakota*

## *An Annual Report on Cancer Incidence and Mortality Among South Dakotans, 2018*



South Dakota Cancer Registry  
April 2021

# Cancer in South Dakota 2018



---

South Dakota Cancer Registry, South Dakota Department of Health



## Preface

“Cancer in South Dakota 2018” is the 26th annual report from the South Dakota Cancer Registry (SDCR) in the Office of Chronic Disease Prevention and Health Promotion in the Division of Family and Community Health within the South Dakota Department of Health (DOH). The report contains 2018 cancer incidence and mortality data of South Dakota residents.

## Acknowledgements

The South Dakota Cancer Registry acknowledges all the Certified Tumor Registrars in hospital cancer registries who work diligently to maintain quality data, hospitals, clinics, physicians, pathology laboratories and other providers and health entities that submit data to the SDCR, and co-workers in the Department of Health.

## Funding Source

South Dakota Cancer Registry funded this publication with a grant from the Centers for Disease Control and Prevention – grant number DP006293.

## Changes in Report

Age-adjusted rates were calculated using the 2000 US standard million and the US Census Bureau 2002 – 2018 population estimates for South Dakota. Reports published before 2012 used the 2000 South Dakota census population for age-adjusted calculations and are not comparable to this report.

## Requests for Information

Data queries, cancer concerns, information, or requests for additional copies (if available) should be addressed to:

Kay Dosch, MA, CTR, SDCR Coordinator  
South Dakota Cancer Registry  
615 East 4th Street  
Pierre, SD 57501-1700  
Phone: (605) 773-6345

## Suggested Citation

*Cancer in South Dakota, 2018*. Department of Health, Pierre, SD. April 2021. All material in this report may be reproduced or copied without permission. However, citation as to source is appreciated.

**Web Access:** This report is available on the internet at the following URL:

<http://getscreened.sd.gov/documents/Cancer2018.pdf>

The following Department of Health personnel were the main contributors to the development of this report:

Mary Sarvis	SDCR Data Manager (Retired)
Morgan Vedvei	SDCR Data Manager
Patricia Da Rosa	DDS, MSc/MPH Data Analyst
Kay Dosch	SDCR Coordinator
Mark Gildemaster	Management Analyst
Rebecca Piroutek	Communications Coordinator



## TABLE OF CONTENTS

PREFACE .....	i
TABLE OF CONTENTS .....	iii
LIST OF TABLES .....	iv
LIST OF FIGURES .....	v
I. EXECUTIVE SUMMARY .....	1
II. INTRODUCTION .....	3
III. TECHNICAL NOTES .....	5
IV. CANCER INCIDENCE .....	8
V. CANCER CASES AND DEATHS BY RANK .....	13
VI. STAGE AT DIAGNOSIS .....	16
VII. CANCER MORTALITY .....	18
VIII. YEARS OF POTENTIAL LIFE LOST .....	23
IX. AVERAGE YEARS OF LIFE LOST .....	25
X. SELECTED CANCER SITES INCIDENCE AND MORTALITY .....	27
Bladder .....	28
Breast (Female) .....	30
Cervix Uteri .....	32
Colorectal .....	34
Corpus and Uterus .....	36
Kidney and Renal Pelvis .....	38
Leukemia .....	40
Lung and Bronchus .....	42
Melanoma (Skin) .....	44
Myeloma .....	46
Non-Hodgkin's Lymphoma .....	48
Ovary .....	50
Pancreas .....	52
Prostate .....	54
Stomach .....	56
Thyroid.....	58
XI. APPENDICES	
Appendix A: 2000 US Standard Million Population.....	60
Appendix B: 2009-2018 South Dakota Estimated Population .....	60
Appendix C: Race in South Dakota by County, 2018 Estimated Population .....	61
Appendix D: SEER Incidence Site Analysis Categories .....	62-64
Appendix E: SEER Cancer Cause of Death Analysis Categories .....	65-66
REFERENCES .....	67

## LIST OF TABLES

Table 1: Cancer Cases and Incidence Rates by County, South Dakota, 2018 and 2014-2018 Average.....	8
Table 2: Age-adjusted Incidence Rates by County for Selected Sites, 2018.....	9
Table 3: Age-adjusted Incidence Rates by Site, Gender, and Race, South Dakota, 2018	10
Table 4: Percentage of Selected Cancers by Age Groups in South Dakota, 2018.....	12
Table 5: South Dakota Stage at Diagnosis, All Cases, 2018.....	16
Table 6: Stage at Diagnosis for Selected Sites by Race, South Dakota, 2018.....	17
Table 7: Cancer Mortality and Rates by County, South Dakota, 2018 and 2014-2018 .....	18
Table 8: Age-adjusted Mortality Rates by County for Selected Sites, South Dakota, 2018	19
Table 9: Age-adjusted Mortality Rates by Site, Gender, and Race, South Dakota, 2018..	20
Table 10: Percentage of Cancer Deaths by Age Groups and Selected Primary Sites, South Dakota, 2018 .....	22
Table 11: Leading Causes of Years of Potential Life Lost (to age 75), South Dakota, 2018	23
Table 12: Bladder Incidence and Mortality Summary, 2018.....	28
Table 13: Breast (Female) Incidence and Mortality Summary, 2018.....	30
Table 14: Cervix Uteri Incidence and Mortality Summary, 2018.....	32
Table 15: Colorectal Incidence and Mortality Summary, 2018 .....	34
Table 16: Corpus and Uterus, NOS Incidence and Mortality Summary, 2018.....	36
Table 17: Kidney and Renal Pelvis Incidence and Mortality Summary, 2018.....	38
Table 18: Leukemia Incidence and Mortality Summary, 2018 .....	40
Table 19: Lung and Bronchus Incidence and Mortality Summary, 2018 .....	42
Table 20: Melanoma Incidence and Mortality Summary, 2018 .....	44
Table 21: Myeloma Incidence and Mortality Summary, 2018 .....	46
Table 22: Non-Hodgkin's Lymphoma Incidence and Mortality Summary, 2018 .....	48
Table 23: Ovarian Incidence and Mortality Summary, 2018 .....	50
Table 24: Pancreas Incidence and Mortality Summary, 2018 .....	52
Table 25: Prostate Incidence and Mortality Summary, 2018.....	54
Table 26: Stomach Incidence and Mortality Summary, 2018.....	56
Table 27: Thyroid Incidence and Mortality Summary, 2018 .....	58

## LIST OF FIGURES

Figure 1:	All Sites Cancer Incidence Rates by Race and Gender, South Dakota, 2018..	12
Figure 2:	Percent Distribution of Cancer Cases and Deaths by Rank and Gender, South Dakota, 2018 .....	13
Figure 3:	Cancer Incidence Rates by County, South Dakota, 2018 (Map) .....	14
Figure 4:	Cancer Mortality Rates by County, South Dakota, 2018 (Map) .....	15
Figure 5:	Stage at Diagnosis, All Sites, South Dakota, 2018.....	16
Figure 6:	Stage at Diagnosis in South Dakota, White, 2018.....	17
Figure 7:	Stage at Diagnosis in South Dakota, American Indian, 2018 .....	17
Figure 8:	All Sites Cancer Mortality Rates by Race and Gender, South Dakota, 2018....	22
Figure 9:	Years of Potential Life Lost (YPLL) Before Age 75 for Leading Causes of Death, by Race, South Dakota, 2018.....	23
Figure 10:	Years of Potential Life Lost (YPLL) Before Age 75 for Leading Causes of Death, by Race, South Dakota, 2014-2018 .....	23
Figure 11:	YPLL for Selected Cancers Among Whites, South Dakota, 2018 .....	24
Figure 12:	YPLL for Selected Cancers Among American Indians, South Dakota, 2018....	24
Figure 13:	AYLL for Selected Cancers Among Whites, South Dakota, 2018 .....	25
Figure 14:	AYLL for Selected Cancers Among American Indians, South Dakota, 2018....	25
Figure 15:	Bladder Cancer Stage at Diagnosis, South Dakota, 2018.....	28
Figure 16:	Bladder Cancer Number of Cases and Deaths by Age, South Dakota, 2018...	29
Figure 17:	Bladder Cancer Cases and Deaths by Year, South Dakota, 2001-2018 .....	29
Figure 18:	Bladder Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018.....	29
Figure 19:	Female Breast Cancer Stage at Diagnosis, South Dakota, 2018 .....	30
Figure 20:	Female Breast Cancer Number of Cases and Deaths by Age, South Dakota, 2018 .....	31
Figure 21:	Female Breast Cancer Cases and Deaths by Year, South Dakota, 2001-2018	31
Figure 22:	Female Breast Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018.....	31
Figure 23:	Cervix Uteri Cancer Stage at Diagnosis, South Dakota, 2018.....	32
Figure 24:	Cervix Uteri Cancer Number of Cases and Deaths by Age, South Dakota, 2018 .....	33
Figure 25:	Cervix Uteri Cancer Cases and Deaths by Year, South Dakota, 2001-2018....	33
Figure 26:	Cervix Uteri Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018.....	33
Figure 27:	Colorectal Cancer Stage at Diagnosis, South Dakota, 2018 .....	34
Figure 28:	Colorectal Cancer Number of Cases and Deaths by Age, South Dakota, 2018	35
Figure 29:	Colorectal Cancer Cases and Deaths by Year, South Dakota, 2001-2018 .....	35
Figure 30:	Colorectal Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018.....	35
Figure 31:	Corpus and Uterus, NOS Cancer Stage at Diagnosis, South Dakota, 2018....	36
Figure 32:	Corpus and Uterus, NOS Cancer Number of Cases and Deaths by Age, South Dakota, 2018 .....	37
Figure 33:	Corpus and Uterus, NOS Cancer Cases and Deaths by Year, South Dakota, 2001-2018 .....	37
Figure 34:	Corpus and Uterus, NOS Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018 .....	37
Figure 35:	Kidney and Renal Pelvis Cancer Stage at Diagnosis, South Dakota, 2018 .....	38
Figure 36:	Kidney and Renal Pelvis Cancer Number of Cases and Deaths by Age, South Dakota, 2018.....	39

## LIST OF FIGURES (cont'd)

Figure 37: Kidney and Renal Pelvis Cancer Cases and Deaths by Year, South Dakota, 2001-2018 .....	39
Figure 38: Kidney and Renal Pelvis Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018 .....	39
Figure 39: Leukemia Number of Cases and Deaths by Age, South Dakota, 2018 .....	41
Figure 40: Leukemia Cases and Deaths by Year, South Dakota, 2001-2018 .....	41
Figure 41: Leukemia Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018 .....	41
Figure 42: Lung and Bronchus Cancer Stage at Diagnosis, South Dakota, 2018 .....	42
Figure 43: Lung and Bronchus Cancer Number of Cases and Deaths by Age, South Dakota, 2018 .....	43
Figure 44: Lung and Bronchus Cancer Cases and Deaths by Year, South Dakota, 2001-2018 .....	43
Figure 45: Lung and Bronchus Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018.....	43
Figure 46: Melanoma of the Skin Stage at Diagnosis, South Dakota, 2018 .....	44
Figure 47: Melanoma of the Skin Number of Cases and Deaths by Age, South Dakota, 2018 .....	45
Figure 48: Melanoma of the Skin Cases and Deaths by Year, South Dakota, 2001-2018 .....	45
Figure 49: Melanoma of the Skin Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018.....	45
Figure 50: Myeloma Number of Cases and Deaths by Age, South Dakota, 2018 .....	47
Figure 51: Myeloma Cases and Deaths by Year, South Dakota, 2001-2018 .....	47
Figure 52: Myeloma Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018 .....	47
Figure 53: Non-Hodgkin's Lymphoma Stage at Diagnosis, South Dakota, 2018.....	48
Figure 54: Non-Hodgkin's Lymphoma Number of Cases and Deaths by Age, South Dakota, 2018.....	49
Figure 55: Non-Hodgkin's Lymphoma Cases and Deaths by Year, South Dakota, 2001-2018 .....	49
Figure 56: Non-Hodgkin's Lymphoma Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018.....	49
Figure 57: Ovarian Cancer Stage at Diagnosis, South Dakota, 2018.....	50
Figure 58: Ovarian Cancer Number of Cases and Deaths by Age, South Dakota, 2018 ..	51
Figure 59: Ovarian Cancer Cases and Deaths by Year, South Dakota, 2001-2018.....	51
Figure 60: Ovarian Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018.....	51
Figure 61: Pancreatic Cancer Stage at Diagnosis, South Dakota, 2018 .....	52
Figure 62: Pancreatic Cancer Number of Cases and Deaths by Age, South Dakota, 2018 .....	53
Figure 63: Pancreatic Cancer Cases and Deaths by Year, South Dakota, 2001-2018.....	53
Figure 64: Pancreatic Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018.....	53
Figure 65: Prostate Cancer Stage at Diagnosis, South Dakota, 2018.....	54
Figure 66: Prostate Cancer Number of Cases and Deaths by Age, South Dakota, 2018..	55
Figure 67: Prostate Cancer Cases and Deaths by Year, South Dakota, 2001-2018.....	55

## LIST OF FIGURES (cont'd)

Figure 68: Prostate Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018.....	55
Figure 69: Stomach Cancer Stage at Diagnosis, South Dakota, 2018 .....	56
Figure 70: Stomach Cancer Number of Cases and Deaths by Age, South Dakota, 2018. ....	57
Figure 71: Stomach Cancer Cases and Deaths by Year, South Dakota, 2001-2018 .....	57
Figure 72: Stomach Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018.....	57
Figure 73: Thyroid Cancer Stage at Diagnosis, South Dakota, 2018 .....	58
Figure 74: Thyroid Cancer Number of Cases and Deaths by Age, South Dakota, 2018 ...	59
Figure 75: Thyroid Cancer Cases and Deaths by Year, South Dakota, 2001-2018 .....	59
Figure 76: Thyroid Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001-2018.....	59



## I. EXECUTIVE SUMMARY

---

This report summarizes the burden of cancer in South Dakota and includes cancer incidence and mortality data. The data will enable the many organizations working with cancer prevention and control to identify public health problems, target goals for cancer control, and to inform citizens and health care professionals about risks, early detection, and treatment.

### Incidence 2018

- South Dakota had 4,820 reportable invasive cases of cancer diagnosed, which excludes the less life-threatening cancers such as *in situ* cancers (except *in situ* bladder cases) and the common skin cancers.
- Each day 13 cases of cancer were diagnosed in residents of South Dakota; this includes only cases of invasive cancer and *in situ* bladder.
- The five most diagnosed cancer sites (prostate, female breast, lung, colorectal, melanoma) accounted for 55% of all cancer cases.
- Prostate cancer was the most common reportable malignancy with 690 cases, 14.3% of all cases and 27.0% of cases for men.
- Female breast cancer was the second most common reportable cancer with 654 cases, 13.6% of all cases and 28.9% of cases for women.
- Lung cancer was the third most common reportable malignancy with 597 cases, accounting for 12.4% of all cases.
- Colon and rectal cancers were the fourth most common malignancy with 409 cases, 8.5% of all cases.
- Melanoma cancers were the fifth most common malignancy with 304 cases, 6.3% of all reported cases.
- Fifty-three percent of all new cancers were diagnosed in males and 47% were in females.
- Males had an age-adjusted incidence rate of 482.9 per 100,000, which was higher than females who had an age-adjusted rate of 422.1 per 100,000.
- Whites accounted for 93.7% of cancer cases with 4,517 cases whereas American Indians were 5.1% with 248 cases.
- The American Indian age-adjusted incidence rate was 499.8, which is higher than the age-adjusted rate among whites of 451.7.
- The South Dakota age-adjusted incidence rate for 2018 was 449.6, not a significantly different from the US SEER 2018 age-adjusted incidence rate of 450.5 per 100,000 persons.

## Mortality 2018

- Overall, cancer was the second leading cause of death in South Dakota.
- In 2018, 1,661 South Dakotans died from cancer, accounting for one in every four deaths in South Dakota.
- Each day over four South Dakotans died from cancer.
- The five cancer sites (lung, colorectal, pancreas, female breast, prostate) caused over half of all cancer deaths.
- Lung and bronchus cancers were the leading cause of cancer deaths at 395 deaths or 23.8% of all cancer deaths. Lung and bronchus cancers caused one in four deaths due to cancer.
- Colorectal cancer was the second leading cause of cancer deaths with 168 deaths, 10.1% of all cancer deaths.
- Pancreatic cancer was the third leading cause of death with 116 deaths, 7.0% of all cancer deaths.
- Female breast cancer was the fourth leading cause of cancer deaths with 112 deaths, 6.7% of all cancer deaths and 14.9% of all female cancer deaths.
- Prostate cancer was the fifth leading cause of death with 92 deaths, 5.5% of all cancer deaths and 10.1% of all male cancer deaths.
- Over half, 55% of all cancer deaths were males and 45% were females.
- Males had an age-adjusted death rate of 179.2 per 100,000 males, 42.1% higher than females with an age-adjusted rate of 126.1 deaths per 100,000 females.
- Whites accounted for 91.8% of deaths with 1,525 deaths, whereas American Indians were 6.9% with 115 deaths.
- The American Indian age-adjusted death rate was 257.8 which is 79.0% higher than the rate among whites at the age-adjusted death rate of 144.0.
- South Dakota's age-adjusted death rate for 2018 was 147.9, with no significant difference from the US SEER 2018 rate of 149.0.

## Trends

- Melanoma incidence cases have increased significantly since 2009.
- For the last decade, female breast cancer mortality rates have remained steady.

## II. INTRODUCTION

---

A limited cancer data collection system was established in 1992 under South Dakota Codified Laws, SDCL 1-43-1 to 1-43-18, and Administrative Rules ARSD 44:22:01 to 44:22:05, inclusive. The South Dakota Cancer Registry (SDCR) was established in 2001 to develop a statewide, population-based cancer surveillance system. However, the state legislature amended the law to expand reporting to reflect statewide surveillance. SDCL 1-43-14 has been in effect since July 1, 2005:

*Any hospital licensed pursuant to chapter 34-12, physician licensed pursuant to chapter 36-4, physician assistant licensed pursuant to chapter 36-4A, nurse practitioner or nurse midwife licensed pursuant to chapter 36-9A, pathology laboratory, or free-standing radiology center that detects, diagnoses, or treats a cancer case in South Dakota shall submit a report to the Department of Health as required by § 1-43-11 to 1-43-17, inclusive.*

Reportable cancers for 2018 include benign brain, benign central nervous system and all malignant neoplasms except basal and squamous cell carcinomas of the skin and *in situ* cervical cancers. In addition to receiving cases from hospital cancer registries the SDCR actively follows back pathology reports and abstracts cases from facilities without tumor registries.

The SDCR performs many quality assurance procedures to ensure that the data are valid. The data are run through numerous edits and consolidated if received from more than one reporting source. In addition, the SDCR links the incidence data with mortality files to identify persons whose death records show cancer as a cause of death, but these cancers were not reported to the central registry. The SDCR also links the incidence file with the Indian Health Service database to identify any American Indian South Dakotan who was misclassified as another race.

The SDCR uses the reported cancer incidence data, the mortality data and health behavior surveys collected by the department's Health Statistics Office and Vital Records to provide useful information for cancer control and prevention programs, researchers, clinicians, and policy makers. The SDCR can answer several epidemiological questions such as:

- How many South Dakotans are diagnosed or die from cancer each year?
- What are the most common cancers?
- When are cancers being diagnosed, i.e. at what stage?
- Which cancers are the deadliest?
- Who is most affected by cancer?
- What are the trends in cancer incidence and mortality?
- Where are cancers occurring?
- Where and what are the disparities?
- Are screening efforts working?

Every life is touched by cancer somehow, whether one is diagnosed with the disease or has a family member or friend with the disease. Although cancer is primarily a disease of people over 50 years old, the younger a person dies from cancer, the more significant the impact on societal and economic costs. Cancer concerns voiced by South Dakotans are a priority for the SDCR.

As the SDCR continues to collect population-based data and as more health care entities and providers report cases, more questions can be answered with the data. For example, SDCR can look at modifiable risk factors such as obesity and exercise. The Harvard Report on Cancer Prevention in 1996 researched the risk factors for cancer. The estimated percentages (which are still relevant today) of total cancer deaths attributed to established causes of cancer were:

Risk Factor	Percentage
Tobacco	30%
Adult diet/obesity	30%
Sedentary lifestyle	5%
Occupational factors	5%
Family history of cancer	5%
Viruses/biological agents	5%
Perinatal factors/growth	5%
Reproductive factors	3%
Alcohol	3%
Socioeconomic status	3%
Environmental pollution	2%
Ionizing /UV radiation	2%
Prescription drugs/medical	1%
Salt/food additives/contaminants	1%

The most successful way to prevent cancer is to limit the number of modifiable risk factors by following these guidelines:

- Do not smoke
- Maintain a healthy weight
- Get at least 30 minutes of physical activity every day
- Eat a healthy diet
- Limit alcohol intake to less than one drink a day
- Protect yourself from the sun

Some risk factors for cancer can be avoided, but many cannot. For example, both smoking and inheriting certain genes are risk factors for some types of cancer, but only smoking can be avoided. Risk factors that a person can control are called modifiable risk factors.

Source: National Cancer Institute

The SDCR continuously expands data collection. This will help prevention and control programs to target at-risk populations as well as support epidemiologic studies. The end goal is to reduce the cancer burden by producing valid and accurate data reflecting the complete assessment of cancer in South Dakota, and to disseminate the information in a timely manner.

### III. TECHNICAL NOTES

---

**Age-adjusted incidence rate:** Age-adjusted incidence rates are calculated using the direct method and standardized to the age distribution of the 2000 US standard population (Appendix A). Age adjustment allows rates for one geographic area to be compared with rates from other geographic areas that may have differences in age distributions. Any observed differences in age-adjusted incidence rates between populations are not due to different age structures. Reports prior to 1999 used the 1970 US standard population. In conformity with the National Cancer Institute's (NCI) Surveillance, Epidemiology, and End Results (SEER) Program guidelines, the incidence rates for cancer sites exclude the following:

- *In situ* cases, except bladder
- Basal and squamous cell skin cancers
- Cases with unknown age
- Cases with unknown gender

**Age-adjusted mortality rates:** Mortality rates are calculated for total cases and separately for males and females. The mortality rates are age-adjusted to the 2000 US standard population using five-year groups and are per 100,000 persons. Rates are presented for 2018 and for the five-year period, 2014-2018.

**Age-specific incidence rates:** Age-specific rates are calculated by dividing the number of cases for a given age group by the total population of that age group and are expressed as an average annual rate per 100,000 persons by age group. Age-specific rates exclude the same types of cases that are excluded from age-adjusted incidence rates. These rates, however, are crude rates, i.e. not age-adjusted.

**Annual percent change (APC):** The annual percent change is the average rate of change in a cancer rate per year in a given time frame indicating how fast or how slowly a cancer rate has increased or decreased each year over a period of years. A negative APC describes a decreasing trend, and a positive APC describes an increasing trend.

**Average years of life lost (AYLL):** This is the extent to which life is cut short due to premature death. This is obtained by dividing the years of potential life lost (YPLL) by the number of deaths. On average each person who dies from cancer loses 15 years of their life.

**Cancer case definitions:** A "cancer case" is defined as the primary cancer site, i.e., the site where the cancer started. Since an individual can have more than one primary cancer site, the number of incident cancer cases could be greater than the number of persons who are diagnosed with cancer. A *metastasis* is not a primary site.

**Changes in diagnostic criteria:** Early detection resulting from either screening or early response to symptoms may result in increasing diagnosis of small tumors that are not yet life-threatening. This may raise incidence and survival rates but without changes in mortality rates. Cancers likely to be affected are breast, colon, cervix uteri, prostate, and melanoma. Prostate cancer is particularly prone to changing diagnostic criteria.

**Confidence intervals (CI):** A confidence interval tells how confident we are of the accuracy of the calculated rates. The SDCR uses a computed interval with a given probability of 95%, i.e., the true value of the calculated rate is contained within the interval. Thus, given a calculated rate of 191.4 and a confidence interval of 182.1 to 200.8, it is better to say that the actual rate will fall between 182.1 and 200.8. The larger the sample size, the shorter the interval size, giving us more certainty that the rate is correct. When CI for percentages contains zero, the rate is considered to be stable. Above zero, the statistical significance is higher and below zero it is lower.

**Data source:** All data, tables, and figures come from the South Dakota Department of Health, *American Cancer Society Facts and Figures 2018* or *SEER Cancer Statistics Review 1975-2017* and should be cited as such if taken out of this report in part. SEER data represents approximately 10% of the U.S population.

**Disparity:** Health disparities are differences in the incidence, prevalence, mortality, and burden of diseases and other adverse health conditions that exist among specific population groups in the United States.<sup>1</sup> Health disparities can be defined as a specific group bearing a disproportionate share of negative health outcomes compared to the general population, i.e., disease, disability, and death.<sup>2</sup> Disparity can occur due to poverty, living in geographically underserved areas and belonging to specific minority groups.

**Early detection/screening:** Improved early detection/screening may increase incidence and survival rates. Increases may occur as a result of the introduction of new procedures. The interval between the time a cancer is diagnosed by a screening procedure and the time when it would have been diagnosed in the absence of screening procedures is called the lead-time. Changes in lead-time, for example, in breast cancer diagnosis, have led to increased survival rates and reduction of mortality.

**Limitations to data interpretation and comparison:** Several factors must be considered when reviewing cancer statistics and interpreting them. A cancer registry database is a fluid and dynamic database, therefore, the reported number of new cases in a particular race, gender, and age cancer category may change for the calendar year for which the data have already been reported in a previous publication. Additional cancer cases previously overlooked for a given diagnosis year may be found and reported to the central registry. There may also be elimination of duplicate records for the same patient, often due to name changes or spelling corrections.

**Metastasis:** When cancer spreads from the primary site to other organs or tissues of the body, it is said to metastasize. Cancer usually spreads through the blood or the lymphatic system.

**Mortality/incidence ratio (M/I):** This ratio is calculated by dividing the number of deaths in a given year by the number of new cancers diagnosed in the same year. The death to case ratio provides a crude indication of the prognosis for patients. A ratio approaching 1.0, when the number of deaths equals the number of cases for a particular type of cancer, indicates a poor prognosis. A lower ratio indicates fewer deaths relative to the number of cases and suggests a better prognosis.

**Percent change:** This is the difference between two rates expressed as a percentage.

**Racial misclassifications:** When race is not specified in a source document and the default is to record these cases as white or unknown, the results are considered biased. Numerator error can occur because of misclassification.

**Rate comparisons:** Rate comparisons are difficult to interpret when comparing age-adjusted and age-specific rates based on fewer than 10 cases. In comparing rates among geographic areas such as counties, states and health districts, the absolute numbers and differences in demographics should be considered, as well as clinical significance of the disease. Data quality indicators for each registry should also be reviewed. Interpretations made without considering these factors may be misleading. There will also be differences between mortality statistics published by various agencies and the mortality rates in this report.

**Risks and associated risk factors:** These were developed using the “American Cancer Society Textbook of Oncology,” and the Harvard Cancer Center, Causes of Human Cancer.

**Stage at time of diagnosis:** Staging is the process of describing the extent or spread of disease from the origin, which is the primary site. Summary staging is the standard used for comparison nationally. SEER Summary Stages 2000 are defined as follows:

- **In Situ:** Malignant cells are within the cell group from which they arose, without penetration of the basement membrane of the tissue and no stromal invasion. *In situ* is “in place”.
- **Localized:** The malignant cells are limited to the organ of origin and have spread no farther than the organ in which they started.
- **Regional:** The tumor is beyond the limits of the organ of origin by direct extension to adjacent areas with or without lymph node involvement.
- **Distant:** The primary tumor has broken away and has traveled, growing secondary tumors in other parts of the body. It has metastasized.

*In situ* and localized stages are the **early stages** of diagnosis. Regional and distant stages are **late stage** diagnoses.

**Staging:** Stage is based on an assessment of the size of the primary tumor, whether it has spread, and, if so, how far. Because an accurate diagnosis is essential to effective treatment, physicians might use physical exams, imaging, lab tests, a biopsy, an analysis of the patient’s body fluids, and surgery in various combinations in the staging process. Advancement in diagnostic procedures may change in due time. These advancements might increase the chance that a given cancer will be diagnosed at a more advanced stage. For example, with new scanning methods metastases can be detected. Therefore, if someone was previously diagnosed with a localized tumor, they may now be staged as distant. This is called stage migration and can affect the analysis of all solid tumors.

**Statistical significance:** This determines whether an event happens by chance alone. The null hypothesis states that in a given place and a period of time, all events occur randomly by chance. If not, then there is statistical significance. Confidence intervals are used to test statistical significance in this report. If the confidence intervals of two different rates intersect each other, then there is no statistical difference between the two rates.<sup>3</sup> However, if the confidence intervals do not intersect one another, there is statistical significance. This report looks at the South Dakota rates as compared to the US national rates using SEER data.

In South Dakota, case counts can be very low; therefore, magnitude bias is inherent with confidence intervals and z- tests. For example, in the year 2001, cervical cancer rates were 10 per 100,000 American Indian women, a cervical cancer age-adjusted rate six times higher than white women in South Dakota. However, the case counts were two for American Indians and 10 whites. Small numbers result in wider confidence intervals, thus less confidence in the data.

**Years of potential life lost (YPLL):** The years of potential life lost are calculated for each individual who dies of cancer by determining the number of years of additional expected life if that person had lived to 75 years. The YPLL in the general population associated with a particular cancer is the sum of this expectation over all those individuals who died of that cancer in a particular year. YPLL reflects the burden of cancer on younger persons while mortality rates reflect the burden on older persons.

---

<sup>1</sup><http://epi.grants.cancer.gov/ResPort/HDoverview.html>

<sup>2</sup><http://www.omni.org/docs/CMHFProceedings.pdf>

<sup>3</sup>BIostatistics *The Bare Essentials*, 2nd edition Norman and Shreiner Page 512

#### IV. CANCER INCIDENCE

South Dakota collected 4,820 new reportable cancer cases in 2018. Data at the county level ranged from a low incidence rate of 39.2 in Corson County to a high of 771.4 in Aurora County. There were eight counties with rates significantly lower than the state incidence rate of 446.7. Four counties had a significantly higher rate, compared to one county in 2017.

The United States incidence rate in 2018 was 450.5 and the South Dakota incidence rate was 446.7 per 100,000 persons.

A map displaying the 2018 incidence rates by county can be located on page 14 of this report. On page 15 is a map displaying the mortality rates by county.

**Table 1 : Cancer Cases and Incidence Rates by County  
South Dakota, 2018 and 2013-2018 Average**

County	2018		2013-2018 <sup>^</sup>	
	Cases	Rate	Cases	Rate
<b>South Dakota</b>	<b>4,820</b>	<b>446.7</b>	<b>4,721</b>	<b>455.4</b>
Aurora	30	771.1 ▲	21	554.1
Beadle	95	385.8	110	481.4
Bennett	12	289.1	13	383.0
Bon Homme	44	476.4	44	441.7
Brookings	137	422.4	147	482.9
Brown	243	508.8	224	471.4
Brule	27	330.4	37	527.5
Buffalo	10	759.2	7	493.3
Butte	53	359.6	61	435.4
Campbell	2	64.1 ▼	8	365.5
Charles Mix	50	431.3	58	484.4
Clark	31	580.3	28	513.5
Clay	68	547.2	64	525.9 ▲
Codington	174	476.0	167	467.0
Corson	2	39.2 ▼	11	287.4 ▼
Custer	62	357.9	57	372.1 ▼
Davison	125	504.9	123	489.9
Day	43	503.6	43	471.5
Deuel	18	244.6 ▼	29	445.0
Dewey	24	481.5	22	444.9
Douglas	20	481.0	22	543.3
Edmunds	28	523.3	30	526.6
Fall River	57	447.5	59	478.9
Faulk	17	451.7	17	475.8
Grant	43	387.8	50	462.8
Gregory	26	441.7	28	432.3
Haakon	15	522.2	15	455.2
Hamlin	35	502.0	33	441.9
Hand	19	284.2 ▼	24	469.4
Hanson	19	567.2	19	568.5
Harding	4	207.4 ▼	4	260.3 ▼
Hughes	116	506.1	106	489.4
Hutchinson	64	593.4	51	468.4
Hyde	12	388.2	11	502.0
Jackson	13	403.8	14	424.3
Jerauld	12	437.9	15	428.3
Jones	8	486.9	8	538.8
Kingsbury	50	576.2	47	609.1 ▲
Lake	73	385.7	76	425.2
Lawrence	150	391.0	140	404.9 ▼
Lincoln	254	416.8	237	439.7
Lyman	19	459.9	20	434.4
McCook	52	638.4 ▲	40	550.3 ▲
McPherson	16	445.7	20	425.6
Marshall	33	414.2	24	374.4 ▼
Meade	117	358.7 ▼	122	395.9 ▼
Mellette	12	512.1	10	425.2
Miner	13	493.7	17	508.7
Minnehaha	1,048	520.7 ▲	986	510.3 ▲
Moody	38	408.3	34	385.4 ▼
Oglala Lakota	35	355.2	41	422.7
Pennington	598	416.0	576	425.1 ▼
Perkins	10	223.7 ▼	14	317.5 ▼
Potter	18	517.2	18	456.5
Roberts	47	314.9 ▼	51	390.0 ▼
Sanborn	15	456.8	15	486.9
Spink	43	470.7	48	530.1 ▲
Stanley	16	451.6	18	445.0
Sully	8	303.2	5	261.6 ▼
Todd	29	456.3	29	448.3
Tripp	36	390.0	39	448.1
Turner	54	424.9	57	474.2
Union	119	576.5 ▲	100	532.3 ▲
Walworth	26	316.9	32	374.3 ▼
Yankton	122	408.0	119	395.0 ▼
Ziebach	11	431.7	5	199.7 ▼

<sup>^</sup> Counts less than three are suppressed. Incidence rates with counts less than 20 are generally considered unstable. ▲Rate significantly higher than the state rate. ▼ Rate significantly lower than the state rate. Number of the cases and rates are averaged over the five-year period. Rates per 100,000 age-adjusted to 2000 US standard population and SD 2018 estimated population. Source: South Dakota Department of Health

	Colorectal		Lung and Bronchus		Female Breast		Prostate		Bladder		NHL	
	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
<b>South Dakota</b>	<b>409</b>	<b>38.5</b>	<b>597</b>	<b>51.9</b>	<b>654</b>	<b>125.2</b>	<b>690</b>	<b>119.5</b>	<b>203</b>	<b>18.3</b>	<b>191</b>	<b>17.7</b>
Aurora	*	21.9	*	47.9	6	464.6	*	85.2	*	49.8	3	95.7
Beadle	6	22.5	12	46.4	8	60.4	26	201.0	4	12.6	*	8.8
Bennett	*	23.5	3	65.4	*	91.0	*	48.5	0	0.0	*	44.0
Bon Homme	*	9.4	5	46.1	6	155.0	12	223.2	4	44.4	*	9.4
Brookings	9	30.2	13	39.0	17	108.8	29	171.4	8	25.8	5	16.8
Brown	16	35.4	28	55.9	36	153.3	38	145.6	9	19.6	9	18.2
Brule	3	32.5	6	79.2	3	69.2	*	50.7	3	34.6	*	18.4
Buffalo	*	132.1	*	93.1	*	110.1	4	743.2	0	0.0	0	0.0
Butte	4	30.6	7	37.2	11	171.9	6	74.3	3	27.0	*	5.0
Campbell	0	0.0	0	0.0	*	79.4	0	0.0	0	0.0	0	0.0
Charles Mix	8	58.9	3	21.8	8	143.8	6	94.3	5	36.7	*	7.5
Clark	6	114.4	7	113.1	*	26.6	5	220.3	0	0.0	0	0.0
Clay	5	51.2	8	59.3	6	119.7	10	147.2	*	11.5	3	28.2
Codington	14	37.2	29	78.8	17	88.0	26	139.0	8	20.5	*	5.2
Corson	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Custer	4	15.1	4	21.7	8	97.3	9	83.9	3	13.2	*	7.0
Davison	7	28.9	21	71.9	20	178.6	15	109.7	6	19.0	5	20.8
Day	6	77.7	4	41.5	6	130.9	8	181.0	0	0.0	*	19.5
Deuel	*	10.8	*	11.7	*	34.6	*	63.2	0	0.0	*	27.4
Dewey	*	13.9	*	42.4	6	252.0	5	195.1	*	27.8	0	0.0
Douglas	0	0.0	*	19.7	*	88.9	6	264.1	*	18.6	*	54.4
Edmunds	0	0.0	*	28.7	4	176.0	8	226.6	0	0.0	*	48.7
Fall River	4	24.3	14	107.5	7	158.9	7	78.9	*	15.1	3	18.0
Faulk	0	0.0	4	80.3	4	268.5	*	71.1	0	0.0	*	65.7
Grant	4	39.1	6	47.3	3	57.8	7	109.1	4	34.0	3	22.6
Gregory	3	38.0	*	15.0	*	75.8	5	143.8	5	73.4	*	24.0
Haakon	*	54.8	*	51.9	6	482.0	0	0.0	0	0.0	0	0.0
Hamlin	*	25.6	5	66.1	3	101.5	4	122.4	0	0.0	*	32.2
Hand	*	30.1	0	0.0	*	12.8	3	89.1	3	42.2	0	0.0
Hanson	*	26.6	*	24.8	5	268.2	*	55.0	*	77.0	0	51.7
Harding	0	0.0	0	0.0	*	90.2	*	113.5	0	0.0	0	0.0
Hughes	7	28.5	9	38.7	11	105.2	17	152.4	6	25.5	5	26.5
Hutchinson	9	93.1	8	58.5	10	180.7	8	141.7	4	22.2	0	0.0
Hyde	*	29.8	*	84.6	4	164.9	*	141.1	0	0.0	0	0.0
Jackson	*	37.5	0	0.0	4	206.2	*	61.5	*	28.3	*	37.5
Jerauld	*	41.4	*	30.7	*	189.8	*	38.1	*	30.7	0	0.0
Jones	*	109.9	0	0.0	*	209.1	0	0.0	0	0.0	*	59.5
Kingsbury	7	72.7	5	52.7	8	186.7	8	182.9	3	34.4	4	45.9
Lake	8	38.8	10	51.1	7	74.3	12	105.3	3	23.8	3	25.6
Lawrence	10	25.9	23	53.0	23	117.9	16	66.5	3	7.1	3	6.4
Lincoln	24	40.1	23	37.0	42	128.8	41	140.2	7	10.6	13	24.3
Lyman	5	125.3	*	58.9	*	65.5	*	35.0	*	18.1	*	21.2
McCook	5	75.9	4	50.8	11	230.6	3	77.9	*	12.7	5	63.4
McPherson	3	75.6	*	44.9	3	143.2	*	45.1	0	0.0	0	0.0
Marshall	*	12.8	*	17.4	3	77.4	8	156.2	4	46.9	*	27.5
Meade	9	29.7	17	52.3	18	132.7	15	81.0	*	3.4	*	2.5
Mellette	0	0.0	0	0.0	2	238.2	*	51.1	0	0.0	*	43.1
Miner	0	0.0	3	100.0	3	232.5	*	127.4	0	0.0	0	0.0
Minnehaha	80	41.2	132	62.3	135	135.4	173	165.9	45	22.8	36	18.1
Moody	3	32.2	8	83.8	3	81.7	*	17.4	0	0.0	*	23.6
Oglala Lakota	5	47.5	4	44.9	4	63.9	6	136.3	0	0.0	*	8.3
Pennington	56	41.7	77	50.6	89	120.8	55	69.1	22	14.9	23	13.2
Perkins	*	17.9	*	22.1	*	36.1	*	33.4	0	0.0	0	0.0
Potter	4	118.2	3	64.3	*	50.4	*	84.2	*	13.8	*	27.6
Roberts	3	20.7	4	24.0	3	37.1	9	134.4	*	12.6	6	40.1
Sanborn	*	86.0	4	113.3	*	41.8	*	102.3	*	21.8	*	18.9
Spink	5	50.0	7	71.5	5	120.5	4	93.4	*	8.2	3	25.5
Stanley	*	16.9	0	0.0	7	498.4	*	30.3	*	20.5	*	21.8
Sully	*	72.2	*	27.6	*	99.1	*	44.5	0	0.0	0	0.0
Todd	*	23.6	7	125.1	*	44.4	*	37.4	*	46.0	0	0.0
Tripp	3	29.9	5	52.9	5	126.7	5	116.4	*	6.6	*	20.8
Turner	5	42.5	11	76.7	4	73.2	3	49.8	*	13.0	*	8.2
Union	15	72.6	13	58.8	15	164.6	15	131.8	6	29.9	8	36.8
Walworth	3	48.5	3	29.4	4	95.4	3	58.8	0	0.0	*	29.5
Yankton	10	36.0	13	38.7	19	115.0	20	119.6	9	27.7	5	15.8
Ziebach	*	111.5	*	96.8	*	44.9	*	78.2	0	0.0	0	0.0

**Table 2: Age-adjusted Incidence Rates by County for Selected Sites, 2018**

Note: \* Counts less than 3 are suppressed. Incidence rates with counts less than 20 are generally considered unstable.

Rates per 100,000 age-adjusted to the 2000 US standard population and 2018 SD estimated population. Source: South Dakota Department of Health

**Table 3: Age-adjusted Incidence Rates by Site, Gender and Race, South Dakota, 2018**

	TOTAL		MALE		FEMALE		WHITE		AMERICAN INDIAN	
	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
<b>Total</b>	<b>4,820</b>	<b>446.7</b>	<b>2,560</b>	<b>482.9</b>	<b>2,260</b>	<b>422.1</b>	<b>4,517</b>	<b>451.7</b>	<b>248</b>	<b>499.8</b>
<b>Oral Cavity</b>	<b>123</b>	<b>11.5</b>	<b>90</b>	<b>17.4</b>	<b>33</b>	<b>5.9</b>	<b>118</b>	<b>11.8</b>	<b>5</b>	<b>8.9</b>
Lip	10	0.9	8	1.7	2	0.4	10	1.0	0	0.0
Tongue	36	3.4	28	5.4	8	1.5	35	3.5	1	2.1
Salivary Gland	12	1.1	7	1.4	5	0.9	12	1.2	0	0.0
Floor of Mouth	4	0.4	1	0.2	3	0.5	4	0.4	0	0.0
Gum and Other Mouth	17	1.5	6	1.1	11	1.9	17	1.6	0	0.0
Nasopharynx	6	0.6	6	1.3	0	0.0	5	0.5	1	1.7
Tonsil	31	2.8	28	4.9	3	0.6	29	2.7	2	3.7
Oropharynx	3	0.3	3	0.7	0	0.0	3	0.4	0	0.0
Hypopharynx	4	0.4	3	0.6	1	0.1	3	0.4	1	1.4
Other Oral Cavity & Pharynx	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
<b>Digestive System</b>	<b>853</b>	<b>78.3</b>	<b>480</b>	<b>91.9</b>	<b>373</b>	<b>65.4</b>	<b>793</b>	<b>78.6</b>	<b>49</b>	<b>102.5</b>
Esophagus	67	6.0	56	10.5	11	1.9	59	5.7	7	12.9
Stomach	47	4.5	34	6.9	13	2.4	43	4.4	2	3.2
Small Intestine	41	3.9	20	3.7	21	4.2	40	4.2	1	1.7
Colorectal	409	38.5	224	44.6	185	32.7	390	39.7	17	36.3
<b>Colon Excluding Rectum</b>	299	27.7	157	31.5	142	24.2	289	28.8	9	20.2
<b>Rectum and Rectosigmoid</b>	110	10.8	67	13.1	43	8.5	101	10.9	8	16.0
Anus, Anal Canal and Anorectum	28	2.4	8	1.4	20	3.3	27	2.4	1	3.6
Liver & Intrahepatic Bile Duct	66	5.5	50	8.6	16	2.6	54	4.8	7	15.5
Gallbladder	11	0.9	2	0.3	9	1.5	9	0.8	2	4.0
Other Biliary	18	1.6	12	2.3	6	1.0	17	1.6	1	3.2
Pancreas	147	12.9	69	12.4	78	13.3	135	12.7	11	22.1
Retroperitoneum	5	0.6	2	0.5	3	0.6	5	0.6	0	0.0
Peritoneum, Omentum and Mesentery	14	1.4	3	0.6	11	2.0	14	1.5	0	0.0
<b>Respiratory</b>	<b>632</b>	<b>55.0</b>	<b>339</b>	<b>62.5</b>	<b>293</b>	<b>49.6</b>	<b>588</b>	<b>54.1</b>	<b>39</b>	<b>89.9</b>
Nose, Nasal Cavity and Middle Ear	3	0.3	3	0.6	0	0.0	3	0.4	0	0.0
Larynx	31	2.7	26	4.8	5	0.8	23	2.1	7	18.3
Lung and Bronchus	597	51.9	310	57.1	287	48.7	561	51.5	32	71.6
Pleura	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Mediastinum and Other Resp Organs	1	0.1	0	0.0	1	0.1	1	0.1	0	0.0
<b>Bones and Joints</b>	<b>3</b>	<b>0.3</b>	<b>2</b>	<b>0.4</b>	<b>1</b>	<b>0.3</b>	<b>3</b>	<b>0.4</b>	<b>0</b>	<b>0.0</b>
<b>Soft Tissue (Including Heart)</b>	<b>27</b>	<b>2.6</b>	<b>16</b>	<b>3.2</b>	<b>11</b>	<b>2.1</b>	<b>27</b>	<b>2.9</b>	<b>0</b>	<b>0.0</b>
<b>Skin</b>	<b>326</b>	<b>31.6</b>	<b>183</b>	<b>35.4</b>	<b>143</b>	<b>29.3</b>	<b>323</b>	<b>34.3</b>	<b>2</b>	<b>5.4</b>
Melanomas of the Skin	304	29.5	173	33.2	131	27.1	303	32.4	1	2.1
Other Skin	22	2.1	10	2.2	12	2.1	20	1.9	1	3.2
<b>Breast</b>	<b>657</b>	<b>64.2</b>	<b>3</b>	<b>0.6</b>	<b>654</b>	<b>125.2</b>	<b>619</b>	<b>65.8</b>	<b>29</b>	<b>56.0</b>
Breast, Female	654	125.2			654	125.2	616	128.7	29	101.7
Breast, Male	3	0.6	3	0.6			3	0.6	0	0.0
<b>Female</b>	<b>260</b>	<b>47.8</b>			<b>260</b>	<b>47.8</b>	<b>238</b>	<b>47.2</b>	<b>19</b>	<b>66.9</b>
Vulva	25	4.5			25	4.5	24	4.7	1	3.0
Vagina	5	1.0			5	1.0	4	0.7	0	0.0
Cervix Uteri	21	5.2			21	5.2	17	4.8	4	16.1
Corpus and Uterus, NOS	143	24.6			143	24.6	131	24.1	11	38.3
<b>Corpus Uteri</b>	141	24.1			141	24.1	129	23.5	11	38.3
<b>Uterus, NOS</b>	2	0.4			2	0.4	2	0.5	0	0.0
Ovary	51	9.8			51	9.8	48	10.2	3	9.5
Other Female Genital Organs	15	2.7			15	2.7	14	2.7	0	0.0

**Table 3: Age-adjusted Incidence Rates by Site, Gender and Race, South Dakota, 2018 (continued)**

	TOTAL		MALE		FEMALE		WHITE		AMERICAN INDIAN	
	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
<b>Male</b>	<b>718</b>	<b>126.0</b>	<b>718</b>	<b>126.0</b>			<b>665</b>	<b>124.4</b>	<b>42</b>	<b>185.5</b>
Penis	7	1.5	7	1.5			5	1.1	1	5.3
Prostate	690	119.5	690	119.5			640	117.6	40	178.2
Testis	19	4.6	19	4.6			18	5.2	1	2.0
Other Male Genital Organs	2	0.4	2	0.4			2	0.5	0	0.0
<b>Urinary</b>	<b>386</b>	<b>36.0</b>	<b>291</b>	<b>57.0</b>	<b>95</b>	<b>18.4</b>	<b>364</b>	<b>36.1</b>	<b>20</b>	<b>38.7</b>
Bladder	203	18.3	165	32.9	38	6.6	200	19.2	2	5.9
Kidney and Renal Pelvis	180	17.4	124	23.8	56	11.6	161	16.7	18	32.9
Ureter	2	0.2	1	0.1	1	0.2	2	0.2	0	0.0
Other Urinary Organs	1	0.1	1	0.2	0	0.0	1	0.1	0	0.0
<b>Eye and Orbit</b>	<b>5</b>	<b>0.5</b>	<b>3</b>	<b>0.5</b>	<b>2</b>	<b>0.5</b>	<b>5</b>	<b>0.6</b>	<b>0</b>	<b>0.0</b>
<b>Brain and CNS</b>	<b>46</b>	<b>4.8</b>	<b>33</b>	<b>7.4</b>	<b>13</b>	<b>2.5</b>	<b>43</b>	<b>5.1</b>	<b>3</b>	<b>2.3</b>
Brain	45	4.7	33	7.4	12	2.2	43	5.1	2	1.5
Meninges and CNS	1	0.1	0	0.0	1	0.2	0	0.0	1	0.8
<b>Endocrine</b>	<b>143</b>	<b>16.3</b>	<b>37</b>	<b>8.0</b>	<b>106</b>	<b>25.1</b>	<b>127</b>	<b>16.2</b>	<b>13</b>	<b>23.8</b>
Thyroid	138	15.7	36	7.8	102	24.2	122	15.6	13	23.8
Other Endocrine	5	0.5	1	0.2	4	0.9	5	0.6	0	0.0
<b>Lymphomas</b>	<b>214</b>	<b>20.2</b>	<b>119</b>	<b>23.2</b>	<b>95</b>	<b>17.6</b>	<b>204</b>	<b>21.1</b>	<b>8</b>	<b>12.2</b>
Hodgkin's Lymphoma	23	2.6	14	2.9	9	2.3	21	2.7	2	3.0
Non-Hodgkin's Lymphoma	191	17.7	105	20.3	86	15.3	183	18.4	6	9.2
<b>Multiple Myeloma</b>	<b>87</b>	<b>8.1</b>	<b>49</b>	<b>9.4</b>	<b>38</b>	<b>7.0</b>	<b>80</b>	<b>8.0</b>	<b>2</b>	<b>3.4</b>
<b>Leukemia</b>	<b>166</b>	<b>16.1</b>	<b>98</b>	<b>19.9</b>	<b>68</b>	<b>12.9</b>	<b>158</b>	<b>16.6</b>	<b>8</b>	<b>17.9</b>
Acute Lymphocytic	14	1.7	8	1.8	6	1.5	11	1.8	3	4.5
Chronic Lymphocytic	80	7.0	52	9.8	28	4.5	80	7.5	0	0.0
Other Lymphocytic	2	0.2	2	0.4	0	0.0	2	0.2	0	0.0
Acute Myeloid	46	4.6	28	6.0	18	3.4	43	4.6	3	7.7
Acute Monocytic	1	0.1	1	0.2	0	0.0	1	0.1	0	0.0
Chronic Myeloid	16	1.8	2	0.4	14	3.0	14	1.7	2	5.6
Other Myeloid/Monocytic	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other Acute Leukemia	2	0.2	2	0.5	0	0.0	2	0.2	0	0.0
Other Leukemia	5	0.5	3	0.7	2	0.4	5	0.5	0	0.0
<b>Myeloproliferative Myelodysplastic</b>	<b>90</b>	<b>8.6</b>	<b>55</b>	<b>11.7</b>	<b>35</b>	<b>6.1</b>	<b>81</b>	<b>8.3</b>	<b>6</b>	<b>17.4</b>
<b>Mesothelioma</b>	<b>4</b>	<b>0.4</b>	<b>3</b>	<b>0.6</b>	<b>1</b>	<b>0.1</b>	<b>4</b>	<b>0.4</b>	<b>0</b>	<b>0.0</b>
<b>Immunoproliferative Diseases</b>	<b>8</b>	<b>0.7</b>	<b>6</b>	<b>1.1</b>	<b>2</b>	<b>0.3</b>	<b>8</b>	<b>0.7</b>	<b>0</b>	<b>0.0</b>
<b>Other Sites</b>	<b>72</b>	<b>6.2</b>	<b>35</b>	<b>6.7</b>	<b>37</b>	<b>6.2</b>	<b>69</b>	<b>6.4</b>	<b>3</b>	<b>4.6</b>

Rates per 100,000 age-adjusted to the 2000 US standard population and 2018 SD estimated population.

Source: South Dakota Department of Health

Table 3 above shows incidence and age-adjusted incidence rates for South Dakota in 2018 by primary sites, gender, and race according to SEER site category recodes (Appendix C).

**Table 4: Percentage of Selected Cancers by Age Groups in South Dakota, 2018**


	0-19	20-34	35-49	50-64	65-74	75-84	85+
<b>All Sites</b>	1%	2%	8%	29%	33%	19%	9%
Bladder	0%	0%	4%	17%	31%	30%	18%
Breast, Female	0%	1%	15%	32%	29%	17%	7%
Colorectal	0%	2%	7%	26%	31%	19%	15%
Corpus and Uterus, NOS	0%	1%	7%	42%	33%	11%	6%
Hodgkin's Lymphoma	4%	35%	17%	17%	22%	4%	0%
Kidney and Renal Pelvis	1%	1%	14%	34%	24%	17%	8%
Leukemia	7%	2%	7%	24%	23%	25%	11%
Lung and Bronchus	0%	0%	1%	26%	40%	24%	9%
Melanomas of the Skin	0%	5%	15%	31%	28%	13%	9%
Non-Hodgkin's Lymphoma	2%	3%	5%	31%	29%	21%	9%
Pancreas	0%	1%	3%	28%	35%	22%	11%
Prostate	0%	0%	1%	34%	46%	15%	4%
Thyroid	0%	14%	30%	33%	16%	7%	1%

Note: Detail may not sum to totals because of rounding. Source: South Dakota Department of Health

In 2018, 62% of all cancers were diagnosed between ages 50 to 74 (Table 4). Notable were the 30% of thyroid and 17% of Hodgkin's lymphoma cancers diagnosed between the ages of 35 to 49. In 2018, 39% of Hodgkin's lymphoma cases were diagnosed in persons under 35 years old, compared to 59% in 2017.

Figure 1 below shows that incidence rates for American Indians in South Dakota were higher than those for whites in 2018. Of the 4,820 newly diagnosed cases in 2018, 248 or 5.1% were American Indians, 123 males, and 125 females.

**Figure 1: All Sites Cancer Incidence Rates by Race and Gender, South Dakota, 2018**




Note: Rates per 100,000 age-adjusted to the 2000 US standard population and 2018 SD estimated population. Source: South Dakota Department of Health

## V. CANCER CASES AND DEATHS BY RANK


Prostate cancer was the most common cancer diagnosed during 2018. The other four most diagnosed cancers were female breast, lung and bronchus, colorectal, and melanoma, which accounted for 55.1% of the new cases diagnosed and 48.2% of cancer deaths. Figure 2 shows the percent of new cancer cases and deaths by rank and gender.

**Figure 2: Percent Distribution of Cancer Cases and Deaths by Rank and Gender, South Dakota, 2018**



Source: South Dakota Department of Health


Figure 3: Cancer Incidence Rates by County, South Dakota, 2018



Note: Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. Source: South Dakota Department of Health

South Dakota has an area of 77,121 square miles with a 2018 estimated population of 882,235 persons resulting in a population density of 11 persons per square mile. Population densities range from a low of 0.5 in Harding County to a high of 210.0 persons per square mile in Minnehaha County.

Figure 4: Cancer Mortality Rates by County, South Dakota, 2018



Note: Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. Source: South Dakota Department of Health


South Dakota has an area of 77,121 square miles with a 2018 estimated population of 882,235 persons resulting in a population density of 11 persons per square mile. Population densities range from a low of 0.5 in Harding County to a high of 210.0 persons per square mile in Minnehaha County.

## VI. STAGE AT DIAGNOSIS

### SEER Summary Staging:

- ***In Situ*** - Malignant cells are within the cell groups from which they arose, without penetration of the basement membrane of the tissue and stromal invasion.
- **Localized** - The malignant cells are limited to the organ of origin and have spread no farther than the organ where they began.
- **Regional** - The tumor is beyond the limits of the organ of origin by direct extension to adjacent areas such as the regional lymph nodes, adjacent organs, or tissue.
- **Distant** - The tumor cells have broken away from the primary tumor and traveled to other parts of the body.
- **Unknown** - If extension or metastatic, there is not sufficient evidence available to assign a stage.

**Figure 5: Stage at Diagnosis, All Sites, South Dakota, 2018**



Source: South Dakota Department of Health


The figure above demonstrates the percentage of cases diagnosed at each stage of disease. For 2018, there were a total of 5,398 cases reported to the South Dakota State Cancer Registry, which includes *in situ* cases. The number of cases is in Table 5.

**Table 5: South Dakota Stage at Diagnosis, All Cases, 2018**

Stage	Number of Cases	Percent of Total
<i>In Situ</i>	695	13%
Localized	2296	43%
Regional	979	18%
Distant	1118	21%
Unknown	310	5%


When analyzed by race, 43% of all cancer cases diagnosed for whites were diagnosed at localized stage and 44% for American Indians. American Indians had another 43% diagnosed at the regional and distant stages combined, while the same stages for whites were a combined 39%. The stage at diagnosis by race was as follows:

**Figure 6: Stage at Diagnosis in South Dakota, White, 2018**



Source: South Dakota Department of Health

**Figure 7: Stage at Diagnosis in South Dakota, American Indians, 2018**



Source: South Dakota Department of Health

Cases of non-Hodgkin’s lymphoma, myeloma, and leukemias are usually at distant stages, and therefore can skew the proportion of all sites diagnosed at distant stages. Some differences in case counts by stage for selected sites are shown in Table 6.

**Table 6: Stage at Diagnosis for Selected Sites by Race, South Dakota, 2018**

	White						American Indian					
	Localized		Regional		Distant		Localized		Regional		Distant	
	Cases	%	Cases	%	Cases	%	Cases	%	Cases	%	Cases	%
Female Breast	422	19.5%	152	16.8%	31	3.0%	18	16.1%	9	17.0%	0	0.0%
Lung and Bronchus	177	8.2%	108	11.9%	242	23.1%	7	6.3%	7	13.2%	12	21.4%
Prostate	488	22.6%	89	9.8%	35	3.3%	29	25.9%	4	7.5%	4	7.1%
Melanoma of the Skin	258	11.9%	26	2.9%	11	1.0%	1	0.9%	0	0.0%	0	0.0%
Colorectal	132	6.1%	165	18.2%	69	6.6%	6	5.4%	9	17.0%	1	1.8%
Bladder	61	2.8%	11	1.2%	7	0.7%	1	0.9%	1	1.9%	0	0.0%
Non-Hodgkin’s Lymphoma	48	2.2%	15	1.7%	103	9.8%	2	1.8%	0	0.0%	3	5.4%
Kidney and Renal Pelvis	110	5.1%	17	1.9%	24	2.3%	14	12.5%	2	3.8%	1	1.8%
Corpus and Uterus, NOS	103	4.8%	17	1.9%	7	0.7%	9	8.0%	1	1.9%	0	0.0%
Leukemia	6	0.3%	1	0.1%	149	14.2%	0	0.0%	0	0.0%	8	14.3%
Thyroid	88	4.1%	31	3.4%	2	0.2%	5	4.5%	6	11.3%	1	1.8%
Pancreas	21	1.0%	44	4.9%	58	5.5%	2	1.8%	3	5.7%	6	10.7%

Source: South Dakota Department of Health



## VII. CANCER MORTALITY

Cancer age-adjusted mortality rates for 2018 ranged from a low of 0.0 in Harding County to a high of 535.2 in Buffalo County. South Dakota's age-adjusted mortality rate was 147.9 in 2018 compared to a five-year mortality rate of 156.5.

In 2018, five counties had a significantly lower rate than the entire state and two had a significantly higher rate. The five-year rates show six counties having significantly lower rates and six counties with significantly higher rates. South Dakota's mortality rate for 2014-2018 was 156.5 per 100,000 persons.

The United States mortality rate in 2018 was 149.0 and the South Dakota rate was 147.9 per 100,000 persons. When comparing the two rates there was not a significant difference.

The South Dakota 2018 mortality rates are displayed in a state map on page 15 of this report.

Table 7 : Cancer Deaths and Mortality Rates by County South Dakota, 2018 and 2014-2018 Average				
County	2018		2014-2018 <sup>^</sup>	
	Deaths	Rate	Deaths	Rate
<b>South Dakota</b>	1,661	147.9	1,689	156.5
Aurora	11	236.1	7	159.7
Beadle	36	142.9	42	171.6
Bennett	3	67.4	7	201.9
Bon Homme	14	125.2	17	145.2
Brookings	38	118.3	41	138.3
Brown	70	130.1	81	157.9
Brule	6	66.8 ▼	11	130.6
Buffalo	7	535.2	5	333.7 ▲
Butte	24	146.2	26	175.2
Campbell	*	37.4 ▼	3	109.8
Charles Mix	25	198.3	25	190.3
Clark	10	192.7	10	170.7
Clay	20	150.2	22	172.4
Codington	53	139.1	63	170.0
Corson	6	141.0	8	196.7
Custer	23	119.6	20	121.1 ▼
Davison	42	164.4 ▼	47	161.8
Day	20	165.9	15	152.6
Deuel	12	170.4	12	167.0
Dewey	10	201.8	9	198.5
Douglas	4	41.9 ▼	8	135.2
Edmunds	8	141.3	7	102.0 ▼
Fall River	28	190.9	25	182.6
Faulk	8	207.5	6	140.3
Grant	13	107.5	16	140.2
Gregory	9	113.2	14	172.8
Haakon	3	86.4	5	149.6
Hamlin	12	137.7	12	144.9
Hand	8	99.8	8	125.0
Hanson	8	339.5	9	322.0 ▲
Harding	0	0.0	*	76.1 ▼
Hughes	26	120.4	31	141.5
Hutchinson	19	142.5	19	135.3
Hyde	3	105.0	4	139.7
Jackson	3	78.5	7	188.1
Jerauld	3	72.8	6	155.6
Jones	3	147.0	3	184.6
Kingsbury	12	136.2	16	190.3
Lake	29	160.7	26	137.6
Lawrence	56	144.8	49	137.4
Lincoln	63	105.9 ▼	61	117.6 ▼
Lyman	13	275.8	9	198.3
McCook	14	159.8	18	228.6 ▲
McPherson	4	82.9	7	111.1 ▼
Marshall	4	52.6 ▼	9	132.1
Meade	61	190.8	53	173.1
Mellette	10	429.7 ▲	5	222.0
Miner	6	179.8	8	207.9
Minnehaha	343	174.6 ▲	323	169.3 ▲
Moody	13	147.2	11	134.3
Oglala Lakota	20	253.7	18	210.5 ▲
Pennington	216	144.1	212	153.1
Perkins	7	120.3	9	164.0
Potter	5	96.6	7	122.1
Roberts	25	168.3	24	163.7
Sanborn	6	169.7	5	138.8
Spink	15	159.2	15	149.6
Stanley	4	93.8	7	172.9
Sully	3	100.2	3	126.4
Todd	17	275.2	15	237.3 ▲
Tripp	11	101.9	13	127.5
Turner	21	160.4	21	164.7
Union	28	142.5	31	158.3
Walworth	20	217.6	15	153.0
Yankton	40	121.1	44	132.5 ▼
Ziebach	6	288.0	*	119.7

\* Counts less than three are suppressed. Mortality rates with counts less than 20 are generally considered unstable. ▲ Rate significantly higher; ▼ Rate significantly lower

<sup>^</sup> Number of the cases and rates are averaged over the five-year period.

Rates per 100,000 age-adjusted to 2000 US standard population and SD 2018 estimated population. Source: South Dakota Department of Health

**Table 8: Age-adjusted Mortality Rates by County for Selected Sites, 2018**

	Colorectal		Lung and Bronchus		Female Breast		Prostate		Bladder		NHL	
	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate
<b>South Dakota</b>	<b>168</b>	<b>15.1</b>	<b>395</b>	<b>35.1</b>	<b>112</b>	<b>19.8</b>	<b>92</b>	<b>19.1</b>	<b>34</b>	<b>3.0</b>	<b>56</b>	<b>5.0</b>
Aurora	*	39.9	*	21.9	*	58.4	*	44.5	0	0.0	3	65.1
Beadle	*	10.9	6	20.7	3	15.3	4	39.0	0	0.0	0	0.0
Bennett	0	0.0	*	21.0	*	44.6	*	48.5	0	0.0	0	0.0
Bon Homme	*	8.5	*	27.9	0	0.0	0	0.0	0	0.0	0	0.0
Brookings	4	12.5	10	33.0	*	6.3	*	7.5	*	4.9	*	5.9
Brown	14	26.6	13	23.3	5	19.7	*	7.4	0	0.0	*	3.3
Brule	*	9.7	*	23.8	*	13.1	0	0.0	0	0.0	*	8.7
Buffalo	3	251.4	2	129.4	0	0.0	0	0.0	0	0.0	0	0.0
Butte	0	0.0	9	56.7	0	0.0	*	27.3	0	0.0	0	0.0
Campbell	0	0.0	*	37.4	0	0.0	0	0.0	0	0.0	0	0.0
Charles Mix	6	51.2	3	23.3	3	36.4	*	14.4	*	5.2	*	26.2
Clark	0	0.0	*	32.6	0	0.0	*	51.0	0	0.0	*	19.1
Clay	*	12.5	4	31.4	*	15.1	*	43.8	0	0.0	*	7.0
Codington	7	20.8	10	24.1	5	18.4	*	5.2	*	4.5	3	8.5
Corson	0	0.0	0	0.0	*	57.4	0	0.0	0	0.0	*	24.8
Custer	*	11.4	8	42.9	*	9.9	*	11.6	0	0.0	*	6.7
Davison	4	17.4	11	41.9	6	46.6	*	9.4	*	6.0	*	4.3
Day	*	6.4	5	41.8	*	9.5	0	0.0	*	17.8	*	8.0
Deuel	*	10.8	3	35.5	0	0.0	*	27.9	0	0.0	0	0.0
Dewey	0	0.0	*	47.1	*	50.4	*	63.7	0	0.0	0	0.0
Douglas	0	0.0	0	0.0	*	14.8	*	36.1	0	0.0	0	0.0
Edmunds	*	13.7	0	0.0	*	44.8	0	0.0	0	0.0	0	0.0
Fall River	3	19.9	12	86.9	0	0.0	*	13.3	0	0.0	*	5.2
Faulk	0	0.0	4	107.4	0	0.0	0	0.0	0	0.0	0	0.0
Grant	0	0.0	3	23.6	0	0.0	*	41.7	0	0.0	*	6.9
Gregory	0	0.0	0	0.0	0	0.0	*	53.1	0	0.0	*	12.7
Haakon	0	0.0	*	27.4	0	0.0	0	0.0	0	0.0	*	28.7
Hamlin	*	11.7	4	47.3	*	13.4	0	0.0	0	0.0	0	0.0
Hand	0	0.0	*	13.4	0	0.0	*	51.2	*	7.9	0	0.0
Hanson	*	35.1	*	63.7	0	0.0	0	0.0	0	0.0	*	51.7
Harding	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Hughes	*	3.2	5	22.2	5	54.8	*	18.8	0	0.0	0	0.0
Hutchinson	*	16.9	6	48.2	0	0.0	*	19.5	0	0.0	*	9.7
Hyde	0	0.0	*	55.7	0	0.0	*	91.2	0	0.0	0	0.0
Jackson	0	0.0	*	28.3	0	0.0	0	0.0	*	26.3	0	0.0
Jerauld	*	15.2	*	30.7	0	0.0	0	0.0	0	0.0	0	0.0
Jones	0	0.0	0	0.0	*	110.5	0	0.0	0	0.0	0	0.0
Kingsbury	*	24.8	3	34.4	0	0.0	*	24.1	0	0.0	0	0.0
Lake	*	2.8	5	25.5	*	15.5	*	17.3	*	15.1	*	5.6
Lawrence	4	9.1	8	18.8	4	26.4	4	20.0	*	2.2	*	5.2
Lincoln	5	8.2	16	28.2	8	25.9	3	12.3	*	3.3	*	4.0
Lyman	*	43.3	6	129.0	*	77.6	0	0.0	0	0.0	0	0.0
McCook	*	12.8	4	50.6	*	26.7	*	33.7	0	0.0	0	0.0
McPherson	0	0.0	3	62.1	0	0.0	0	0.0	0	0.0	0	0.0
Marshall	0	0.0	0	0.0	0	0.0	0	0.0	*	9.5	0	0.0
Meade	7	22.2	18	55.3	5	36.8	*	7.8	*	5.5	*	5.0
Mellette	0	0.0	*	76.0	0	0.0	*	51.1	0	0.0	*	43.1
Miner	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Minnehaha	37	18.3	77	39.1	24	24.6	18	24.3	6	3.7	8	4.5
Moody	0	0.0	4	44.0	*	59.7	*	27.8	0	0.0	*	11.2
Oglala Lakota	*	33.8	4	60.9	0	0.0	*	37.5	0	0.0	0	0.0
Pennington	20	14.8	59	38.3	14	17.8	15	22.8	6	3.8	7	4.2
Perkins	0	0.0	*	10.2	0	0.0	*	39.7	0	0.0	*	22.1
Potter	*	13.8	*	13.8	0	0.0	*	55.1	0	0.0	0	0.0
Roberts	4	37.2	5	29.3	*	24.8	*	11.6	0	0.0	*	7.0
Sanborn	0	0.0	3	94.8	*	46.7	0	0.0	0	0.0	0	0.0
Spink	3	20.9	2	20.0	0	0.0	0	0.0	*	11.7	*	7.9
Stanley	*	20.5	0	0.0	*	68.6	*	38.0	0	0.0	0	0.0
Sully	0	0.0	*	27.6	0	0.0	*	77.5	0	0.0	0	0.0
Todd	*	23.6	3	64.5	0	0.0	2	88.4	*	22.3	0	0.0
Tripp	3	25.9	3	23.5	*	9.5	0	0.0	0	0.0	0	0.0
Turner	*	6.5	7	49.1	*	29.6	*	31.7	0	0.0	0	0.0
Union	3	13.4	5	27.4	*	8.6	*	23.5	0	0.0	*	4.6
Walworth	*	12.4	4	38.3	0	0.0	0	0.0	*	10.8	0	0.0
Yankton	5	18.5	15	46.9	*	3.2	*	11.6	*	3.3	3	8.3
Ziebach	3	133.7	*	57.4	0	0.0	0	0.0	0	0.0	0	0.0

Note: \* Counts less than 3 are suppressed. Mortality rates with counts less than 20 are generally considered unstable.  
 Rates per 100,000 age-adjusted to the 2000 US standard population and 2018 SD estimated population.  
 Source: South Dakota Department of Health

**Table 9: Age-adjusted Mortality Rates by Site, Gender, and Race, South Dakota, 2018**

	TOTAL		MALE		FEMALE		WHITE		AMERICAN INDIAN	
	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate
<b>Total</b>	<b>1,661</b>	<b>147.9</b>	<b>907</b>	<b>179.2</b>	<b>754</b>	<b>126.1</b>	<b>1,525</b>	<b>144.0</b>	<b>115</b>	<b>257.8</b>
<b>Oral Cavity</b>	<b>18</b>	<b>1.6</b>	<b>16</b>	<b>2.8</b>	<b>2</b>	<b>0.5</b>	<b>14</b>	<b>1.2</b>	<b>4</b>	<b>6.9</b>
Lip	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Tongue	5	0.4	4	0.6	1	0.2	4	0.3	1	1.7
Salivary Gland	2	0.2	2	0.5	0	0.0	2	0.2	0	0.0
Floor of Mouth	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Gum and Other Mouth	3	0.3	2	0.4	1	0.3	1	0.1	2	3.7
Nasopharynx	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Hypopharynx	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Tonsil	4	0.3	4	0.7	0	0.0	4	0.4	0	0.0
Oropharynx	4	0.3	4	0.7	0	0.0	3	0.3	1	1.4
Other Oral Cavity and Pharynx	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
<b>Digestive System</b>	<b>427</b>	<b>37.4</b>	<b>257</b>	<b>48.3</b>	<b>170</b>	<b>27.7</b>	<b>382</b>	<b>35.7</b>	<b>39</b>	<b>84.6</b>
Esophagus	49	4.3	41	7.7	8	1.3	46	4.3	3	6.9
Stomach	20	1.8	13	2.5	7	1.3	14	1.4	5	8.3
Small Intestine	7	0.6	4	0.8	3	0.6	5	0.5	0	0.0
Colorectal	168	15.1	92	18.1	76	12.4	153	14.8	13	31.1
Colon Excluding Rectum	123	10.8	63	12.4	60	9.3	111	10.5	11	28.3
Rectum and Rectosigmoid	45	4.2	29	5.7	16	3.1	42	4.3	2	2.9
Anus, Anal Canal and Anorectum	3	0.3	1	0.1	2	0.4	1	0.1	2	3.4
Liver and Intrahepatic Bile Duct	56	4.5	43	7.6	13	1.8	47	3.9	8	17.0
Gallbladder	3	0.3	1	0.2	2	0.4	2	0.2	1	1.7
Other Biliary	5	0.5	2	0.5	3	0.5	4	0.4	1	3.2
Pancreas	116	10.0	60	10.7	56	9.1	110	10.1	6	12.7
Retroperitoneum	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Peritoneum, Omentum and Mesentery	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
<b>Respiratory</b>	<b>404</b>	<b>35.9</b>	<b>219</b>	<b>42.6</b>	<b>185</b>	<b>31.3</b>	<b>371</b>	<b>34.7</b>	<b>27</b>	<b>66.5</b>
Nose, Nasal Cavity and Middle Ear	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Larynx	8	0.7	6	1.1	2	0.3	7	0.6	1	2.3
Lung and Bronchus	395	35.1	212	41.3	183	31.0	363	33.9	26	64.3
Pleura	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Mediastinum and Other Resp Organs	1	0.1	1	0.2	0	0.0	1	0.1	0	0.0
<b>Bones and Joints</b>	<b>4</b>	<b>0.4</b>	<b>3</b>	<b>0.6</b>	<b>1</b>	<b>0.2</b>	<b>4</b>	<b>0.5</b>	<b>0</b>	<b>0.0</b>
<b>Soft Tissue</b>	<b>14</b>	<b>1.3</b>	<b>6</b>	<b>1.5</b>	<b>8</b>	<b>1.2</b>	<b>13</b>	<b>1.3</b>	<b>0</b>	<b>0.0</b>
<b>Skin</b>	<b>46</b>	<b>4.4</b>	<b>35</b>	<b>7.5</b>	<b>11</b>	<b>2.1</b>	<b>45</b>	<b>4.6</b>	<b>1</b>	<b>1.4</b>
Melanoma of the Skin	34	3.4	25	5.3	9	1.9	33	3.6	1	1.4
Other Nonepithelial Skin	12	1.0	10	2.2	2	0.3	12	1.1	0	0.0
<b>Breast</b>	<b>113</b>	<b>10.7</b>	<b>1</b>	<b>0.2</b>	<b>112</b>	<b>19.8</b>	<b>106</b>	<b>10.8</b>	<b>5</b>	<b>13.0</b>
Breast, Female	112	19.8			112	19.8	106	20.2	5	20.7
Breast, Male	1	0.2	1	0.2			0	0.0	0	0.0
<b>Female</b>	<b>84</b>	<b>14.7</b>			<b>84</b>	<b>14.7</b>	<b>78</b>	<b>14.4</b>	<b>5</b>	<b>18.2</b>
Vulva	3	0.5			3	0.5	3	0.5	0	0.0
Vagina	1	0.1			1	0.1	1	0.1	0	0.0
Cervix Uteri	9	2.0			9	2.0	6	1.5	3	12.1
Corpus and Uterus, NOS	21	3.7			21	3.7	21	3.9	0	0.0
Corpus Uteri	10	1.7			10	1.7	10	1.9	0	0.0
Uterus, NOS	11	1.9			11	1.9	11	2.1	0	0.0
Ovary	48	8.0			48	8.0	46	8.2	2	6.1
Other Female Genital Organs	2	0.4			2	0.4	1	0.2	0	0.0

**Table 9: Age-adjusted Mortality Rates by Site, Gender, and Race, South Dakota, 2018  
(continued)**

	TOTAL		MALE		FEMALE		WHITE		AMERICAN INDIAN	
	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate	Deaths	Rate
<b>Male</b>	<b>94</b>	<b>19.8</b>	<b>94</b>	<b>19.8</b>			<b>84</b>	<b>18.4</b>	<b>9</b>	<b>44.6</b>
Penis	0	0.0	0	0.0			0	0.0	0	0.0
Prostate	92	19.1	92	19.1			83	18.1	8	40.1
Testis	2	0.7	2	0.7			1	0.4	1	4.4
Other Male Genital Organs	0	0.0	0	0.0			0	0.0	0	0.0
<b>Urinary</b>	<b>82</b>	<b>7.1</b>	<b>52</b>	<b>10.3</b>	<b>30</b>	<b>4.9</b>	<b>75</b>	<b>6.9</b>	<b>6</b>	<b>14.7</b>
Bladder	34	3.0	23	4.6	11	1.8	31	2.8	3	8.3
Kidney and Renal Pelvis	46	3.9	28	5.4	18	2.9	43	4.0	2	5.0
Ureter	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other Urinary Organs	2	0.1	1	0.2	1	0.1	1	0.1	1	1.4
<b>Eye and Orbit</b>	<b>2</b>	<b>0.2</b>	<b>2</b>	<b>0.4</b>	<b>0</b>	<b>0.0</b>	<b>2</b>	<b>0.2</b>	<b>0</b>	<b>0.0</b>
<b>Brain and CNS</b>	<b>49</b>	<b>4.9</b>	<b>37</b>	<b>7.8</b>	<b>12</b>	<b>2.1</b>	<b>43</b>	<b>4.8</b>	<b>5</b>	<b>9.3</b>
Brain	48	4.8	36	7.6	12	2.1	42	4.6	5	9.3
Meninges and CNS	1	0.1	1	0.2	0	0.0	1	0.1	0	0.0
<b>Endocrine</b>	<b>7</b>	<b>0.5</b>	<b>5</b>	<b>0.9</b>	<b>2</b>	<b>0.2</b>	<b>7</b>	<b>0.5</b>	<b>0</b>	<b>0.0</b>
Thyroid	7	0.5	5	0.9	2	0.2	7	0.5	0	0.0
Other Endocrine	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
<b>Lymphomas</b>	<b>58</b>	<b>5.2</b>	<b>33</b>	<b>6.8</b>	<b>25</b>	<b>3.9</b>	<b>54</b>	<b>5.0</b>	<b>3</b>	<b>5.5</b>
Hodgkin's Disease	2	0.2	1	0.2	1	0.2	2	0.2	0	0.0
Non-Hodgkin's Lymphomas	56	5.0	32	6.5	24	3.7	52	4.8	3	5.5
<b>Multiple Myeloma</b>	<b>39</b>	<b>3.3</b>	<b>23</b>	<b>4.8</b>	<b>16</b>	<b>2.4</b>	<b>39</b>	<b>3.5</b>	<b>0</b>	<b>0.0</b>
<b>Leukemia</b>	<b>58</b>	<b>5.1</b>	<b>37</b>	<b>7.4</b>	<b>21</b>	<b>3.3</b>	<b>54</b>	<b>5.0</b>	<b>4</b>	<b>12.0</b>
Acute Lymphocytic	1	0.1	1	0.1	0	0.0	1	0.1	0	0.0
Chronic Lymphocytic	14	1.1	9	1.9	5	0.6	13	1.1	1	3.6
Other Lymphocytic	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Acute Myeloid	30	2.6	20	3.9	10	1.7	30	2.8	0	0.0
Acute Monocytic	3	0.3	1	0.2	2	0.4	3	0.4	0	0.0
Chronic Myeloid	3	0.3	2	0.5	1	0.2	3	0.3	0	0.0
Other Myeloid/Monocytic	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other Acute Leukemia	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other Leukemia	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
<b>Myeloproliferative &amp; Myelodysplastic</b>	<b>29</b>	<b>2.6</b>	<b>16</b>	<b>3.4</b>	<b>13</b>	<b>2.2</b>	<b>27</b>	<b>2.5</b>	<b>2</b>	<b>6.9</b>
<b>Mesothelioma</b>	<b>4</b>	<b>0.4</b>	<b>2</b>	<b>0.5</b>	<b>2</b>	<b>0.4</b>	<b>4</b>	<b>0.4</b>	<b>0</b>	<b>0.0</b>
<b>III-Defined and Unspecified Sites</b>	<b>129</b>	<b>11.1</b>	<b>69</b>	<b>13.7</b>	<b>60</b>	<b>9.1</b>	<b>123</b>	<b>11.2</b>	<b>5</b>	<b>7.7</b>

Rates per 100,000 age-adjusted to the 2000 US standard population and 2018 SD estimated population.  
Source: South Dakota Department of Health

Table 9 shows death and age-adjusted mortality rates by SEER recode primary sites (Appendix D), gender and race. Approximately 1,700 persons die from cancer in South Dakota each year with a slight change in counts. Overall, more males than females die from cancer.


**Table 10: Percentage of Cancer Deaths by Age Groups and Selected Primary Sites, South Dakota, 2018**

Age Group	0-19	20-34	35-49	50-64	65-74	75-84	85+
All Sites	0%	0%	4%	21%	28%	25%	21%
Bladder	0%	0%	3%	12%	26%	24%	35%
Female Breast	0%	0%	10%	26%	17%	27%	21%
Colorectal	0%	0%	7%	19%	24%	24%	26%
Corpus and Uterus, NOS	0%	10%	0%	5%	38%	33%	14%
Meninges, Brain and CNS	6%	8%	12%	33%	18%	18%	4%
Kidney and Renal Pelvis	0%	0%	7%	24%	30%	11%	28%
Leukemia	2%	0%	3%	17%	24%	26%	28%
Lung and Bronchus	0%	0%	2%	23%	31%	28%	16%
Melanoma of the Skin	0%	0%	15%	21%	21%	26%	18%
Non-Hodgkin's Lymphoma	0%	2%	0%	7%	27%	32%	32%
Pancreas	0%	0%	3%	25%	36%	22%	14%
Prostate	0%	0%	0%	8%	20%	28%	45%

Source: South Dakota Department of Health

Overall, in 2018 more persons 65 to 74 years of age died from cancer in South Dakota than any other age group (Table 10).

**Figure 8: All Sites Cancer Mortality Rates by Race, and Gender, South Dakota, 2018**



Note: Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population.  
Source: South Dakota Department of Health

Figure 8 illustrates that American Indian males had a higher mortality rate than any other group. In 2012 American Indian females had a higher mortality rate than their male counterparts.

## VIII. YEARS OF POTENTIAL LIFE LOST


In 2018, cancer out ranked accidents in years of potential years lost (YPLL). For the five-year period 2014 to 2018, accidents ranked number one in YPLL. The number of years lost are shown in Figures 9 and 10 by race. There were 10,592 years of potential years lost due to cancer in 2018, compared to 9,904 years in 2008.

**Table 11: Leading Causes of Years of Potential Life Lost (to 75 years of age), South Dakota, 2018**

Malignant Neoplasms	10,592
Accidents	9,938
Heart Disease	8,365
Suicide & Self-Inflicted Injury	5,918
Chronic Liver Disease and Cirrhosis	3,711
Diabetes Mellitus	1,771
Chronic Lower Respiratory Diseases	1,540
Cerebrovascular Diseases	1,275
Homicide	1,243
Influenza and Pneumonia	886
Septicemia	637
All Other Causes	13,419


Source: South Dakota Department of Health

**Figure 9: Years of Potential Life Lost (YPLL) Before Age 75 for the Leading Causes of Death by Race, South Dakota, 2018**



Source: South Dakota Department of Health


**Figure 10: Years of Potential Life Lost (YPLL) Before Age 75 for the Leading Causes of Death by Race, South Dakota 2014-2018 Average\***



\* Number of years are averaged over the five-year period.  
Source: South Dakota Department of Health


The differences in YPLL in Figures 11 and 12 reflect the number of cancer cases by primary sites by race. For example, the YPLL in 2018 for lung and bronchus for whites was 2,011 for 202 deaths whereas the YPLL for American Indians was 178 years for 15 deaths. Not all cancers were present among the American Indian population during 2018, hence the differences in the cancer sites presented.

**Figure 11: Years of Potential Life Lost for Selected Cancers Among Whites, South Dakota, 2018**



Source: South Dakota Department of Health

**Figure 12: Years of Potential Life Lost for Selected Cancers Among American Indians, South Dakota, 2018**




Source: South Dakota Department of Health

## IX. AVERAGE YEARS OF LIFE LOST


South Dakota's average years of life lost (AYLL) due to cancer in 2018 was 11.8 years, a slight decrease from 12 years in 2017. Brain and nervous system cancer ranked first among cancer sites for American Indians at 36.3 years compared to whites where it ranked second with an average of 22.9 years.

**Figure 13: Average Years of Life Lost for Selected Cancers Among Whites, South Dakota, 2018**



Source: South Dakota Department of Health

**Figure 14: Average Years of Life Lost for Selected Cancers Among American Indians, South Dakota, 2018**



Source: South Dakota Department of Health

This page intentionally left blank.

## X. SELECTED CANCER SITES INCIDENCE AND MORTALITY

---

This section covers the following cancers: bladder, female breast, cervix uteri, colorectal cancer, corpus uterus, kidney and renal pelvis, leukemia, lung and bronchus, melanoma (skin), myeloma, non-Hodgkin's lymphoma, ovary, pancreas, prostate, stomach, and thyroid.

These cancers were selected because of the ranking in the cancer sites reported as well as the importance and impact to society.


Topics for each cancer include incidence and mortality data along with age-adjusted rates, trends, comparison with national data (if available), risks, associated risk factors, and prevention.

Table 12: Bladder Incidence and Mortality Summary, 2018

Bladder Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	203	165	38	34	23	11
		Age-Adjusted Rate	18.3	32.9	6.6	3.0	4.6	1.8
	White	# Cases / Deaths	200	164	36	31	22	9
		Age-Adjusted Rate	19.2	34.5	6.7	2.8	4.7	1.5
	American Indian	# Cases / Deaths	2	0	2	3	1	2
		Age-Adjusted Rate	5.9	0.0	10.3	8.3	3.0	11.9
United States	Total	Age-Adjusted Rate	19.0	32.8	8.4	4.3	7.1	2.0
	White	Age-Adjusted Rate	21.2	36.3	9.3	4.4	7.6	2.1
	American Indian	Age-Adjusted Rate	9.8	16.9	5.6	1.8	2.1	1.5

<sup>1</sup>Includes *in situ* bladder; rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source: South Dakota Department of Health

Figure 15: Bladder Cancer Stage at Diagnosis, South Dakota, 2018



Source: South Dakota Department of Health

**Descriptive Epidemiology**

**Stage at Diagnosis:** Cancer is categorized as noninvasive and invasive. There were 117 noninvasive bladder cancers and 88 invasive cancers reported in 2018. More than half, 58%, of bladder cancer cases in South Dakota were diagnosed at noninvasive, *in situ* stage. Nationally, 50% of the cases of urinary bladder cancer are diagnosed at the *in situ* stage. In South Dakota, 3% of the cases were not diagnosed until the disease had spread to distant sites. In the United States, distant stages accounted for 5% of bladder cancers reported.

**Incidence:** In 2018, it was estimated that almost 81,190 cases of bladder cancer would be diagnosed in the United States. There were


193 cases of bladder cancer reported in South Dakota. There were 155 men and 38 women diagnosed with bladder cancer in 2018. Statistically, men were diagnosed about four times as often as women. There were only five American Indian cases diagnosed in 2018. In the United States it was the sixth most frequent cancer. In South Dakota it was also the sixth most frequent cancer diagnosed.

**Mortality:** Advances in intravesical therapy and in the treatment of advanced disease with chemotherapy have reduced the percentage of mortality from bladder cancer. In South Dakota, ages from 75 and above have the highest mortality rate. In 2018, the South Dakota and US mortality rate was 4.3.

**Risk and Associated Factors:** Bladder cancer was one of the first malignancies associated with industrialization. Cigarette smoking increases the risk of bladder cancer by two times that of a nonsmoker. Work exposure to certain chemicals also increases risk. Those with the highest risk are makers of rubber, leather, textiles, paint products, and printing compounds.


**Prevention and Early Detection:** Avoiding exposure to chemicals and cigarette smoking are two of the most common recommendations for prevention.

**Figure 16: Bladder Cancer Number of Cases and Deaths by Age, South Dakota, 2018**



Source: South Dakota Department of Health


**Figure 17: Bladder Cancer Cases and Deaths by Year, South Dakota, 2001 - 2018**



Bladder cancer cases were at an all-time high in 2011 with 234 cases.

Source: South Dakota Department of Health

**Figure 18: Bladder Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health


## BREAST (FEMALE)

**Table 13: Female Breast Incidence and Mortality Summary, 2018**

Female Breast Cancer			Incidence	Mortality
South Dakota	Total	# Cases / Deaths	654	112
		Age-Adjusted Rate	125.2	19.8
	White	# Cases / Deaths	616	106
		Age-Adjusted Rate	128.7	20.2
	American Indian	# Cases / Deaths	29	5
		Age-Adjusted Rate	101.7	20.7
United States	Total	Age-Adjusted Rate	131.3	19.7
	White	Age-Adjusted Rate	133.5	19.2
	American Indian	Age-Adjusted Rate	89.9	10.6

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source: South Dakota Department of Health

**Figure 19: Female Breast Cancer Stage at Diagnosis, South Dakota, 2018**



Source: South Dakota Department of Health

### Descriptive Epidemiology

**Stage at Diagnosis:** Including *in situ* female breast cancer cases there were 809 cases diagnosed in 2018, of which 444 cases were diagnosed at a localized stage. This represents 55% of all reported breast cancer cases. There were 165 cases that had progressed beyond the breast. There were 31 that were diagnosed as a distant stage and 14 that were staged as unknown. The 155 *in situ* female breast cancer cases were reported but were not used in calculating incidence rates.

**Incidence:** National statistics report that female breast cancer is the most common malignant tumor among women. The incidence rate increased from 1947-1990. The rates fell 3.5% per year from 2001-2005. This decrease may be in part due to the lower number of women using hormone replacement therapy. South Dakota had 654 cases of invasive


female breast cancer reported in 2018. These cases represented 13.6% of all invasive cancer cases reported in 2018 versus 14.8% nationally. Looking at South Dakota women, breast cancer represented 28.9% of the cancer cases for woman diagnosed in 2018.

**Mortality:** Breast cancer is the fourth leading cause of death attributed to cancer in South Dakota. Nationwide, breast cancer mortality has been relatively stable overall since 1950. In cancers only of women, it is the second leading cause of cancer deaths. Although mortality has increased among women older than 55 years, it has decreased among women younger than 55. In 2018, there were 112 deaths. Of those deaths, 106 were white and five were American Indian.


**Risk and Associated Factors:** Studies have shown that breast cancer risk is due to a combination of factors. The main factors that influence your risk include being a woman and getting older. Most breast cancers are found in women who are 50 years old or older. Lifestyle factors that contribute to an increased risk include: drinking alcohol, being overweight or obese, and being physically inactive. If you have a family history of breast cancer or inherited changes in your BRCA1 and BRCA2 genes, you may have a high risk of getting breast cancer.

**Prevention and Early Detection:** Prevention and early detection is the key to the survival of breast cancer. Breast cancers found during mammograms are more likely to be smaller and still confined to the breast. The US Preventive Services Task Force (USPSTF) recommends biennial screening mammography for women aged 50 to 74 years.

**Figure 20: Female Breast Cancer Number of Cases and Deaths by Age, South Dakota, 2018**




Source: South Dakota Department of Health



**Figure 21: Female Breast Cancer Cases and Deaths by Year, South Dakota, 2001 - 2018**

The incidence of female breast cancer was at an all-time high in 2013.

Source: South Dakota Department of Health



**Figure 22: Female Breast Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**


Rates per 100,00 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health

**Table 14: Cervix Uteri Incidence and Mortality Summary, 2018**

Cervix Uteri Cancer			Incidence	Mortality
South Dakota	Total	# Cases / Deaths	21	9
		Age-Adjusted Rate	5.2	2.0
	White	# Cases / Deaths	17	6
		Age-Adjusted Rate	4.8	1.5
American Indian		# Cases / Deaths	4	3
		Age-Adjusted Rate	16.1	12.1
United States	Total	Age-Adjusted Rate	7.5	2.2
		White	Age-Adjusted Rate	7.3
	American Indian	Age-Adjusted Rate	7.7	2.2

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source: South Dakota Department of Health

**Figure 23: Cervix Uteri Cancer Stage at Diagnosis, South Dakota, 2018**



Source: South Dakota Department of Health

**Descriptive Epidemiology**

**Stage at Diagnosis:** Early stage of diagnosis provides the best opportunity for cure. In South Dakota, 43% of the cases reported were diagnosed at a localized stage. SEER reports that 44% of the cases diagnosed nationally were at a localized stage.

**Incidence:** In 2018, the incidence rate in South Dakota was 5.2 and the United States rate was 7.5. Both nationally and in South Dakota, cervical cancer was the third most common female genital tract malignancy. Invasive cervical cancer accounted for 0.4% of all cases reported and 0.9% of all females diagnosed with cancer in South Dakota in 2018. SEER incidence reports that 0.1% of cases were younger than 20 years of age.


**Mortality:** In 2018, the mortality rate in South Dakota was 2.0 for cancer of the cervix uteri. The United States rate was 2.2. The stage of disease at diagnosis affects the mortality rate. Cases diagnosed at a localized stage have a 92% survival rate according to the American Cancer Society. Nationally, when diagnosed at a distant stage, the percentage of survival drops to 17.6% at five years. In South Dakota, there were five cases in 2018 diagnosed at a distant stage.

**Risk and Associated Factors:** Almost all cervical cancers are caused by human papillomavirus (HPV). HPV is so common that most people get it at some time in their lives.

**Prevention and Early Detection:** The US Preventive Services Task Force (USPSTF) recommends screening for cervical cancer in women age 21 to 65 years with cytology (Pap test) every three years or, for women age 30 to 65 years who want to lengthen the screening interval, screening with a combination of cytology and human papillomavirus (HPV) testing every five years.


The HPV vaccine protects against the types of HPV that most often cause cervical, vaginal, and vulvar cancers. Many of these cancers could be prevented with vaccination. Vaccination is recommended for boys and girls ages 11 to 12 years but can be given at age 9 through 26.

**Figure 24: Cervix Uteri Cancer Number of Cases and Deaths by Age, South Dakota, 2018**



Source: South Dakota Department of Health


Half of the incidence of cervical cancer occurred in women under the age of 50.



Source: South Dakota Department of Health

**Figure 25: Cervix Uteri Cancer Cases and Deaths by Year, South Dakota, 2001 - 2018**

The incidence peak for female cervix uteri cancer was in 2014.



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health


**Figure 26: Cervix Uteri Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**

**Table 15: Colorectal Incidence and Mortality Summary, 2018**

Colorectal Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	<b>Total</b>	# Cases / Deaths Age-Adjusted Rate	<b>409</b> <b>38.5</b>	224 44.6	185 32.7	<b>168</b> <b>15.1</b>	92 18.1	76 12.4
	<b>White</b>	# Cases / Deaths Age-Adjusted Rate	<b>390</b> <b>39.7</b>	212 45.5	178 34.1	<b>153</b> <b>14.8</b>	84 17.8	69 12.1
	<b>American Indian</b>	# Cases / Deaths Age-Adjusted Rate	<b>17</b> <b>36.3</b>	10 51.1	7 24.3	<b>13</b> <b>31.1</b>	7 40.8	6 26.0
United States	<b>Total</b>	Age-Adjusted Rate	<b>36.8</b>	42.0	32.3	<b>13.1</b>	15.8	10.9
	<b>White</b>	Age-Adjusted Rate	<b>36.5</b>	41.4	32.2	<b>12.9</b>	15.5	10.8
	<b>American Indian</b>	Age-Adjusted Rate	<b>38.7</b>	39.3	38.0	<b>10.0</b>	11.7	8.5

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source: South Dakota Department of Health

**Figure 27: Colorectal Cancer Stage at Diagnosis, South Dakota, 2018**



Source: South Dakota Department of Health

**Descriptive Epidemiology**

**Stage at Diagnosis:** The patient’s prognosis is greatly influenced by the stage of disease at diagnosis. In 2018, 33% (138) of the cases of colorectal cancer were diagnosed at a localized stage. Localized is defined as when the disease is still confined to the colon. The remaining 246 invasive cases (59%) were diagnosed after the disease had spread beyond the colon. Of those 246 cases, 70 were diagnosed at a distant stage when the disease had spread further involving other organs. The SEER National Cancer Institute website states that the five-year survival rate for those who have a distant stage at diagnosis is 14.7% for the 2011-2017 time period.


**Incidence:** Colorectal cancer accounted for 8.5% of all cases reported in South Dakota in 2018. The median age at diagnosis was 70. There were 224 men and 185 women diagnosed with colorectal cancer in 2018 in South Dakota. Overall, colorectal cancer was the fourth most diagnosed cancer. When reviewed by gender, it was the third most diagnosed cancer with 8.8% of the cancers reported in males and 8.2% of the cancers reported in females.

**Mortality:** Overall incidence and mortality rates for colorectal cancer are decreasing. The overall five-year survival rate for 2011-2017 from SEER was 64.7% for men and women. In 2018, there were a total of 168 deaths that were attributed to colorectal cancer in South Dakota; about 16 more men than women. Of that number, 153 were white and thirteen were American Indian. The median age at death was 75. The SEER National Cancer Institute website states that the United States mortality rate was 13.1.


**Risk and Associated Factors:** Risk for colorectal cancer increases with age. Over 90% of cases occur in people who are 50 years old or older. Lifestyle factors that may increase risk include lack of regular physical activity, a diet low in fruit and vegetables, a low-fiber and high-fat diet, overweight and obesity, alcohol consumption, and tobacco use.

**Prevention and Early Detection:** The USPSTF recommends screening for colorectal cancer starting at age 50 and continuing until age 75. Recommended screening methods include high-sensitivity fecal occult blood test (FOBT), fecal immunochemical test (FIT), FIT-DNA, colonoscopy, CT colonography, & flexible sigmoidoscopy. The risks and benefits of these screening methods vary.

**Figure 28: Colorectal Cancer Number of Cases and Deaths by Age, South Dakota, 2018**




Source: South Dakota Department of Health



Source: South Dakota Department of Health

**Figure 29: Colorectal Cancer Cases and Deaths by Year, South Dakota, 2001 – 2018**

The incidence peak for colorectal cancer occurred in 2002. Incidences of colorectal cancer appear to escalate after age 50.



**Figure 30: Colorectal Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**

Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health


## CORPUS and UTERUS, NOS

**Table 16: Corpus and Uterus, NOS Incidence and Mortality Summary, 2018**

Corpus and Uterus, NOS Cancer			Incidence	Mortality
South Dakota	Total	# Cases / Deaths	143	21
		Age-Adjusted Rate	24.6	3.7
	White	# Cases / Deaths	131	21
		Age-Adjusted Rate	24.1	3.9
American Indian	# Cases / Deaths	11	0	
	Age-Adjusted Rate	38.3	0.0	
United States	Total	Age-Adjusted Rate	28.5	5.0
	White	Age-Adjusted Rate	28.6	4.6
	American Indian	Age-Adjusted Rate	26.2	2.2

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source: South Dakota Department of Health

**Figure 31: Corpus and Uterus, NOS Cancer Stage at Diagnosis, South Dakota, 2018**



Source: South Dakota Department of Health

### Descriptive Epidemiology

**Stage at Diagnosis:** Cancer in the uterus is treated surgically. Staging for these diseases is done following surgery, unless it is evident that the disease has progressed and advanced. Cases with evident advanced disease do not benefit from surgical procedures and are staged by physical examination. These cases are treated without operative staging. In South Dakota, during 2018, 79% of corpus uteri cases were diagnosed at a localized stage. Seven cases were diagnosed at a distant stage, less than in 2017.

**Incidence:** The uterine cervix is the small cylindrical neck that leads from the uterus, or womb, into the vagina. A knob of the cervix protrudes into the vagina and can be visualized on physical examination. It is lined with epithelial and stromal cells creating a site for epithelial, stromal, and mixed cell malignancies.


Endometrial carcinoma is one of the female genital cancers. It is ranked fourth among females reported with cancer in South Dakota in 2018. Cancer of the corpus uteri represented 6.3% of all the cancers diagnosed in South Dakota females in 2018. Endometrial cancer affects primarily postmenopausal women. The median age at diagnosis in South Dakota is 64 years of age.

**Mortality:** The mortality rate in South Dakota for the reporting period was 3.7 for deaths attributed to uterine cancer. In the United States the rate was 5.0. South Dakota had 21 female deaths attributed to cancer of the uterus in 2018. The stage of disease at diagnosis affects the mortality rate. Overall (all stages included), the five-year relative survival rate was 81.1% in the United States.

**Risk and Associated Factors:** Risk factors associated with corpus uteri cancer suggest that exposure to estrogen for long periods of time play a critical role. The use of exogenous estrogen replacement therapy accounted for a dramatic rise in the incidence of endometrial cancer in the United States in the 1970s. The use of combination estrogen-progesterone oral contraceptive pills confers protection against endometrial hyperplasia and subsequent development of cancer.


**Prevention and Early Detection:** Other factors associated with an increased risk of developing uterine cancer include obesity, a high-fat diet and prolonged exposure to the female hormone, estrogen. One pregnancy appears to lower the risk of uterine cancer by 50%.

**Figure 32: Corpus and Uterus, NOS Cancer Number of Cases and Deaths by Age, South Dakota, 2018**



Source: South Dakota Department of Health


In South Dakota, in 2018 the incidences were the highest in the 60-64 and 65-69 age groups.



Source: South Dakota Department of Health

**Figure 33: Corpus and Uterus, NOS Cancer Cases and Deaths by Year, South Dakota, 2001 - 2018**

The incidence peak for female corpus and uterus, NOS cancer was in 2017



**Figure 34: Corpus and Uterus, NOS Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**

Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health


## KIDNEY AND RENAL PELVIS

**Table 17: Kidney and Renal Pelvis Incidence and Mortality Summary, 2018**

Kidney and Renal Pelvis Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	174	110	64	43	29	14
		Age-Adjusted Rate	16.8	21.9	12.8	3.9	5.6	2.4
	White	# Cases / Deaths	161	101	60	39	26	13
Age-Adjusted Rate		16.9	21.7	13.2	3.8	5.3	2.5	
American Indian	# Cases / Deaths	11	8	3	3	3	0	
	Age-Adjusted Rate	22.8	47.2	9.1	6.5	15.0	0.0	
United States	Total	Age-Adjusted Rate	16.8	22.9	11.5	3.5	5.1	2.2
		Age-Adjusted Rate	17.5	23.8	12.0	3.7	5.3	2.3
	American Indian	Age-Adjusted Rate	22.9	31.8	16.0	4.1	6.3	2.4

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population.  
US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source: South Dakota Department of Health

**Figure 35: Kidney and Renal Pelvis Cancer Stage at Diagnosis, South Dakota, 2018**



Source: South Dakota Department of Health

### Descriptive Epidemiology

**Stage at Diagnosis:** As with all malignancies, early diagnosis is the key to a better prognosis and possible cure. In 2018, 67% of the cases were diagnosed at a localized stage; another 14% were diagnosed at a distant stage. Unfortunately, symptoms do not always reflect the stage of the disease. Blood in the urine is one of the symptoms that is frequently present at diagnosis. As with other cancers, renal cancer can spread through the bloodstream and/or lymphatic system. Survival rates associated with kidney cancer depend on how far the disease has progressed, the size of the tumor, and whether or not it has metastasized. The latest five-year survival rate for localized stage kidney cancer is 92.7%. The survival rate for distant stage is 13.9%.


**Incidence:** In 2018, the American Cancer Society estimated there would be 65,340 new cases of kidney cancer in the United States. This accounts for 3.8% of all reported malignancies in the United States. In South Dakota there were 180 reported cases of kidney cancer in 2018, representing 3.7% of all cancer cases with an age-adjusted rate of 17.4 per 100,000 persons. Kidney cancer develops most often in people over 40. There is no known cause of this disease. Doctors can seldom explain why one person develops kidney cancer and another does not. The median age at diagnosis is 64 in South Dakota and the United States.

**Mortality:** This cancer was the ninth leading cause of cancer death for South Dakota in 2018. In the United States, it was the 12th leading cause of death. The South Dakota median age of death was 70 years.

**Risk and Associated Factors:** Cigarette smoking increases the risk of developing kidney cancer. The risk seems to increase by the amount one smokes. Obesity is associated with risk, as are exposures to occupational substances such as aniline dyes, benzene, and naphthalene.


**Prevention and Early Detection:** The primary preventive measure is to stop smoking and maintain a healthy weight. It is difficult to diagnose kidney cancer until it becomes symptomatic. There are no known screenings recommended at this time.

**Figure 36: Kidney and Renal Pelvis Cancer Number of Cases and Deaths by Age, South Dakota, 2018**



Source: South Dakota Department of Health


**Figure 37: Kidney and Renal Pelvis Cancer Cases and Deaths by Year, South Dakota, 2001 - 2018**



Source: South Dakota Department of Health

The incidence peak for kidney and renal pelvis cancer occurred in 2016

**Figure 38: Kidney and Renal Pelvis Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.

## LEUKEMIA

**Table 18: Leukemia Incidence and Mortality Summary, 2018**

Leukemia			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	166	98	68	58	37	21
		Age-Adjusted Rate	16.1	19.9	12.9	5.1	7.4	3.3
	White	# Cases / Deaths	158	95	63	54	35	19
Age-Adjusted Rate		16.6	20.8	13.1	5.0	7.4	3.3	
American Indian	# Cases / Deaths	8	3	5	4	2	2	
	Age-Adjusted Rate	17.9	12.0	21.3	12.0	13.6	9.3	
United States	Total	Age-Adjusted Rate	14.7	18.9	11.4	6.0	8.1	4.5
		Age-Adjusted Rate	15.9	20.3	12.3	6.3	8.4	4.7
	American Indian	Age-Adjusted Rate	11.2	13.7	9.4	3.0	4.3	1.9

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source: South Dakota Department of Health

### Descriptive Epidemiology

**Stage at Diagnosis:** Leukemias are not staged because they may involve bone marrow throughout the body. Doctors classify them by type and subtype in an attempt to determine the prognosis and a recommended level of treatment. Chronic myelogenous leukemia is grouped by phases and chronic lymphocytic leukemia (CLL) uses a Rai classification. Leukemia is a type of cancer of the blood. It is defined by how quickly the disease progresses. Leukemia is either chronic (disease progresses slowly) or acute (progresses quickly).

**Incidence:** Leukemias are a diverse group of cancers and are subtyped by histology. Subtypes have different etiology, treatment, and prognosis. Leukemias accounted for 3.4% of the cancers reported in 2018 for South Dakota. The American Cancer Society estimated that there would be 190 new cases of leukemia in South Dakota during 2018 and 60,300 cases nationwide.

**Mortality:** Leukemia accounted for 3.5% of the cancer deaths in South Dakota in 2018. The subtype of acute myeloid leukemia was the most frequent cause of leukemia death. About 80% of the deaths associated with leukemia occurred at the age of 65 or older.

Leukemia is clinically and pathologically subdivided into a variety of large groups. The first division is between the acute and chronic forms.

**Chronic Leukemia:** Early in the disease process, the abnormal blood cells still have normal processes. Slowly, chronic leukemia does get worse. It causes symptoms as the number of abnormal cells in the blood rises. In South Dakota in 2018, there were 96 new cases of chronic leukemia.


**Acute Leukemia:** The blood cells are very abnormal. The blood cells cannot carry out their normal processes. The number of abnormal cells increases rapidly. Acute leukemia worsens quickly as do the symptoms. There were 63 new cases of acute leukemia in South Dakota in 2018.

These types of leukemia are further divided by the type of white blood cell that is affected.

**Risk and Associated Factors:** People who are exposed to very high levels of radiation are more likely to develop leukemia. Working with certain chemicals and exposure to high levels of benzene in the workplace can cause leukemia. Benzene is used widely in the chemical industry. Workers exposed to formaldehyde may also be at greater risk of leukemia. Chromosomal abnormalities, such as Down’s syndrome and certain other genetic diseases may increase the risk of leukemia.

**Prevention and Early Detection:** There are no early detection or prevention strategies. Often symptoms are the same as for many other health problems, thus early detection is difficult. Diagnosis is made using blood tests and bone marrow biopsies.


**Figure 39: Leukemia Number of Cases and Deaths by Age, South Dakota, 2018**



Source: South Dakota Department of Health

Leukemia is frequently diagnosed in children, but as with all malignancies it is a disease of the elderly. Deaths from leukemia increase after the age of 64.


**Figure 40: Leukemia Cases and Deaths by Year, South Dakota, 2001 - 2018**



The incidence peak for leukemia occurred in 2016.

Source: South Dakota Department of Health

**Figure 41: Leukemia Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health


## LUNG AND BRONCHUS

**Table 19: Lung and Bronchus Incidence and Mortality Summary, 2018**

Lung and Bronchus Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	<b>Total</b>	# Cases / Deaths Age-Adjusted Rate	<b>597</b> <b>51.9</b>	310 57.1	287 48.7	<b>395</b> <b>35.1</b>	212 41.3	183 31.0
	<b>White</b>	# Cases / Deaths Age-Adjusted Rate	<b>561</b> <b>51.5</b>	294 57.0	267 48.1	<b>363</b> <b>33.9</b>	196 39.8	167 30.3
	<b>American Indian</b>	# Cases / Deaths Age-Adjusted Rate	<b>32</b> <b>71.6</b>	15 76.7	17 70.7	<b>26</b> <b>64.3</b>	11 61.3	15 64.6
United States	<b>Total</b>	Age-Adjusted Rate	<b>50.8</b>	56.3	46.7	<b>34.8</b>	41.8	29.3
	<b>White</b>	Age-Adjusted Rate	<b>52.5</b>	56.8	49.4	<b>35.6</b>	42.1	30.6
	<b>American Indian</b>	Age-Adjusted Rate	<b>39.2</b>	39.9	38.8	<b>23.5</b>	27.2	20.5

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source: South Dakota Department of Health

**Figure 42: Lung and Bronchus Cancer Stage at Diagnosis, South Dakota, 2018**



Source: South Dakota Department of Health

### Descriptive Epidemiology

**Stage at Diagnosis:** The presentation of lung cancer is extremely variable and depends on local manifestations of the tumor, distant metastases or associated paraneoplastic syndromes. In 2018, 31% of lung cancer patients were diagnosed at a localized stage. The more advanced the stage, the poorer the prognosis is for the patient. In 2018, 257 (43%) cases were diagnosed when the disease had progressed beyond the lung and metastasized to a distant location. Approximately 62% of cases in 2018 were diagnosed after the disease had progressed beyond the lung to lymph nodes, regional areas, or distant sites, such as brain or bone.

**Incidence:** Lung cancer is a significant public health concern, with an estimated 234,030 new


cases in the United States in 2018. Despite the well documented link between tobacco product use and respiratory diseases, including cancer, the outcomes of such efforts to curb the use of tobacco products have been mixed. In South Dakota, there were 597 new invasive lung cancer cases diagnosed in 2018.

**Mortality:** There were 395 lung cancer deaths in South Dakota in 2018. Incidence and mortality rates have significantly increased during the last century. Lung cancer accounts for approximately 21.7% of all United States deaths attributed to cancer. In South Dakota, lung cancer accounts for 23.8% of deaths from cancer. Lung cancer is the leading cause of cancer deaths in both men and women.


**Risk and Associated Factors:** Cigarette smoking is the number one risk factor for lung cancer and is linked to approximately 90% of lung cancers. Other risk factors include secondhand smoke exposure, occupational or environmental exposures to substances such as radon, arsenic, benzene, and asbestos, a personal or family history of lung cancer, and previous radiation therapy to the chest.

**Prevention and Early Detection:** The USPSTF recommends annual screening for lung cancer with low-dose computed tomography (LDCT) in adults aged 50 to 80 years who have a 20 pack/ year smoking history and currently smoke or have quit within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.

**Figure 43: Lung and Bronchus Cancer Number of Cases and Deaths by Age, South Dakota, 2018**




Source: South Dakota Department of Health



Source: South Dakota Department of Health

**Figure 44: Lung and Bronchus Cancer Cases and Deaths by Year, South Dakota, 2001 - 2018**

The number of cases and deaths associated with lung and bronchus cancer remain constant.



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health

**Figure 45: Lung and Bronchus Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**


## MELANOMA OF THE SKIN

**Table 20: Melanoma of the Skin Incidence and Mortality Summary, 2018**

Melanoma of the Skin			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	304	173	131	34	25	9
		Age-Adjusted Rate	29.5	33.2	27.1	3.4	5.3	1.9
	White	# Cases / Deaths	303	172	131	33	25	8
		Age-Adjusted Rate	32.4	35.5	30.6	3.6	5.7	1.9
	American Indian	# Cases / Deaths	1	1	0	1	0	1
		Age-Adjusted Rate	2.1	4.4	0.0	1.4	0.0	2.7
United States	Total	Age-Adjusted Rate	22.7	28.9	18.1	2.1	3.0	1.4
		White	Age-Adjusted Rate	27.9	35.1	22.6	2.4	3.5
	American Indian	Age-Adjusted Rate	6.0	0.0	5.6	*	*	*

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) \*=rates suppressed. Source: South Dakota Department of Health

**Figure 46: Melanoma of the Skin Stage at Diagnosis, South Dakota, 2018**



Source: South Dakota Department of Health

### Descriptive Epidemiology

**Stage at Diagnosis:** Melanoma is staged by the depth of invasion and the extension of the lesion. In 2018, 43% of the melanoma of the skin cases reported for South Dakota were localized. Another 51% were staged as *in situ* disease. The survival rate for localized melanoma is 99.4%. For distant disease, the five-year survival rate is 29.8%.

**Incidence:** In the United States in 2018, the American Cancer Society estimated that there would be 91,270 new cases of melanoma of the skin. There are three forms of skin cancer: basal cell, squamous cell, and melanoma. Melanoma is by far the most dangerous form of skin cancer.

Melanoma is primarily a cancer of the white populations. In 2018, South Dakota's incidence rate was 29.5 and the United States had an incidence rate of 22.7.

**Mortality:** There were 34 deaths attributed to melanoma of the skin in South Dakota in 2018 with a mortality rate of 3.4 and the United States mortality rate was 2.3. The median age for death in South Dakota for this cancer was 70.5. The median age for the United States was 71.


**Risk and Associated Factors:** Certain factors are more likely to contribute to a higher risk:

- Lighter natural skin color
- Family history of skin cancer
- Personal history of skin cancer
- Exposure to the sun
- History of sunburns early in life
- Skin that burns, freckles, reddens easily
- Blue or green eyes, blond or red hair
- A large number of moles


**Prevention and Early Detection:** The best way to prevent skin cancer is to protect the skin from the sun. The CDC recommends five easy options for protection from sunburn:

- use sunglasses
- use sunscreen regularly
- stay in the shade
- cover skin
- wear a hat

**Figure 47: Melanoma of the Skin Number of Cases and Deaths by Age, South Dakota, 2018**




Source: South Dakota Department of Health



Source: South Dakota Department of Health

**Figure 48: Melanoma of the Skin Cases and Deaths by Year, South Dakota, 2001 - 2018**

The incidence count for melanoma cancers reached an all-time high in 2017.



**Figure 49: Melanoma of the Skin Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**

Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health

**Table 21: Myeloma Incidence and Mortality Summary, 2018**

Myeloma §			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	<b>87</b>	49	38	<b>39</b>	23	16
		Age-Adjusted Rate	<b>8.1</b>	9.4	7.0	<b>3.3</b>	4.8	2.4
	White	# Cases / Deaths	<b>80</b>	44	36	<b>39</b>	23	16
		Age-Adjusted Rate	<b>8.0</b>	9.1	7.2	<b>3.5</b>	5.0	2.5
	American Indian	# Cases / Deaths	<b>2</b>	1	1	<b>0</b>	0	0
		Age-Adjusted Rate	<b>3.4</b>	4.0	2.7	<b>0.0</b>	0.0	0.0
United States	Total	Age-Adjusted Rate	<b>7.3</b>	9.0	6.0	<b>3.1</b>	4.1	2.6
		White	Age-Adjusted Rate	<b>6.7</b>	8.5	5.2	<b>2.9</b>	3.8
	American Indian	Age-Adjusted Rate	<b>7.6</b>	8.6	6.5	<b>4.3</b>	3.9	3.1

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population.

§ can include NOS, multiple, plasma cell and solitary. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source:

South Dakota Department of Health

### Descriptive Epidemiology

**Stage at Diagnosis:** Stage of disease for myeloma is always distant per the SEER Summary Staging Manual.

**Incidence:** Myeloma is a systemic malignancy of plasma cells that is highly treatable, but rarely curable. It is potentially curable when it presents as a solitary plasmacytoma of the bone or as an extramedullary plasmacytoma. In South Dakota during 2018, myeloma accounted for 1.8% of total cancer cases reported. The 2018 South Dakota median age at diagnosis was 69 and the United States median age was 69. The national incidence rate is higher in men (9.0) than women (6.0). In South Dakota the incidence rate is also higher in men (9.4) than women (7.0). Myeloma is more common among the elderly. African Americans have approximately twice the incidence and mortality rates of whites.


**Mortality:** The median survival prior to the common use of chemotherapy was about seven months. After the introduction of chemotherapy, prognosis improved significantly with a median survival of 24 to 30 months and a 10-year survival of 3%. During 2018, there were 39 deaths attributed to myeloma in South Dakota. Twenty-three were male and 16 were female. The mortality rate for South Dakota was 3.3. The rate for men was 4.8 and 2.4 for women.

These rates compare to United States mortality rates of 3.1 overall, 4.1 for men and 2.6 for women.

**Risk and Associated Factors:** The etiology of myeloma is relatively unknown. There are many research studies evaluating the exposure of individuals with myeloma to various substances.


**Prevention and Early Detection:** There is no known test for screening for early detection. Some cases of myeloma progress very slowly, and they are referred to as smoldering or indolent myeloma. The presence of plasma cells and proteinuria do not automatically lead to myeloma, but it can be an early symptom. This disease is often asymptomatic in early stages of the disease. Myeloma is most often diagnosed clinically by radiological procedures and through cytology.

**Figure 50: Myeloma Number of Cases and Deaths by Age, South Dakota, 2018**



Source: South Dakota Department of Health


**Figure 51: Myeloma Cancer Cases and Deaths by Year, South Dakota, 2001 – 2018**



Source: South Dakota Department of Health

The incidence count for myeloma cancers took a sharp drop from 2004 to 2005 and 2013 to 2014, with an all-time high in 2018. Death counts were at an all-time high in 2016.

**Figure 52: Myeloma Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health


## NON-HODGKIN'S LYMPHOMA

**Table 22: Non-Hodgkin's Lymphoma Incidence and Mortality Summary, 2018**

Non-Hodgkin's Lymphoma			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	<b>Total</b>	# Cases / Deaths Age-Adjusted Rate	<b>191</b> <b>17.7</b>	105 20.3	86 15.3	<b>56</b> <b>5.0</b>	32 6.5	24 3.7
	<b>White</b>	# Cases / Deaths Age-Adjusted Rate	<b>183</b> <b>18.4</b>	102 21.4	81 15.7	<b>52</b> <b>4.8</b>	30 6.4	22 3.6
	<b>American Indian</b>	# Cases / Deaths Age-Adjusted Rate	<b>6</b> <b>9.2</b>	2 6.1	4 11.8	<b>3</b> <b>5.5</b>	2 7.3	1 3.9
United States	<b>Total</b>	Age-Adjusted Rate	<b>19.4</b>	23.3	16.3	<b>5.1</b>	6.8	3.8
	<b>White</b>	Age-Adjusted Rate	<b>20.5</b>	24.6	17.2	<b>5.4</b>	7.1	4.0
	<b>American Indian</b>	Age-Adjusted Rate	<b>11.9</b>	18.1	7.4	<b>3.0</b>	4.0	2.3

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source: South Dakota Department of Health

**Figure 53: Non-Hodgkin's Lymphoma Stage at Diagnosis, South Dakota, 2018**



Source: South Dakota Department of Health

### Descriptive Epidemiology

**Stage at Diagnosis:** Stage is based on where lymphoma cells are found (in the lymph or in other organs or tissues). The stage also depends on how many areas are involved. Localized stage only involves a single lymph node region or single extra lymphatic organ. When two or more lymph node regions are involved and the regions are on both sides of the diaphragm, the cancer is staged as distant. In 2018, 56% of the cases were diagnosed at a distant stage, the same as in 2017 but increased from 2011 when 47% were diagnosed at a distant stage.

**Incidence:** Non-Hodgkin's lymphoma is a cancer that originates in the lymphatic system, the disease-fighting network that spreads throughout the body. It develops in lymphocytes, a type of white blood cell. Non-Hodgkin's lymphoma is more than five times as


typical as the other general type of lymphoma, Hodgkin's disease. The incidence rate has been increasing in the United States since the 1970s. The incidence of non-Hodgkin's lymphoma in South Dakota was 191 cases in 2018. The 2018 median age at diagnosis in South Dakota was 68.

**Mortality:** There were 56 deaths reported in South Dakota that were attributed to non-Hodgkin's lymphoma. The median age at death for those whose death was attributed to non-Hodgkin's lymphoma in South Dakota was 79 years of age and 76 for the United States. Nationally, the five-year survival rate is 73.2% for non-Hodgkin's lymphoma.

**Risk and Associated Factors:** Age is a strong risk factor for this disease, with most cases occurring from age 60 and older. Some studies suggest that exposure to chemicals such as benzene and certain herbicides and insecticides may be linked to an increased risk. Some chemotherapy used to treat other cancers can increase the risk as well as patients having been treated with radiation. The risk is higher for those having been treated with both. Certain infections increase the risk, such as HIV, Epstein - Barr virus, H. pylori bacteria, and Hepatitis C virus.


**Prevention and Early Detection:** Non-Hodgkin's lymphoma may present with various symptoms. Symptoms may include signs resulting from local effects of cancer growth. Non-Hodgkin's lymphoma can also produce generalized symptoms, such as unexplained weight loss, fever, drenching night sweats, and severe itching.

**Figure 54: Non-Hodgkin's Lymphoma Number of Cases and Death by Age, South Dakota, 2018**



Source: South Dakota Department of Health


**Figure 55: Non-Hodgkin's Lymphoma Cases and Deaths by Year, South Dakota, 2001 – 2018**



Source: South Dakota Department of Health

The number of cases associated with non-Hodgkin's lymphoma cancer remains constant.

**Figure 56: Non-Hodgkin's Lymphoma Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**




Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health

**Table 23: Ovarian Incidence and Mortality Summary, 2018**

Ovarian Cancer			Incidence	Mortality
South Dakota	Total	# Cases / Deaths	51	48
		Age-Adjusted Rate	9.8	8.0
	White	# Cases / Deaths	48	46
		Age-Adjusted Rate	10.2	8.2
	American Indian	# Cases / Deaths	3	2
		Age-Adjusted Rate	9.5	6.1
United States	Total	Age-Adjusted Rate	10.3	6.3
		White	Age-Adjusted Rate	10.5
	American Indian	Age-Adjusted Rate	11.6	3.4

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source: South Dakota Department of Health

**Figure 57: Ovarian Cancer Stage at Diagnosis, South Dakota, 2018**



Source: South Dakota Department of Health

**Descriptive Epidemiology**

**Stage at Diagnosis:** Staging of ovarian cancer is done by a surgical procedure to remove as much of the cancer as possible. Surgical staging is of critical importance in the management of this disease. The morbidity associated with ovarian carcinoma is partially attributable to the fact that in the United States two-thirds of the patients present with advanced-stage disease at the time of diagnosis. In 2018, in South Dakota, 19 (37%) of the 51 cases were diagnosed at a distant stage.

**Incidence:** The incidence of ovarian cancer varies greatly. There were 51 invasive cases of ovarian cancer reported in 2018 in South Dakota. This accounted for 2.3% of the cancer cases diagnosed in 2018 for South Dakota women. The lifetime risk of a woman developing ovarian

cancer is 1.2%. There were twelve cases diagnosed at younger than 49 years of age. There were 16 cases diagnosed in the 60-69 age group. The median age at diagnosis in South Dakota was 62 and 63 in the United States.


**Mortality:** Doctors are using dramatic new therapies to fight ovarian cancer, extending the lives of women who five or 10 years ago would have died from the disease. Survival rates for the last several decades are only about 25% for those with advanced disease. Most ovarian cancer presents at advanced disease. Only 30.3% of those diagnosed at late stage survive five years. For those who are diagnosed early, before the disease spreads beyond the ovaries, the disease is 92.6% curable. In South Dakota, 48 patients died in 2018. The mortality rate was 8.0 for women in South Dakota and 6.3 in the United States.

**Risk and Associated Factors:** Women who have a higher risk for developing ovarian cancer have a family history of the disease, those who have used fertility drugs, those who had their first baby after age 30, and those over the age of 65.

**Prevention and Early Detection:** Factors that may reduce the risk of ovarian cancer are pregnancy lasting full term, use of oral contraceptives, breastfeeding, tubal ligation, hysterectomy, or removal of ovaries in women with inherited risks.


No early-stage screening tests have been proven for ovarian cancer and it can be challenging to detect until it has advanced. A combination of imaging and lab tests are the most useful diagnostic tools.

**Figure 58: Ovarian Cancer Number of Cases and Deaths by Age, South Dakota, 2018**



Source: South Dakota Department of Health


**Figure 59: Ovarian Cancer Cases and Deaths by Year, South Dakota, 2001 - 2018**



Source: South Dakota Department of Health

The incidence count for ovarian cancer peaked in 2013. The number of deaths rose to an all-time high in 2017.

**Figure 60: Ovarian Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**




Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health

**Table 24: Pancreas Incidence and Mortality Summary, 2018**

Pancreas Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	147	69	78	116	60	56
		Age-Adjusted Rate	12.9	12.4	13.3	10.0	10.7	9.1
	White	# Cases / Deaths	135	63	72	110	57	53
		Age-Adjusted Rate	12.7	12.1	13.3	10.1	10.8	9.2
	American Indian	# Cases / Deaths	11	5	6	6	3	3
		Age-Adjusted Rate	22.1	19.7	23.1	12.7	12.4	13.0
United States	Total	Age-Adjusted Rate	13.7	15.6	12.1	11.0	12.7	9.6
	White	Age-Adjusted Rate	13.7	15.8	11.9	11.0	12.7	9.5
	American Indian	Age-Adjusted Rate	10.8	13.6	8.6	6.0	6.6	5.5

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source: South Dakota Department of Health

**Figure 61: Pancreatic Cancer Stage at Diagnosis, South Dakota, 2018**



Source: South Dakota Department of Health

**Descriptive Epidemiology**

**Stage at Diagnosis:** Often pancreatic cancer is diagnosed late in the disease process. Patients who have local stage disease may be acceptable for resection. Only 10 to 20% of patients with pancreatic cancer are candidates for surgical resection. In South Dakota, 75% of new cases were diagnosed at a late stage (regional and distant) in 2018.

**Incidence:** The incidence of pancreatic cancer increases steadily with age. An estimated 55,440 new cases of pancreatic cancer were expected to be diagnosed in 2018 in the United States. The majority of the cases occurred in South Dakotans 65 years old or older. One hundred cases (68.0%) were diagnosed in 2018 in that age group. This cancer usually occurs more in males than in


females but in 2018 there were 69 men and 78 women diagnosed with pancreatic cancer in South Dakota. Nationally, there was a higher incidence rate in blacks of both genders. The median age at diagnosis was 69 years in South Dakota and 70 in the United States.

**Mortality:** The overall survival for cancer of the pancreas is poor. Studies reveal that the five-year survival rate is approximately 10.0%. More recently, prospective studies show survival improvement with postoperative chemotherapy. In 2018, there were 116 deaths and the median age at death was 70.5 in South Dakota and 72 in the United States.


**Risk and Associated Factors:** The exact causes of pancreatic cancer are unknown. Studies have found that certain factors increase a person's risk of developing pancreatic cancer. As one ages, the incidence of pancreatic cancer increases, especially after the age of 60. Cigarette smokers are two to three times more likely than nonsmokers to develop this cancer. Pancreatic cancer frequently occurs in those with diabetes. Also, African Americans are more likely than Asians, Hispanics, American Indians, or whites to have pancreatic cancer. The risk triples if the person's mother, father, sister, or brother had the disease. Also, a history of colon or ovarian cancer increases the risk. Some evidence shows that chronic pancreatitis may increase the risk.

**Prevention and Early Detection:** Currently, there are no known screenings for pancreatic cancer. Also, there is no specific prevention except to avoid smoking.

**Figure 62: Pancreatic Cancer Number of Cases and Deaths by Age, South Dakota, 2018**




Source: South Dakota Department of Health



**Figure 63: Pancreatic Cancer Cases and Deaths by Year, South Dakota, 2001 – 2018**

The number of cases and deaths associated with cancer of the pancreas reached an all-time high in 2016

Source: South Dakota Department of Health



**Figure 64: Pancreatic Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**

Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health


## PROSTATE

**Table 25: Prostate Incidence and Mortality Summary, 2018**

Prostate Cancer			Incidence	Mortality
South Dakota	<b>Total</b>	# Cases / Deaths Age-Adjusted Rate	690 119.5	92 19.1
	<b>White</b>	# Cases / Deaths Age-Adjusted Rate	640 117.6	83 18.1
	<b>American Indian</b>	# Cases / Deaths Age-Adjusted Rate	40 178.2	8 40.1
United States	<b>Total</b>	Age-Adjusted Rate	119.3	18.9
	<b>White</b>	Age-Adjusted Rate	114.1	17.7
	<b>American Indian</b>	Age-Adjusted Rate	54.8	12.8

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source: South Dakota Department of Health

**Figure 65: Prostate Cancer Stage at Diagnosis, South Dakota, 2018**



Source: South Dakota Department of Health

### Descriptive Epidemiology

**Stage at Diagnosis:** The greatest number of cases was diagnosed at an early stage. In 2018, 76% of the cases were diagnosed as localized (not extending outside the prostate). Frequently older cases may simply be monitored (watchful waiting) by their physician to assess the rate of growth; others may be given hormonal therapy. New treatments for prostate cancer include the da Vinci Robotic assisted prostatectomy, proton therapy, and brachytherapy radiation.

**Incidence:** Carcinoma of the prostate is predominately a tumor of older men. The median age at diagnosis in South Dakota is 67. Also, in South Dakota the incidence of prostate cancer begins to increase in the 60's age group. Nationwide, eight out of 10 men diagnosed with prostate cancer are over the age of 65. Prostate


cancer is the most diagnosed site of all cancers reported in the state in 2018.

**Mortality:** Prostate cancer was the third leading cancer death in men in South Dakota in 2018. Prostate cancer can be a slow progressing disease and can be cured or at least controlled in the early stages. The median age of death in South Dakota in 2018 was 82.5 years old. Many patients have co-morbid conditions and will die of other causes rather than prostate cancer.

**Risk and Associated Factors:** A number of risk factors for prostate cancer have been identified. Studies suggest that prostate cancer risk is increased two to five-fold in relatives of men with prostate cancer. Environmental factors, including exposure to heavy metals may increase risk. Smoking has also been indicated as a risk. Diets high in saturated fat intake may also contribute.


**Prevention and Early Detection:** The availability of the Prostate Specific Antigen (PSA) test as a diagnostic test coupled with increased awareness by the public of the disease has produced an increase in the number of new cases diagnosed each year in the United States. Disease detected by elevated PSA in the presence of a normal palpable gland is the most common presentation of prostate cancer. The American Cancer Society recommends the PSA and digital rectal exam should be offered annually beginning at the age of 50.

**Figure 66: Prostate Cancer Number of Cases and Deaths by Age, South Dakota, 2018**



Source: South Dakota Department of Health


**Figure 67: Prostate Cancer Cases and Deaths by Year, South Dakota, 2001 - 2018**



The number of deaths associated with prostate cancer remains constant

Source: South Dakota Department of Health

**Figure 68: Prostate Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota 2001 - 2018**




Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations. Source: South Dakota Department of Health

**Table 26: Stomach Incidence and Mortality Summary, 2018**

Stomach Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	<b>Total</b>	# Cases / Deaths Age-Adjusted Rate	<b>52</b> <b>4.7</b>	37 7.0	15 2.6	<b>20</b> <b>2.0</b>	11 2.4	9 1.5
	<b>White</b>	# Cases / Deaths Age-Adjusted Rate	<b>46</b> <b>4.6</b>	35 7.1	11 2.2	<b>16</b> <b>1.7</b>	8 1.9	8 1.5
	<b>American Indian</b>	# Cases / Deaths Age-Adjusted Rate	<b>4</b> <b>8.4</b>	0 0.0	4 13.6	<b>3</b> <b>8.1</b>	2 9.2	1 5.5
United States	<b>Total</b>	Age-Adjusted Rate	<b>7.0</b>	9.3	5.3	<b>2.8</b>	3.7	2.1
	<b>White</b>	Age-Adjusted Rate	<b>6.4</b>	8.5	4.6	<b>2.5</b>	3.3	1.8
	<b>American Indian</b>	Age-Adjusted Rate	<b>7.5</b>	9.1	6.1	<b>2.7</b>	3.3	2.3

Rates per 100,000 age-adjusted to 2000 US standard population and 2013 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) Source: South Dakota Department of Health

**Figure 69: Stomach Cancer Stage of Diagnosis, South Dakota, 2018**



Source: South Dakota Department of Health

**Descriptive Epidemiology**

**Stage at Diagnosis:** In 2018, data demonstrates that 15 (31%) cases were diagnosed at a localized stage. When a patient is diagnosed at an early stage, the prognosis is much better. Fourteen cases (29%) were diagnosed at a regional stage. There were 12 (25%) of the cases in South Dakota diagnosed at a distant stage. The prognosis for the distant stage is very poor. The stage is based on whether the tumor has invaded nearby tissues, where the cancer has spread, and if so, to what extent.

**Incidence:** Stomach cancer only accounted for approximately 1.0% of all cancers in South Dakota in 2018. Of the 47 cases diagnosed in 2018, 34 were male and 13 were female. It is

predominately a disease of men. Gastric (stomach) cancer is found more commonly in people between the ages of 50 and 70 years of age. The median age at diagnosis was 69 in South Dakota and 68 in the United States.


**Mortality:** Stomach cancer accounted for 1.2% of cancer deaths in South Dakota in 2018. The median age at death was 73 in South Dakota and 71 in the United States. The age-adjusted mortality rate was 2.5 for men and 1.3 in women in South Dakota. These rates are based on patients who died in 2018 in South Dakota. There were five American Indian stomach cancer deaths.

**Risk and Associated Factors:** Men have twice the risk of women for developing stomach cancer. In recent years, Helicobacter pylori bacteria have received considerable attention as a potential factor. Some researchers suspect this bacterium, which causes stomach inflammation and ulcers, may be an important stomach cancer risk factor. Individuals with pernicious anemia (a vitamin B-12-related disorder) and achlorhydria or gastric atrophy, both of which result in lower than normal amounts of gastric juices, may be at higher risk.

**Prevention and Early Detection:** Excessive salt intake has been identified as a possible risk factor for stomach cancer. Having a high intake of fresh fruits and vegetables may be associated with a decreased risk of stomach cancer. Studies have suggested that eating foods that contain **beta-carotene**<sup>1</sup> and **vitamin C**<sup>2</sup> may decrease the risk of stomach cancer.


<sup>1</sup><http://www.cancer.gov/Common/PopUps/popDefinition.aspx?id=45328&version=Patient&language=English>  
<sup>2</sup><http://www.cancer.gov/Common/PopUps/popDefinition.aspx?id=439435&version=Patient&language=English>

**Figure 70: Stomach Cancer Number of Cases and Deaths by Age, South Dakota, 2018**



Source: South Dakota Department of Health


**Figure 71: Stomach Cancer Cases and Deaths by Year, South Dakota, 2001 - 2018**



Source: South Dakota Department of Health

The incidence peak for stomach cancer was in 2010.

**Figure 72: Stomach Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**




Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health

**Table 27: Thyroid Incidence and Mortality Summary, 2018**

Thyroid Cancer			Incidence			Mortality		
			Total	Male	Female	Total	Male	Female
South Dakota	Total	# Cases / Deaths	138	36	102	7	5	2
		Age-Adjusted Rate	15.7	7.8	24.2	0.5	0.9	0.2
	White	# Cases / Deaths	122	32	90	7	5	2
		Age-Adjusted Rate	15.6	7.8	23.9	0.5	0.9	0.3
	American Indian	# Cases / Deaths	13	3	10	0	0	0
		Age-Adjusted Rate	23.8	11.1	36.6	0.0	0.0	0.0
United States	Total	Age-Adjusted Rate	14.9	7.8	21.8	0.5	0.5	0.5
	White	Age-Adjusted Rate	15.5	8.3	22.6	0.5	0.5	0.5
	American Indian	Age-Adjusted Rate	11.4	5.0	17.4	*	*	*

Rates per 100,000 age-adjusted to 2000 US standard population and 2018 SD estimated population. US rates [www.seer.cancer.gov](http://www.seer.cancer.gov) \*=rate suppressed. Source: South Dakota Department of Health

**Figure 73: Thyroid Cancer Stage at Diagnosis, South Dakota, 2018**



Source: South Dakota Department of Health

**Descriptive Epidemiology**

**Stage at Diagnosis:** In 2018, data demonstrates that 94 (68%) of cases were diagnosed at a localized stage. When a patient is diagnosed at an early stage, the prognosis is much better for a cure. There were 38 (27%) cases diagnosed at a regional stage. Four cases (3%) were diagnosed at a distant stage.

**Incidence:** The American Cancer Society estimated 53,990 thyroid cancer cases would be diagnosed in the United States in 2018. Thyroid cancer continues to account for approximately 2.9% of all cancers in South Dakota. Of the 138 cases diagnosed in 2018, 36 were male and 102 were female. The median age at diagnosis was 52.5 for South Dakota and 51 for the United States. Thyroid cancer is found more commonly

in people between the ages of 45 and 70 years of age, with 77% diagnosed before age 65. It is predominately a disease of females as the statistics for South Dakota confirm.


**Mortality:** South Dakota had only seven deaths attributed to thyroid cancer in 2018. Nationally, the five-year relative survival rates were 99.9% for localized, 98.3% for regional, and 53.3% for distant stage.

**Risk and Associated Factors:** Thyroid cancer accounted for only 2.9% of the cancer cases in South Dakota in 2018. Risk factors include being exposed to radiation to the head and neck in childhood. Other risk factors for the development of thyroid cancer include a history of goiter, family history of thyroid disease, and Asian race.


**Early Detection and Prevention:** Early detection of cancer of the thyroid is extremely important. There are currently no tests or screenings for early detection of thyroid cancer. Physical examinations may reveal a lump on the side of neck, hoarseness of the voice, and difficulty swallowing. Most cancerous thyroid tumors are slow growing and curable. Prompt attention to signs and symptoms is the best approach to early diagnosis of most thyroid cancers. Signs or symptoms include:

- A lump in the neck, sometimes growing rapidly
- Pain in the neck, sometimes going up to the ears
- Hoarseness
- Trouble swallowing
- Breathing problems (feeling as if one were breathing through a straw)
- A cough that persists and is not due to a cold

**Figure 74: Thyroid Cancer Number of Cases and Deaths by Age, South Dakota, 2018**




Source: South Dakota Department of Health



Source: South Dakota Department of Health

**Figure 75: Thyroid Cancer Cases and Deaths by Year, South Dakota, 2001 - 2018**

The incidence count for thyroid cancers peaked in 2018.



Rates per 100,000 age-adjusted to 2000 US standard population and SD estimated populations.  
Source: South Dakota Department of Health

**Figure 76: Thyroid Cancer Age-Adjusted Rates, Cases, and Deaths by Year, South Dakota, 2001 - 2018**

## XI: APPENDICES

### Appendix A: 2000 United States Standard Million Population

Age Group	Number in Group
Total	1,000,000
<5	69,135
5-9	72,533
10-14	73,032
15-19	72,169
20-24	66,478
25-29	64,529
30-34	71,044
35-39	80,762
40-44	81,851
45-49	72,118
50-54	62,716
55-59	48,454
60-64	38,793
65-69	34,264
70-74	31,773
75-79	26,999
80-84	17,842
85+	15,508

### Appendix B: 2009-2018 South Dakota Estimated Population

Year	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
Total	812,383	814,180	824,082	833,354	844,877	853,175	858,469	865,454	869,666	882,235
<5	59,640	59,621	59,591	59,202	59,957	60,610	61,244	61,369	61,759	62,132
5-9	53,496	55,531	56,389	58,010	59,832	60,148	60,266	60,477	60,372	60,762
10-14	52,576	53,960	54,096	53,956	54,899	55,807	56,525	57,842	59,303	60,882
15-19	58,571	57,628	57,916	57,439	57,419	57,823	56,581	56,838	56,675	57,674
20-24	62,919	57,596	58,178	59,174	60,849	61,697	61,382	61,366	59,550	59,585
25-29	56,270	55,570	56,020	56,397	56,103	55,859	55,656	55,852	57,257	58,754
30-34	46,540	49,859	52,216	53,875	55,411	56,075	56,436	56,414	56,348	57,341
35-39	45,707	45,766	45,524	46,326	47,452	49,007	50,813	52,857	54,536	56,007
40-44	47,626	47,346	47,375	47,570	47,255	46,709	46,136	45,735	46,458	47,902
45-49	57,845	57,519	54,849	52,681	50,182	48,446	47,650	47,699	47,425	47,241
50-54	57,850	59,399	59,960	60,037	59,699	58,976	57,094	54,496	52,109	49,840
55-59	52,996	54,231	56,261	57,577	58,434	58,768	59,517	59,850	59,476	59,311
60-64	42,615	43,573	47,054	48,927	51,222	53,027	54,749	55,854	56,774	57,950
65-69	32,303	31,944	33,144	35,738	37,753	40,473	43,381	46,846	48,277	50,160
70-74	25,236	25,683	25,882	26,396	27,834	28,875	29,750	30,618	32,952	34,890
75-79	21,866	21,724	21,660	21,766	21,939	22,209	22,305	22,243	22,463	23,666
80-84	18,122	18,004	18,102	18,112	18,017	17,795	17,616	17,582	17,151	17,147
85+	20,205	19,226	19,865	20,171	20,620	20,871	21,368	21,516	20,781	20,991

## Appendix C: Race in South Dakota by County, 2018 Estimated Population

	Total	White	Black	American Indian	Asian	Other
South Dakota	882,235	744,733 84%	21,000 2%	79,412 9%	15,093 2%	21,997 2%
Aurora	2,801	2,633 94%	21 1%	76 3%	20 1%	51 2%
Beadle	18,883	15,732 83%	262 1%	284 2%	2149 11%	456 2%
Bennett	3,468	1,169 34%	18 1%	2,077 60%	15 0%	189 5%
Bon Homme	6,980	6,167 88%	95 1%	596 9%	18 0%	104 1%
Brookings	35,232	32,602 93%	579 2%	419 1%	1031 3%	601 2%
Brown	39,316	34,800 89%	1022 3%	1,353 3%	1225 3%	916 2%
Brule	5,229	4,480 86%	29 1%	516 10%	20 0%	184 4%
Buffalo	2,036	322 16%	16 1%	1,645 81%	1 0%	52 3%
Butte	10,222	9,580 94%	62 1%	249 2%	46 0%	285 3%
Campbell	1,377	1,341 97%	4 0%	14 1%	6 0%	12 1%
Charles Mix	9,338	5,949 64%	45 0%	2,997 32%	25 0%	322 3%
Clark	3,739	3,585 96%	75 2%	18 0%	24 1%	37 1%
Clay	14,041	12,462 89%	239 2%	571 4%	405 3%	364 3%
Codington	28,015	26,426 94%	175 1%	762 3%	214 1%	438 2%
Corson	4,165	1,252 30%	20 0%	2,731 66%	22 1%	140 3%
Custer	8,726	8,105 93%	50 1%	321 4%	56 1%	194 2%
Davison	19,790	18,411 93%	181 1%	663 3%	127 1%	408 2%
Day	5,505	4,814 87%	21 0%	514 9%	47 1%	109 2%
Deuel	4,337	4,216 97%	38 1%	24 1%	11 0%	48 1%
Dewey	5,904	1,236 21%	30 1%	4,370 74%	12 0%	256 4%
Douglas	2,935	2,795 95%	18 1%	73 2%	5 0%	44 1%
Edmunds	3,875	3,732 96%	12 0%	45 1%	46 1%	40 1%
Fall River	6,758	5,910 87%	87 1%	446 7%	94 1%	221 3%
Faulk	2,330	2,277 98%	12 1%	14 1%	15 1%	12 1%
Grant	7,147	6,868 96%	48 1%	107 1%	26 0%	98 1%
Gregory	4,212	3,717 88%	17 0%	328 8%	19 0%	131 3%
Haakon	1,918	1,787 93%	5 0%	50 3%	5 0%	71 4%
Hamlin	6,111	5,953 97%	31 1%	52 1%	22 0%	53 1%
Hand	3,262	3,192 98%	4 0%	18 1%	13 0%	35 1%
Hanson	3,376	3,296 98%	17 1%	14 0%	14 0%	35 1%
Harding	1,249	1,180 94%	12 1%	29 2%	3 0%	25 2%
Hughes	17,650	14,760 84%	135 1%	2,102 12%	137 1%	516 3%
Hutchinson	7,380	7,103 96%	73 1%	102 1%	19 0%	83 1%
Hyde	1,282	1,130 88%	7 1%	110 9%	3 0%	32 2%
Jackson	3,307	1,362 41%	40 1%	1,730 52%	5 0%	170 5%
Jerauld	2,043	1,989 97%	8 0%	14 1%	4 0%	28 1%
Jones	928	842 91%	3 0%	42 5%	0 0%	41 4%
Kingsbury	4,919	4,734 96%	23 0%	54 1%	31 1%	77 2%
Lake	13,057	12,409 95%	148 1%	162 1%	133 1%	205 2%
Lawrence	25,741	24,024 93%	214 1%	607 2%	382 1%	514 2%
Lincoln	58,807	55,642 95%	899 2%	423 1%	797 1%	1046 2%
Lyman	3,821	2,153 56%	28 1%	1,495 39%	12 0%	133 3%
McCook	5,546	5,337 96%	48 1%	74 1%	14 0%	73 1%
McPherson	2,407	2,349 98%	18 1%	6 0%	9 0%	25 1%
Marshall	5,112	4,472 87%	36 1%	487 10%	10 0%	107 2%
Meade	28,294	25,611 91%	577 2%	849 3%	322 1%	935 3%
Mellette	2,042	797 39%	6 0%	1,114 55%	5 0%	120 6%
Miner	2,213	2,130 96%	23 1%	16 1%	10 0%	34 2%
Minnehaha	192,876	165,092 86%	12,664 7%	5,364 3%	4,751 2%	5,005 3%
Moody	6,579	5,178 79%	96 1%	890 14%	200 3%	215 3%
Oglala Lakota	14,309	762 5%	44 0%	13,238 93%	23 0%	242 2%
Pennington	111,729	92,960 83%	1626 1%	11305 10%	1454 1%	4384 4%
Perkins	2,922	2,795 96%	15 1%	58 2%	11 0%	43 1%
Potter	2,207	2,085 94%	11 0%	53 2%	17 1%	41 2%
Roberts	10,447	5,998 57%	77 1%	4008 38%	25 0%	339 3%
Sanborn	2,429	2,365 97%	5 0%	11 0%	7 0%	41 2%
Spink	6,495	6,203 96%	44 1%	175 3%	6 0%	67 1%
Stanley	3,022	2,707 90%	21 1%	208 7%	8 0%	78 3%
Sully	1,392	1,326 95%	6 0%	34 2%	2 0%	24 2%
Todd	10,283	894 9%	60 1%	8,814 86%	269 3%	246 2%
Tripp	5,478	4,475 82%	29 1%	803 15%	21 0%	150 3%
Turner	8,424	8,168 97%	49 1%	94 1%	18 0%	95 1%
Union	15,619	14,793 95%	196 1%	131 1%	246 2%	253 2%
Walworth	5,587	4,393 79%	26 0%	800 14%	187 3%	181 3%
Yankton	22,869	21,024 92%	487 2%	733 3%	207 1%	418 2%
Ziebach	2,742	682 25%	13 0%	1,930 70%	12 0%	105 4%

US Census Bureau Estimated Race Population

### Appendix D: SEER Incidence Site Analysis Categories

Site Group	ICD-O-3 Site	ICD-O-3 Histology (Type)	Recode	
<b>Oral Cavity and Pharynx</b>				
Lip	C000-C009	excluding 9050-9055, 9140, 9590-9992	20010	
Tongue	C019-C029		20020	
Salivary Gland	C079-C089		20030	
Floor of Mouth	C040-C049		20040	
Gum and Other Mouth	C030-C039, C050-C059, C060-C069		20050	
Nasopharynx	C110-C119		20060	
Tonsil	C090-C099		20070	
Oropharynx	C100-C109		20080	
Hypopharynx	C129, C130-C139		20090	
Other Oral Cavity and Pharynx	C140, C142, C148		20100	
<b>Digestive System</b>				
Esophagus	C150-C159	excluding 9050-9055, 9140, 9590-9992	21010	
Stomach	C160-C169		21020	
Small Intestine	C170-C179		21030	
<b>Colon and Rectum</b>				
Colon excluding Rectum				
Cecum	C180	excluding 9050-9055, 9140, 9590-9992	21041	
Appendix	C181		21042	
Ascending Colon	C182		21043	
Hepatic Flexure	C183		21044	
Transverse Colon	C184		21045	
Splenic Flexure	C185		21046	
Descending Colon	C186		21047	
Sigmoid Colon	C187		21048	
Large Intestine, NOS	C188-C189, C260		21049	
<b>Rectum and Rectosigmoid Junction</b>				
Rectosigmoid Junction	C199	excluding 9050-9055, 9140, 9590-9992	21051	
Rectum	C209		21052	
Anus, Anal Canal and Anorectum	C210-C212, C218		21060	
<b>Liver and Intrahepatic Bile Duct</b>				
Liver	C220	excluding 9050-9055, 9140, 9590-9992	21071	
Intrahepatic Bile Duct	C221		21072	
Gallbladder	C239		21080	
Other Biliary	C240-C249		21090	
Pancreas	C250-C259		21100	
Retroperitoneum	C480		21110	
Peritoneum, Omentum and Mesentery	C481-C482		21120	
Other Digestive Organs	C268-C269, C488		21130	
<b>Respiratory System</b>				
Nose, Nasal Cavity and Middle	C300-C301, C310-C319		excluding 9050-9055, 9140, 9590-9992	22010
Larynx	C320-C329	22020		
Lung and Bronchus	C340-C349	22030		
Pleura	C384	22050		
Trachea, Mediastinum and Other Respiratory Organs	C339, C381-C383, C388, C390, C398, C399	22060		

**Appendix D: SEER Incidence Site Analysis Categories (continued)**

Site Group	ICD-O-3 Site	ICD-O-3 Histology (Type)	Recode
<b>Bones and Joints</b>	<b>C400-C419</b>	<b>excluding 9050-9055, 9140, 9590-9992</b>	<b>23000</b>
<b>Soft Tissue including Heart</b>	<b>C380, C470-C479, C490-C499</b>	<b>excluding 9050-9055, 9140, 9590-9992</b>	<b>24000</b>
<b>Skin excluding Basal and Squamous</b>			
Melanoma of the Skin	C440-C449	8720-8790	25010
Other Non-Epithelial Skin	C440-C449	excluding 8000-8005, 8010-8046, 8050-8084, 8090-8110, 8720-8790, 9050-9055, 9140, 9590-9992	25020
<b>Breast</b>	<b>C500-C509</b>	<b>excluding 9050-9055, 9140, 9590-9992</b>	<b>26000</b>
<b>Female Genital System</b>			
Cervix Uteri	C530-C539	excluding 9050-9055, 9140, 9590-9992	27010
Corpus and Uterus, NOS			
Corpus Uteri	C540-C549	excluding 9050-9055, 9140, 9590-9992	27020
Uterus, NOS	C559		27030
Ovary	C569		27040
Vagina	C529		27050
Vulva	C510-C519		27060
Other Female Genital Organs	C570-C579, C589		27070
<b>Male Genital System</b>			
Prostate	C619	excluding 9050-9055, 9140, 9590-9992	28010
Testis	C620-C629		28020
Penis	C600-C609		28030
Other Male Genital Organs	C630-C639		28040
<b>Urinary System</b>			
Urinary Bladder	C670-C679	excluding 9050-9055, 9140, 9590-9992	29010
Kidney and Renal Pelvis	C649, C659		29020
Ureter	C669		29030
Other Urinary Organs	C680-C689		29040
<b>Eye and Orbit</b>	<b>C690-C699</b>	<b>excluding 9050-9055, 9140, 9590-9992</b>	<b>30000</b>
<b>Brain and Other Nervous System</b>			
Brain	C710-C719	excluding 9050-9055, 9140, 9530-9539, 9590-9992	31010
Cranial Nerves Other Nervous System	C710-C719	9530-9539	31040
	C700-C709, C720-C729	excluding 9050-9055, 9140, 9590-9992	
<b>Endocrine System</b>			
Thyroid	C739	excluding 9050-9055, 9140, 9590-9992	32010
Other Endocrine including Thymus	C379, C740-C749, C750-C759		32020

**Appendix D: SEER Incidence Site Analysis Categories (continued)**

Site Group	ICD-O-3 Site	ICD-O-3 Histology (Type)	Recode
<b>Lymphoma</b>			
Hodgkin's Lymphoma			
Hodgkin's - Nodal	C024,C098-C099,C111, C142,C379,C422,C770-C779	9650-9667	33011
Hodgkin's - Extranodal	All other sites		33012
<b>Non-Hodgkin's Lymphoma</b>			
NHL - Nodal	C024, C098, C099, C111, C142,C379, C422, C770-C779	9590-9597, 9670-9671, 9673, 9675, 9678-9680, 9684, 9687-9691, 9695, 9698-9702, 9705, 9708-9709, 9712, 9714-9719, 9724-9729, 9735, 9737-9738, 9811-9818, 9823, 9827, 9837	33041
NHL - Extranodal	All sites except C024, C098-C099, C111, C142, C379, C422, C770-C779	9590-9597, 9670-9671, 9673, 9675, 9678-9680, 9684, 9687, 9688, 9689-9691, 9695, 9698-9702,9705, 9708-9709, 9712, 9714-9719, 9724-9729, 9735, 9737,9738	33042
	All sites except C024, C098-C099, C111, C142, C379, C420-C422, C424, C770-C779	9811-9818, 9823, 9827, 9837	
<b>Myeloma</b>		<b>9731-9732, 9734</b>	<b>34000</b>
<b>Leukemia</b>			
Lymphocytic Leukemia			
Acute Lymphocytic Leukemia		9826, 9835-9836	35011
	C420, C421, C424	9811-9818, 9837	
Chronic Lymphocytic Leukemia	C420, C421, C424	9823	35012
Other Lymphocytic Leukemia		9820, 9832-9834, 9940	35013
<b>Myeloid and Monocytic Leukemia</b>			
Acute Myeloid Leukemia		9840, 9861, 9865-9867, 9869, 9871-9874, 9895-9897, 9898, 9910-9911, 9920	35021
Acute Monocytic Leukemia		9891	35031
Chronic Myeloid Leukemia		9863, 9875-9876, 9945-9946	35022
Other Myeloid/Monocytic Leukemia		9860, 9930	35023
<b>Other Leukemia</b>			
Other Acute Leukemia		9801, 9805-9809, 9931	35041
Aleukemic, subleukemic and NOS		9733, 9742, 9800, 9831, 9870, 9948, 9963-9964	35043
	C420, C421, C424	9827	
<b>Mesothelioma +</b>		<b>9050-9055</b>	<b>36010</b>
<b>Kaposi Sarcoma +</b>		<b>9140</b>	<b>36020</b>
<b>Miscellaneous</b>		<b>9740-9741, 9750-9769, 9950, 9960-9962, 9965-9967, 9970-9971, 9975, 9980, 9982-9987, 9989, 9991-9992</b>	<b>37000</b>
	<b>C760-C768, C809</b>	<b>Excluding 9050-9055, 9140, 9590-9992</b>	
	<b>C420-C424</b> <b>C770-C779</b>		
<b>Invalid</b>	<b>Site or histology code not within valid range or site code not found in this table.</b>		<b>99999</b>

Source: <http://seer.cancer.gov/siterecode>

+ The Site Recode variable can be created with or without Mesothelioma (9050-9055) and Kaposi Sarcoma (9140) as separate groupings. The table above documents both possibilities.

## **Appendix E: SEER Cancer Cause of Death Analysis Categories**

Cancer Causes of Death	ICD-10
<b>All Malignant Cancers</b>	<b>C00-C97</b>
<b>Oral Cavity and Pharynx</b>	
Lip	C00
Tongue	C01-C02
Salivary Gland	C07-C08
Floor of Mouth	C04
Gum and Other Mouth	C03, C05-C06
Nasopharynx	C11
Tonsil	C09
Oropharynx	C10
Hypopharynx	C12-C13
Other Oral Cavity and Pharynx	C14
<b>Digestive System</b>	
Esophagus	C15
Stomach	C16
Small Intestine	C17
Colon and Rectum	
Colon Excluding Rectum	C18, C26.0
Rectum and Rectosigmoid Junction	C19-C20
Anus, Anal Canal and Anorectum	C21
<b>Liver and Intrahepatic Bile Duct</b>	
Liver	C22.0, C22.2-C22.4, C22.7, C22.9
Intrahepatic Bile Duct	C22.1
Gallbladder	C23
Other Biliary	C24
Pancreas	C25
Retroperitoneum	C48.0
Peritoneum, Omentum and Mesentery	C45.1+, C48.1-C48.2
Other Digestive Organs	C26.8-C26.9, C48.8
<b>Respiratory System</b>	
Nose, Nasal Cavity and Middle Ear	C30-C31
Larynx	C32
Lung and Bronchus	C34
Pleura	C38.4, C45.0+
Trachea, Mediastinum and Other Respiratory Organs	C33, C38.1-C38.3, C38.8, C39
<b>Bones and Joints</b>	C40-C41
<b>Soft Tissue including Heart</b>	C47, C49, C38.0, C45.2+
<b>Skin excluding Basal and Squamous</b>	
Melanoma of the Skin	C43
Other Non-Epithelial Skin	C44, C46+
<b>Breast</b>	C50
<b>Female Genital System</b>	
Cervix Uteri	C53
Corpus and Uterus, NOS	
Corpus Uteri	C54
Uterus, NOS	C55
Ovary	C56
Vagina	C52
Vulva	C51
Other Female Genital Organs	C57-C58

**Appendix E: SEER Cancer Cause of Death Analysis Categories (Continued)**

Cancer Causes of Death	ICD-10
<b>Male Genital System</b>	
Prostate	C61
Testis	C62
Penis	C60
Other Male Genital Organs	C63
<b>Urinary System</b>	
Bladder	C67
Kidney and Renal Pelvis	C64-C65
Ureter	C66
Other Urinary Organs	C68
<b>Eye and Orbit</b>	C69
<b>Brain and Other Nervous System</b>	C70, C71, C72
<b>Endocrine System</b>	
Thyroid	C73
Other Endocrine Including Thymus	C37, C74-C75
<b>Lymphoma</b>	
Hodgkin's Lymphoma	C81
Non-Hodgkin's Lymphoma	C82-C85, C96.3
<b>Myeloma</b>	C90.0, C90.2
<b>Leukemia</b>	
Lymphocytic Leukemia	
Acute Lymphocytic Leukemia	C91.0
Chronic Lymphocytic Leukemia	C91.1
Other Lymphocytic Leukemia	C91.2-C91.4, C91.7, C91.9
Myeloid and Monocytic Leukemia	
Acute Myeloid	C92.0, C92.4-C92.5, C94.0, C94.2
Acute Monocytic Leukemia	C93.0
Chronic Myeloid Leukemia	C92.1
Other Myeloid/Monocytic Leukemia	C92.2-C92.3, C92.7, C92.9, C93.1-C93.2, C93.7, C93.9
Other Acute Leukemia	C94.4, C94.5, C95.0
Aleukemic, Sub leukemic and NOS	C90.1, C91.5, C94.1, C94.3, C94.7, C95.1, C95.2, C95.7, C95.9
<b>Mesothelioma (ICD-10 only)+</b>	C45+
<b>Kaposi Sarcoma (ICD-10 only)+</b>	C46+
<b>Miscellaneous Malignant Cancer</b>	C26.1, C45.7+, C45.9+, C76-C80, C88, C96.0-C96.2, C96.7, C96.9, C97

Source: <http://seer.cancer.gov/codrecode>

## REFERENCES

---

- Bast et Al. Cancer Edition 5. American Cancer Society, 2000.
- Beahrs, O.H. et al, AJCC Staging Manual, 5<sup>th</sup> Edition. American Joint Committee on Cancer, Lippincott, Williams and Wilkins. 1995.
- Cancer Facts and Figures, 2018. Atlanta. American Cancer Society. 2018.
- Cancer Progress Report, US Department of Health and Human Services, National Cancer Institute. 2005.
- Devita, Vincent et al. CANCER Principles and Practice of Oncology. Lippincott. Williams & Wilkins. 2001.
- Harvard Center for Cancer Prevention. Harvard Report on Cancer Prevention Volume 1: Causes of Human Cancer, 1996:7 (S1): 7-15.
- Howlander N, Noone AM, Krapcho M, Garshell J, Miller D, Altekruse SF, Kosary CL, Yu M, Ruhl J, Tatalovich Z, Mariotto A, Lewis DR, Chen HS, Feuer EJ, Cronin KA (eds). SEER Cancer Statistics Review, 1975-2011, National Cancer Institute. Bethesda, MD, [http://seer.cancer.gov/csr/1975\\_2011/](http://seer.cancer.gov/csr/1975_2011/), based on November 2013 SEER data submission, posted to the SEER web site, April 2018.
- Jemal A, Simard EP, Dorell C, Noone AM, Markowitz LE, Kohler B, Ehemann C, Saraiya M, Bandi P, Saslow D, Cronin KA, Watson M, Schiffman M, Henley SJ, Schymura MJ, Anderson RN, Yankey D, and Edwards BK. Annual Report to the Nation on the Status of Cancer, 1975-2009, Featuring the Burden and Trends in HPV-Associated Cancers and HPV Vaccination Coverage Levels. *J Natl Ca Inst* 2013 Feb;105(3). Epub 2013 Jan 7. Lenhard, Raymond et al. Clinical Oncology. The American Cancer Society. Bethesda, MD. 2001.
- Nascar, Philip and Harris Pastides. Fundamentals of Cancer Epidemiology. Aspen Publications, Maryland. 2001.
- Schottenfeld, David and Joseph Fraumenis, Cancer Epidemiology and Prevention. 2<sup>nd</sup> Edition. Oxford University Press. 1996.
- Surveillance, Epidemiology, and End Results (SEER) Program of the National Cancer Institute, Cancer Query Systems, <http://seer.cancer.gov/canques/>

This document is available online at  
<http://getscreened.sd.gov/documents/Cancer2018.pdf>

A limited number were printed and publication, funded by the South Dakota Cancer Registry with a grant from the Centers for Disease Control and Prevention – grant number DP006293. For more information or additional copies (if available), contact the South Dakota Department of Health at 605.773.3361

Per SDCL 5-18D-15: 75 copies of this publication have been printed on recycled paper by the South Dakota Department of Health at a cost of \$        each.